

SPECIAL LOCAL NEED 24(C)LABELING

VYDATE® L
INSECTICIDE/NEMATICIDE
BULB ONIONS
STATE OF COLORADO

RESTRICTED USE PESTICIDE

Due to Acute Toxicity And Toxicity to Birds and Mammals.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF COLORADO

VYDATE® L INSECTICIDE/NEMATICIDE

EPA Reg. No. 352 -372

FOR USE ON BULB ONIONS IN THE STATE OF COLORADO

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

VYDATE® L Insecticide/Nematicide is recommended for use on bulb onions for suppression of onion thrips and western flower thrips in the State of Colorado.

FOLIAR APPLICATION

Apply by ground, air or overhead sprinkler chemigation at 2 to 4 pts/Acre (minimum 5 gal water/acre by air). Apply before insects reach the economic thresholds. Repeat applications at 14 day intervals.

Use of a nonionic surfactant at 0.25% v/v (1 qt. per 100 gal spray solution) or 0.5% (2 qts. per 100 gal. spray solution) under arid conditions will improve performance. Acidify the spray solution to a pH lower than 6.0.

DRIP APPLICATION

Apply 2 to 4 pts/acre through the drip irrigation system. Apply before insects reach the economic threshold. Repeat applications at 7 to 14 day intervals. Apply in enough water to thoroughly wet the root zone. Always buffer the VYDATE® L injection solution to a pH of 5 or lower.

AT PLANTING AND SIDE DRESS APPLICATION

Apply up to 1 gallon per acre with or without liquid fertilizer. Apply before insects reach economic threshold.

Inject into soil 2-4 inches below onion plants. Repeat applications can be made by soil injection or at 2-4 pints per acre via foliar spray (normally at 14-day intervals).

Irrigate with enough water to thoroughly wet the root zone once application is complete.

Acidify the solution to a pH lower than 6.0.

NOTE: Do not harvest tops of treated bulb onions.

Do not apply within 14 days of harvest.

Do not use on green onions.

Do not apply more than 18 pts (2 1/4 gals) VYDATE® L per acre per season.

Do not apply more than 8 applications of VYDATE® L per acre per growing season.

VYDATE® L should not be applied at the same time that a drip/irrigation line clean out product is being used as performance may be reduced. Soil applications must be incorporated into the soil by water or mechanical means.

Refer to the VYDATE® L package label for specific precautions regarding additional chemigation use and required safety devices.

IMPORTANT

BEFORE USING VYDATE® L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners