



**Galigan® 2E Herbicide**  
EPA REG NO. 66222-28

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **GALIGAN® 2E HERBICIDE** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

“Label” as used in this supplemental labeling refers to the label booklet for Galigan 2E Herbicide and this supplement.

---

**APRICOTS, NECTARINES, OLIVES, PEACHES, PLUMS, AND PRUNES**  
**CALIFORNIA ONLY**

---

### **KEEP OUT OF REACH OF CHILDREN**

### **WARNING**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

---

### **DIRECTIONS FOR USE**

---

#### **Nondormant Application to Apricots, Nectarines, Olives, Peaches, Plums, and Prunes in California.**

##### **Product Information**

Galigan 2E herbicide provides effective postemergence control of cheeseweed (malva), fleabane, and marestalk (horseweed) young broadleaf weed seedlings in non-dormant apricots, nectarines, olives, peaches, plums, and prunes. For enhanced postemergence activity against these target weeds as well as other weed species, tank mix Galigan 2E herbicide with either paraquat (Parazone 3SL) or glyphosate (Glyphogan) to increase the spectrum of weed control by either of these tank mix partners. Compatibility of each mixture must be established before tank mixing and application must be applied by ground equipment. Follow all precautions and restrictions on the labeling of the products to be tank mixed.

##### **Dosage**

Use Galigan 2E herbicide for postemergence suppression of the target weeds at 0.5 to 1 pint (0.125 to 0.25 lbs active) per broadcast acre when applied to susceptible weed seedlings less than 4 inches in height. Repeat applications may be required. Do not exceed 6 pints (1.5 lbs active) of Galigan 2E herbicide during the non-dormant stage of apricots, nectarines, olives, peaches, plums, and prunes. For a broader spectrum of grass weeds and broadleaf weeds control in the tree row middles, a tank mixture of Galigan 2E herbicide with either paraquat (Parazone 3SL) or glyphosate (Glyphogan) pesticide product which is to be tank mixed with Galigan 2E Herbicide.

##### **Method of Application**

**Ground Application:** Apply a minimum spray volume of 10 gallons of water per acre. Use higher volumes to ensure adequate coverage in high densities of emerged weeds or heavy trash. Use conventional low-pressure ground spray equipment with flat fan spray nozzles at 20 to 40 psi. Position an off-center nozzle at the end of the boom. Calibrate spray equipment carefully before each use.

**3120 Highwoods Blvd, Suite 100, Raleigh NC 27604**  
**T: 919.256.9305 ♦ F: 919.256.9308**

**Chemigation Application:** Apply this product only through flood (basin) irrigation system, or low-volume sprinkler (microsprinkler) and drip (trickle) irrigation systems designed to distribute irrigation water beneath the tree canopy. For additional information on these systems, see the **APPLICATION THROUGH IRRIGATION SYSTEMS-CHEMIGATION** section of this label.

**Cultural Considerations for All Applications:** In order to provide maximum effectiveness of preemergence activity of Galigan 2E herbicide, the berm or soil surface must be level, smooth, and free of crop or weed trash (decaying leaves, clippings, dead weeds, etc.). Remove leaves and trash by blowing the area to be treated or by thoroughly mixing the trash into the soil through cultivation prior to herbicide applications.

Cultural practices that result in redistribution or disturbance of the soil surface after treatment will decrease the herbicidal effectiveness of Galigan 2E herbicide. Cutting water furrows or cultivations that mix untreated soil into treated areas will also reduce the effectiveness of the treatment. For best results, apply to established berms or soil surfaces that are left undisturbed during the time period for which weed control is desired.

**Apricots, Nectarines, Olives, Peaches, Plums, and Prunes in California  
Nondormant Application  
Specific Use Restrictions**

In addition to the following, also observe **PRODUCT USE RESTRICTIONS** listed at the beginning of this label.

- Read and observe all label direction before using. When tank mixing, always read all individual manufacturer labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- When applied as a non-dormant treatment, Galigan 2E herbicide can only be applied to apricots, peaches, nectarines, plums, and prunes after May 1. Galigan 2E Herbicide can only be applied as a non-dormant treatment to olives after bloom.
- Do not apply Galigan 2E Herbicide within 14 days of harvest of fruit.
- Do not apply more than 6 pints (1.5 lb ai) per broadcast acre of Galigan 2E Herbicide during the non-dormant season.
- Apply Galigan 2E Herbicide only to healthy trees.
- Direct spray toward the base of the tree. Avoid direct herbicide contact with foliage and fruit.

Read the “**LIMITATION OF WARRANTY AND LIABILITY**” in the label booklet for Galigan 2E before using.

Galigan is a registered trademark of Agan Chemical Manufacturers, Ltd.

05/16/12  
EPA 041111



**3120 Highwoods Blvd, Suite 100, Raleigh NC 27604  
T: 919.256.9305 ♦ F: 919.256.9308**