



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

**Proline® 480 SC Fungicide**

**EPA Reg No. 264-825**

**plus**

**Folicur® 3.6 F Foliar Fungicide**

**EPA Reg No. 264-752**

**Reduced Application Rates for Control of Fusarium Head Blight and Leaf Diseases in Wheat**

***PRODUCT BULLETIN***

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF MINNESOTA, MISSOURI, NEBRASKA, NEW JERSEY, NEW MEXICO, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**Proline + Folicur can be applied at reduced rates of 3 + 3 fl oz/acre or 4 + 4 fl oz/acre for control of Fusarium head blight (Fusarium spp.) and certain leaf diseases in wheat.**

**Directions for Use**

<b>Fusarium Head Blight Control:</b>	The optimal time to apply Proline + Folicur is at 15% flower (Feekes 10.51). Application may be made using either ground or aerial application equipment. Spray equipment must be set up to provide good coverage to wheat heads. For thorough coverage of the wheat head using ground application equipment, use forward, forward and backward mounted nozzles or nozzles that have a two-directional spray. Operate nozzles within the spray pressure recommendations suggested by the manufacturer.
<b>Leaf Disease Control (Leaf Rust, Stem Rust, Stripe Rust, Septoria Leaf and Glume Blotch, Tan Spot):</b>	Apply Proline + Folicur as a preventative foliar spray when the earliest disease symptoms appear on the leaves or stems.

**RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 4 fl oz/acre of Folicur per acre per crop year.
- Do not apply more than 4 fl oz/acre of Proline in tank-mixture with Folicur per acre per crop year.
- Do not apply within 30 days of harvest.
- Refer to the Federal Section 3 labels of Proline and Folicur for additional use directions, precautions and/or restrictions.

**For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577**

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

**or**

**Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)**

*As with any crop-protection product, always read and follow label directions.  
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*