

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW JERSEY
FOR CONTROL OF CUCUMBER BEETLE IN CUCURBITS**

RESTRICTED USE PESTICIDE

Due to Reproductive Effects

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during mixing, loading, equipment repair and equipment cleaning. Certified applicators must ensure that all persons involved in these activities under their direct supervision are informed of the precautionary statements.

Metasystox-R®

**SPRAY CONCENTRATE
SYSTEMIC INSECTICIDE
FOR EFFECTIVE SYSTEMIC CROP PROTECTION
EPA REG. No. 10163-220**

ACTIVE INGREDIENT:	% By Wt.
S-[2-(Ethylsulfanyl)ethyl] O,O-dimethyl phosphorothioate.....	25.0%
INERT INGREDIENTS	75.0%
	TOTAL 100.0%

Contains 2 lbs. S-[2-(Ethylsulfanyl)ethyl] O,O-dimethyl phosphorothioate per U.S. gallon
(This product contains aromatic petroleum distillates.)

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

This recommendation must be in the possession of the user at the time of pesticide application.

CROP	PEST	PINTS PER ACRE	COMMENTS
Cucumbers	Cucumber Beetle	1½ - 2	Apply specified dosage per acre in sufficient water for complete coverage but not less than one gallon per acre. For application by irrigation systems, apply specified dosage per acre. Follow all directions given under the CHEMIGATION section of this label.
	<ul style="list-style-type: none"> Do not apply more than 2 time per season. Do not apply within 3 days of harvest. 		
Pumpkins Summer Squash Winter Squash	Cucumber Beetle	1½ - 2	Apply specified dosage per acre in sufficient water for complete coverage but not less than one gallon per acre. For application by irrigation systems, apply specified dosage per acre. Follow all directions given under the CHEMIGATION section of this label.
	<ul style="list-style-type: none"> Do not apply more than 1 time per season. Do not apply to Summer Squash within 3 days of harvest. Do not apply to Pumpkins and Winter Squash within 14 days of harvest. 		
Muskmelons (Cantaloupes), Other Melons	Cucumber Beetle	1½ - 2	Apply specified dosage per acre in sufficient water for complete coverage but not less than one gallon per acre. For application by irrigation systems, apply specified dosage per acre. Follow all directions given under the CHEMIGATION section of this label.
	<ul style="list-style-type: none"> Do not apply more than 3 times per season. Do not apply within 14 days of harvest. 		

FOR CONTROL OF CUCUMBER BEETLE IN CUCURBITS (Continued)

CROP	PEST	PINTS PER ACRE	COMMENTS
Watermelons	Cucumber Beetle	1½ - 2	Apply specified dosage per acre in sufficient water for complete coverage but not less than one gallon per acre. For application by irrigation systems, apply specified dosage per acre. Follow all directions given under the CHEMIGATION section of this label.
			<ul style="list-style-type: none">• Do not apply more than 2 times per season.• Do not apply within 7 days of harvest.

2(ee) REGISTRATION

Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

02-26-01
Metasystox-R ® is a trademark of Bayer