



Recommendation

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision, and only for the uses covered by the certified applicator's certification.

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN ALABAMA, DELAWARE, FLORIDA, GEORGIA, LOUISIANA, MARYLAND, MISSISSIPPI, NEW JERSEY, NORTH CAROLINA, PENNSYLVANIA, SOUTH CAROLINA, TENNESSEE AND VIRGINIA

**Bifenture® EC Agricultural Insecticide
EPA Reg. No. 70506-57**

**For control of Kudzu bug in
COTTON, POTATO, SOYBEANS, and SUCCULENT BEANS AND PEAS**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Cotton	0.1 lb. ai (6.4 fl. oz product)	Apply Bifenture EC in water in a minimum of 5 gallons/A by ground or 1 gallon/A by air. Do not apply more than 0.5 lb. ai (32 fl. oz. of product) per acre per season. Use an intensive scouting program for appropriate application timing. Repeat applications may be necessary for control. Do not graze livestock in treated areas or cut treated crops for feed.	14



Recommendation

CROP	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Potato	0.1 lb. ai (6.4 fl. oz. product)	Apply Bifenture EC in water in a minimum of 10 gallons/A by ground or 3 gallons/A by air. Do not make more than 2 foliar applications per season. Do not make applications less than 21 days apart. Do not apply more than 0.5 lb. ai (32 fl. oz. of product) per acre per season.	21
Soybeans	0.1 lb. ai (6.4 fl. oz product)	Apply Bifenture EC in water in a minimum of 10 gallons/A for foliar applications. Do not apply more than 0.3 lb. ai (19.2 fl. oz. of product) per acre per season. Do not apply more than once every 30 days.	18
Succulent Peas and Beans	0.1 lb ai (6.4 fl. oz product)	Apply Bifenture EC in water in a minimum of 10 gallons/A by ground or 2 gallons/A by air. When applying by air, substitute 1-2 quarts of emulsified oil for 1-2 quarts water, if desired. For best control, thorough coverage is necessary. Do not apply more than 0.2 lb. ai (12. 8 fl. oz. product) per acre per season.	3

Rev. 07/16/2013