




Metribuzin 75DF

HERBICIDE

PULL HERE TO OPEN

FOR CONTROL OF CERTAIN GRASSES AND BROADLEAF WEEDS

| | |
|--|---------------------|
| ACTIVE INGREDIENT: | % BY WEIGHT |
| Metribuzin: 4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one | 75.0% |
| INERT INGREDIENTS: | 25.0% |
| | TOTAL 100.0% |

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

| FIRST AID | |
|----------------------|--|
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |

(continued)

| FIRST AID <i>(continued)</i> | |
|--|---|
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.</p> | |
| <p>NOTE TO PHYSICIAN: Treat patient symptomatically.</p> | |

For additional precautionary, handling and use statements, see inside of this booklet.

EPA Reg. No. 66222-106 EPA Est. No. 5905-IA-001 ^{HD}
 39578-TX-001 ^S
 37429-GA-001 ^{BT}
 37429-GA-002 ^{BB}

Letter(s) in lot number correspond(s) to superscript in EPA Est. No.



M A N A

Manufactured for:
**Makhteshim Agan
 of North America, Inc.**
 4515 Falls of Neuse Road
 Suite 300
 Raleigh, NC 27609

EPA 110504/092605/051309/Notif 060508/Rev B

Net Contents: 20 Pounds

14030

PF-52561

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors, Unwind, and Dieline.**

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors. **THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.**

WE CANNOT PROCESS THIS ORDER WITHOUT AN AUTHORIZED SIGNATURE

ARTWORK IS APPROVED

REVISED PROOF NEEDED

Signed _____


Date _____

Please Return To: _____



4201 North Westport Avenue, Sioux Falls, South Dakota 57107
 Phone: (605) 978-0451 • Fax: (605) 978-0463

Please indicate the correct unwind position



PROOF DATE: 1/20/11
 CUSTOMER: MAKHTESHIM
 JOB NUMBER: 52561
 LABEL SIZE: 6.75" x 6.75"
 LEAFLET FLAT SIZE: 6.5" x 11.5"
 LEAFLET FOLDED SIZE: 6.5" x 5.75"
 LABEL COLORS: BLACK PMS 1235
VARNISH
 LEAFLET "IN" COLORS: BLACK
 LEAFLET "OUT" COLORS: BLACK PMS 1235

dieline does not print





Metribuzin 75DF

HERBICIDE

FOR CONTROL OF CERTAIN GRASSES AND BROADLEAF WEEDS

| ACTIVE INGREDIENT: | % BY WEIGHT |
|--|-------------|
| Metribuzin: 4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4H)-one | 75.0% |
| INERT INGREDIENTS: | 25.0% |
| TOTAL 100.0% | |

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

| FIRST AID | |
|----------------------|--|
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |

(continued)

EPA Reg. No. 66222-106

EPA Est. No. 5905-IA-001 ^(HD)

39578-TX-001 ^(S)

37429-GA-001 ^(BT)

37429-GA-002 ⁽⁸⁰⁾

Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

FIRST AID (continued)

| | |
|--------------------------------|---|
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact PROSAR at 1-877-250-9291 for emergency medical treatment information.

NOTE TO PHYSICIAN: Treat patient symptomatically.

For additional precautionary, handling and use statements, see inside of this booklet.



M A N A

Manufactured for:
**Makhteshim Agan
of North America, Inc.**
4515 Falls of Neuse Road
Suite 300
Raleigh, NC 27609

EPA 110504/092605/051309/Notif 060508/Rev B

Net Contents: 20 Pounds

14030





**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May be harmful if swallowed or absorbed through skin. Causes moderate eye irritation.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made out of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Do not contaminate feed or food. Keep out of reach of children.

Obtain prompt medical aid if poisoning should occur.

Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Poisoning is accompanied by breathing difficulties and sedation.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment wash waters.

GROUND WATER ADVISORY

Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface and where the soils are very permeable, i.e., well-drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Applicators and other handlers must use chemical-resistant gloves such as butyl rubber, or nitrile rubber, or neoprene rubber, or natural rubber
- Shoes plus socks

IMPORTANT: Read these entire Directions and Conditions of Sale before using Metribuzin 75DF.

GENERAL INFORMATION

MIXING

When using Metribuzin 75DF make sure the sprayer is completely clean, free of rust or corrosion which occurs from winter storage. Examine strainers and screens to be sure the sprayer is clean from previously used pesticides.

Any tank mixes containing Metribuzin 75DF should be kept agitated and sprayed out immediately. Do not allow tank mixes to stand for prolonged periods of time.

The proper mixing procedure for Metribuzin 75DF alone or in tank mix combinations with other herbicides is:

1. Fill the spray tank 1/4 to 1/3 full with clean water.
2. Add recommended rate of Metribuzin 75DF while recirculating and with agitator running.
3. Follow the triple rinse procedure described under STORAGE AND DISPOSAL to insure that all product is removed from the container.
4. Mix thoroughly and add clean water to fill spray tank to desired level.
5. Add the other herbicide to tank last and agitate thoroughly.
6. Continue agitation during application and until sprayer tank is empty.

When an adjuvant is to be used with this product, Makhteshim Agan of North America, Inc. suggests the use of a Chemical Products and Distributors Association certified adjuvant.

SOIL TEXTURE

As used on this label, **coarse soils** are loamy sand or sandy loam soils. **Medium soils** are loam, silt loam, silt, sandy clay, or sandy clay loam. **Fine soils** are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.



GENERAL PRECAUTIONS AND RESTRICTIONS

Do not allow sprays to drift on to adjacent desirable plants.

Apply this product only as specified on this label.

Do not rotate any crop not listed on this label for 18 months following application of Metribuzin 75DF.

Do not use on other crops grown for food or forage. Observe all cautions and limitations on labeling of all products used in mixtures.

For all uses: Low-pressure and high-volume hand-wand equipment is prohibited.

CHEMIGATION

Metribuzin 75DF is recommended for application through sprinkler irrigation equipment to potatoes, soybeans, tomatoes, and asparagus as directed on this label. Refer to the crop sections of this label for recommended rates, weeds controlled or suppressed, restrictions, and special precautions.

Apply this product only through sprinkler including center pivot, lateral move, or solid set irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

Calibration: Center Pivot and Self-Propelled Lateral Move Systems: Sprinkler irrigation systems must be accurately calibrated for application of Metribuzin 75DF. Greater accuracy in calibration (and distribution) will be achieved by injecting a larger volume of a more dilute mixture of product and water per hour. Follow the steps below to calibrate center pivot and lateral move systems.

1. Determine number of minutes required to make one complete revolution while applying 1/4 to 3/4 inch of water per acre.
2. With the system at operating pressure determine the exact number of minutes required to inject one gallon of water.
3. Divide the time required for one revolution (step 1) by the time required to inject one gallon (step 2). This gives total gallons of product-water mixture to be added to nurse tank.
4. Add required amount of water to nurse tank and start the agitation system. Then add sufficient Metribuzin 75DF at the recommended rate (See RECOMMENDED BROADCAST APPLICATIONS) to the nurse tank.

EXAMPLE: If 20 hours (1200 minutes) were required for one revolution and if 2 minutes were required to inject one gallon, then a total of 600 gallons of product-water mixture are required (1200/2 = 600); to treat 135 acres at 2/3 lb./acre, 90.5 lb. of Metribuzin 75DF are required.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection

pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in the injection nurse tanks during the herbicide application sufficient to keep herbicide in suspension.

Apply specified dosage in 1/4 to 3/4 inch of water (1/4 to 1/2 inch of water on sandy soils) per acre as a continuous injection in center pivot and lateral move systems or in the last 15 to 30 minutes of set in permanent solid set sprinkler systems. Application of more than the quantity of irrigation water recommended on this label may result in decreased product performance by removing the chemical from the zone of effectiveness. Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap, excessive crop injury may result. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. To ensure that lines are flushed and free of remaining pesticide, an indicator dye may be injected into the lines to mark the end of the application period.

Use a minimum of 1 part water to 1 part herbicide for injection. The use of a larger volume of water will insure greater accuracy and more uniform distribution.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
3. Where states have more stringent regulations, they should be observed.
4. The applicator should be familiar with and take into account the information covered in the AERIAL DRIFT REDUCTION ADVISORY INFORMATION.

INFORMATION ON DROPLET SIZE: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see WIND, TEMPERATURE AND HUMIDITY, and TEMPERATURE INVERSIONS).



CONTROLLING DROPLET SIZE:

- Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure-Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles-Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation-Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH: For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

WIND: Drift potential is lowest between wind speeds of 2 to 10 mph; however, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS: Metribuzin 75DF should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

APPLICATION OF METRIBUZIN 75DF WITH HERBICIDE SPRAY EQUIPMENT

Use a standard low pressure (20 to 40 psi) herbicide boom sprayer equipped with suitable nozzles and screens no finer than 50-mesh in nozzle and in-line strainers. Agitate thoroughly before and during application with bypass agitation.

GROUND APPLICATION: Apply the proper rate of Metribuzin 75DF in a minimum of 10 to 40 gallons of spray mixture per acre broadcast.

BANDED APPLICATION: Use proportionally less Metribuzin 75DF per acre in a band versus a broadcast application. For band application, use 1/4 to 1 gallon of spray mix per inch of band width regardless of row spacing.

Examples: (1) To treat a 15-inch band on rows 30 inches apart, use one-half of the broadcast rate of Metribuzin 75DF. (2) To treat a 14-inch band on rows 42 inches apart, use one-third of the broadcast rate of Metribuzin 75DF.

AERIAL APPLICATION: Where permitted, apply specified rate in a minimum of 2 to 10 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

NOTE: Do not apply aerially when Metribuzin 75DF is tank mixed with Lasso®.

For All Applications of Metribuzin 75DF: Sprayer must be accurately calibrated before applying Metribuzin 75DF. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases. Avoid over application, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result.) Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops, such as cole crops, onions, peas, or strawberries are present in adjacent fields or in areas where wheat is growing in coarse-textured soils.

SPRAYER CLEANUP: Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution of Metribuzin 75DF from the spray tank and dispose of according to label disposal instructions. Rinse the spray tank and refill with water adding a heavy-duty detergent at the rate of one cup per 20 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles, or spray rig. All rinse water must be disposed of in compliance with local, state, and Federal guidelines.

APPLICATION OF METRIBUZIN 75DF IN FLUID FERTILIZERS

Metribuzin 75DF may be applied in fluid fertilizer solutions to alfalfa and soybeans by following the appropriate mixing procedures and compatibility check. When using tank mix combinations, be sure all components are compatible.

Compatibility checks of Metribuzin 75DF and tank mix combinations which include Metribuzin 75DF should be made for each batch of fluid fertilizer because of the variability of these fertilizers.

Compatibility Check:

1. Premix 2 teaspoons of Metribuzin 75DF with 8 teaspoons of water (1:4 ratio) in a quart jar by adding the water first and follow with Metribuzin 75DF. Mix thoroughly. If a second herbicide is to be used, double the amount of water (1:8 ratio) and add the second herbicide after mixing Metribuzin 75DF first.
2. Then pour 1 pint of fluid fertilizer into the quart jar and shake well.
3. Allow to stand for 5 minutes.

THIS COMPATIBILITY CHECK SHOULD ONLY BE USED WHEN MIXING WITH FLUID FERTILIZERS.





Interpretation of Results: If the solution in the jar appears to be uniform without signs of agglomeration or without a separation of an oily film on top of the fertilizer, the mixture may be used. If not, repeat the compatibility check using twice the amount of water or add a compatibility agent to the water. If separation occurs but the mixture can be resuspended by shaking, then application is possible with good agitation in the spray tank.

Tank Mixing Guidelines:

1. Add the required amount of water and compatibility agent (if required) to the tank. Start agitation system while adding Metribuzin 75DF and follow by adding the fluid fertilizer and agitate.
2. If a second herbicide is to be used, follow as above in 1, but use twice the amount of water. Start agitation and add Metribuzin 75DF and follow by adding the second herbicide and then continue filling the tank with fluid fertilizer.
3. Maintain continuous agitation to assure uniform spray mixture until the tank is emptied.

COMMERCIAL IMPREGNATION AND APPLICATION OF METRIBUZIN 75DF ON DRY BULK FERTILIZER

Dry bulk fertilizer may be impregnated or coated with Metribuzin 75DF for application to established alfalfa and to soybeans. All recommendations, cautions, and special precautions on this label must be followed along with state regulations relating to dry bulk fertilizer blending, impregnating, and labeling.

IMPREGNATION: To impregnate, use a system consisting of a belt, conveyor, or closed drum which is used for dry bulk fertilizer blending. Any commonly used fertilizer can be impregnated with Metribuzin 75DF except ammonium nitrate or fertilizers containing ammonium nitrate, potassium nitrate, or sodium nitrate. Do not use on powder limestone.

Apply using a minimum of 200 lb. dry bulk fertilizer per acre and up to a maximum of 450 lb. per acre. To impregnate or coat dry bulk fertilizer, mix Metribuzin 75DF with sufficient water to form a sprayable slurry. The delivery nozzles must be directed to deliver a fine spray toward the fertilizer for thorough coverage while avoiding spray contact with mixing equipment. Uniform impregnation of Metribuzin 75DF to dry bulk fertilizer will vary and if the absorptivity is not adequate, an absorbent powder may be added to produce a dry, free-flowing mixture. Micro-Cel E (Johns-Manville Product Corporation) is the recommended absorbent powder. When another herbicide is used with Metribuzin 75DF, mix and impregnate immediately.

Apply immediately after impregnation unless experience has shown that impregnated fertilizer can be stored without becoming lumpy and difficult to spread.

RATES: Select the recommended rate of Metribuzin 75DF per acre from the appropriate section of this label and refer to the formula below to determine the amount of Metribuzin 75DF which is to be impregnated on a ton of dry bulk fertilizer based on the amount of fertilizer which will be distributed on one acre.

$$\frac{\text{Lb. Metribuzin 75DF}}{\text{Acre}} \times \frac{2000 \text{ Lb. Fertilizer}}{\text{Acre}} = \frac{\text{Lb. Metribuzin 75DF}}{\text{Ton of Fertilizer}}$$

APPLICATION: Uniform application is essential for satisfactory weed control. Accurate calibration of fertilizer application equipment is essential for uniform distribution to the soil surface. The recommended method of application is to apply 1/2 the recommended rate and overlap 50 percent or to double apply by splitting the middles to obtain the best distribution pattern.

If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

INCORPORATION AND COMBINATION USES: When Metribuzin 75DF is to be used in combination with another herbicide, follow directions on this label for combinations, rates, crops, incorporation, and special precautions.

SOYBEANS (Except California)

Metribuzin 75DF herbicide tank mix combinations are recommended for preplant incorporated applications, preemergence surface applications, Split-Shot applications, and Extended Split-Shot application. Metribuzin 75DF is also recommended as an overlay application following a preplant incorporated application of a recommended grass herbicide and alone as a preemergence surface application. All these applications can be applied with ground equipment and some can be applied with aerial spray equipment. In addition, Metribuzin 75DF can be applied as a postemergence directed spray to soybeans in certain states.

SPECIAL PRECAUTIONS (Soybeans): Injury to soybeans may occur when Metribuzin 75DF is used under the following conditions:

1. When soils have a calcareous surface area or a pH of 7.5 or higher.
2. Due to the sensitivity of certain soybean varieties, Metribuzin 75DF is not recommended for use on Altona, AP 55, AP 71, Asgrow 6520, Burlison, Coker 102, Coker 156, Dassel, GL 3202, Govan, Maple Amber, NB 3665, NKS 1884, Paloma 350, Portage, Regal, Semmes, Terra-Vig 505, Terra-Vig 606, Tracy, Vansoy, and Vinton 81. Consult your Makhteshim Agan of North America, Inc. Representative or your seed supplier for information on the tolerance to Metribuzin 75DF of newly released soybean varieties prior to use of Metribuzin 75DF.
3. When applied in conjunction with soil-applied organic phosphate pesticides.
4. Over application of boom overlapping may result in stand loss and soil residues.
5. Uneven application or improper incorporation can decrease the level of weed control and/or increase the level of injury.
6. When applied to any soil with less than 1/2% organic matter.
7. Soil incorporation deeper than recommended.
8. When sprayers are not calibrated accurately.
9. When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
10. When soybeans are planted less than 1 1/2 inches deep, particularly in preemergence application.

Activation: A minimum amount of soil moisture is required to activate Metribuzin 75DF. In areas of low rainfall, applications to dry soil should be followed with light irrigation of 1/4 acre inch of water. Do not apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

Grazing and Feeding Treated Vines: Treated vines may be grazed or fed to livestock 40 days after application when Metribuzin 75DF is applied alone or with Treflan®, Dual®, Prowl®, or Lasso®.

Do not use treated vines for feed or forage when Metribuzin 75DF is applied with Sonalan®, linuron plus Lasso, or linuron plus Dual.



| WEEDS CONTROLLED BY METRIBUZIN 75DF AND METRIBUZIN 75DF HERBICIDE TANK MIX COMBINATIONS | | | | | | | | | |
|---|-----------------------------------|-------------------------------|---|---|--|---|---|---|---|
| C= Control | S= Suppression or Erratic Control | P= Poor or No Control | | | O= No information (Control may range from poor to excellent) | | | | |
| 1= Metribuzin 75DF Alone | | 4= Metribuzin 75DF plus Dual | | | 7= Extended Split-Shot | | | | |
| 2= Metribuzin 75DF Split-Shot | | 5= Metribuzin 75DF plus Prowl | | | 8= Metribuzin 75DF plus Sonalan | | | | |
| 3= Metribuzin 75DF plus Treflan | | 6= Metribuzin 75DF plus Lasso | | | 9= Metribuzin 75DF plus linuron plus (Lasso or Dual) | | | | |
| ANNUAL BROADLEAF WEEDS | | | | | | | | | |
| Black Nightshade (<i>Solanum nigrum</i>) | P | P | P | C | P | C | C | P | S |
| Bristly Starbur (<i>Acanthospermum hispidum</i>) | C | C | C | C | C | C | C | C | C |
| Buffalobur (<i>Solanum rostratum</i>) | C | C | P | P | P | P | C | P | O |
| Carpetweed (<i>Mollugo verticillata</i>) | C | C | C | C | C | C | C | C | C |
| Cocklebur (<i>Xanthium pensylvanicum</i>) | S | C | S | S | S | S | C | S | S |
| Copperleaf, Hophombeam (<i>Acalypha ostryaefolia</i>) | C | C | C | C | C | C | C | C | C |
| Florida Beggarweed (<i>Desmodium tortuosum</i>) | C | C | C | C | C | C | C | C | C |
| Florida Pusley (<i>Richardia scabra</i>) | C | C | C | C | C | C | C | C | C |
| Galinsoga (<i>Galinsoga</i> spp.) | C | C | C | C | C | C | C | C | C |
| Horseweed, Marestalk (<i>Conyza canadensis</i>) | O | O | O | O | O | O | C | O | O |
| Jimsonweed (<i>Datura stramonium</i>) | C | C | C | C | C | C | C | C | S |
| Knotweed (<i>Polygonum</i> spp.) | C | C | C | C | C | C | C | C | C |
| Kochia (<i>Kochia scoparia</i>) | C | C | C | C | C | C | C | C | C |
| Lambsquarters (<i>Chenopodium</i> spp.) | C | C | C | C | C | C | C | C | C |
| Morningglory, Ivyleaf (<i>Ipomoea hederacea</i>) | P | P | S | P | P | P | P | P | P |
| Morningglory, Pitted (<i>Ipomoea lacunosa</i>) | P | P | S | P | P | P | P | P | P |
| Morningglory, Smallflower (<i>Jacquemontia tamnifolia</i>) | P | P | C | P | P | P | P | P | P |
| Morningglory, Tall (<i>Ipomoea purpurea</i>) | P | P | S | P | P | P | P | P | P |
| Pigweeds (<i>Amaranthus</i> spp.) | C | C | C | C | C | C | C | C | C |
| Prickly Sida/Teaweed (<i>Sida spinosa</i>) | C | C | C | C | C | C | C | C | C |
| Purslane (<i>Portulaca oleracea</i>) | C | C | C | C | C | C | C | C | C |
| Ragweed, Common (<i>Ambrosia artemisiifolia</i>) | C | C | C | C | C | C | C | C | C |
| Redweed (<i>Melochia corchorifolia</i>) | C | C | C | C | C | C | C | C | C |
| Russian Thistle (<i>Salsola kali</i>) | C | C | C | C | C | C | C | C | C |
| Sesbania (<i>Sesbania</i> spp.) | C | C | C | C | C | C | C | C | C |
| Shepherdspurse (<i>Capsella bursa-pastoris</i>) | C | C | C | C | C | C | C | C | C |
| Sicklepod (<i>Cassia obtusifolia</i>) | C | C | S | C | S | C | C | S | S |
| Smartweeds (<i>Polygonum</i> spp.) | C | C | C | C | C | C | C | C | S |
| Spotted Spurge (<i>Euphorbia maculata</i>) | C | C | P | C | P | C | C | P | O |
| Spurred Anoda (<i>Anoda cristata</i>) | C | C | C | C | C | C | C | C | O |
| Sunflower (<i>Helianthus</i> spp.) | C | C | S | S | S | S | C | S | P |
| Velvetleaf (<i>Abutilon theophrasti</i>) | C | C | C | C | C | C | C | C | C |
| Venice Mallow (<i>Hibiscus trionum</i>) | C | C | C | C | C | C | C | C | C |
| Wild Mustards (<i>Brassica</i> spp.) | C | C | C | C | C | C | C | C | C |



| WEEDS CONTROLLED BY METRIBUZIN 75DF AND METRIBUZIN 75DF HERBICIDE TANK MIX COMBINATIONS | | | | | | | | | |
|---|-----------------------------------|---|---|--|---|---|--|---|---|
| C= Control | S= Suppression or Erratic Control | | | P= Poor or No Control | | | O= No information (Control may range from poor to excellent) | | |
| 1= Metribuzin 75DF Alone | 4= Metribuzin 75DF plus Dual | | | 7= Extended Split-Shot | | | | | |
| 2= Metribuzin 75DF Split-Shot | 5= Metribuzin 75DF plus Prowl | | | 8= Metribuzin 75DF plus Sonalan | | | | | |
| 3= Metribuzin 75DF plus Treflan | 6= Metribuzin 75DF plus Lasso | | | 9= Metribuzin 75DF plus linuron plus (Lasso or Dual) | | | | | |
| ANNUAL GRASSES | | | | | | | | | |
| Barnyardgrass (<i>Echinochloa crus-galli</i>) | S | C | C | C | C | C | C | C | C |
| Bluegrass (<i>Poa annua</i>) | C | C | C | C | C | C | C | C | C |
| Broadleaf Signalgrass (<i>Brachiaria platyphylla</i>) | C | C | C | C | C | C | C | C | O |
| Browntop Millet (<i>Panicum ramosum</i>) | C | C | C | P | C | S | C | O | O |
| Crabgrass (<i>Digitaria</i> spp.) | C | C | C | C | C | C | C | C | C |
| Crowfootgrass (<i>Dactyloctenium aegyptium</i>) | C | C | C | C | C | C | C | O | O |
| Cupgrass (<i>Echinochloa gracilis</i>) | P | C | P | P | P | P | C | O | O |
| Foxtails (<i>Setaria</i> spp.) | S | C | C | C | C | C | C | C | C |
| Goosegrass (<i>Eleusine indica</i>) | C | C | C | C | C | C | C | C | C |
| Johnsongrass, Seedling (<i>Sorghum halepense</i>) | C | C | C | C | C | C | C | C | O |
| Junglerice (<i>Echinochloa colonum</i>) | C | C | C | C | C | C | C | C | O |
| Nutsedge, Yellow (<i>Cyperus esculentus</i>) | P | P | P | C | P | C | C | P | O |
| Panicum, Fall (<i>Panicum dichotomiflorum</i>) | P | C | C | C | C | C | C | C | C |
| Panicum, Texas (<i>Panicum texanum</i>) | P | C | C | P | C | S | S | C | O |
| Red Rice (<i>Oryza sativa</i>) | P | C | C | C | P | C | C | O | O |
| Sandbur (<i>Cenchrus</i> spp.) | P | C | C | P | C | S | S | O | O |
| Shattercane (<i>Sorghum bicolor</i>) | P | C | C | P | P | P | P | C | O |
| Sorghum, Volunteer (<i>Sorghum</i> spp.) | P | C | C | P | P | P | P | O | P |
| Sprangletop (<i>Leptochloa</i> spp.) | P | C | C | P | P | P | P | O | P |
| Stinkgrass (<i>Eragrostis</i> spp.) | P | C | C | P | P | P | P | O | P |
| Wheat, Volunteer (<i>Triticum</i> spp.) | P | P | P | P | P | P | P | O | P |
| Witchgrass (<i>Panicum capillare</i>) | P | C | C | C | C | C | C | C | O |

Rate Ranges: Where a rate range is shown, use a lower rate on soils that are coarse textured or low in organic matter. Use a higher rate on soils that are relatively fine textured or high in organic matter.

Replanting: If replanting is necessary in fields treated with Metribuzin 75DF as directed on this label, the field may be replanted to soybeans. When replanting, a minimum of tillage is recommended. Do not apply a second treatment as injury to soybeans may occur.





METRIBUZIN 75DF ALONE

Metribuzin 75DF (Alone) Pre-emergence Application: The following rates of Metribuzin 75DF may be applied preemergence to soybeans through center pivot or lateral move sprinkler irrigation systems that apply water in a uniform manner. Refer to CHEMIGATION section of this label for directions.

Metribuzin 75DF can be applied broadcast or banded. This application may be

made during planting or as a separate operation after planting but before crop emergence. See the GENERAL INFORMATION section in the front of this label.

Do not apply to sand soils or to sandy loam or loamy sand soils containing less than 2% organic matter. Do not incorporate into soil or apply more than once per season.

| LB. OF METRIBUZIN 75DF PER ACRE | | | |
|---|-------------------------|------------|------------|
| SOIL TEXTURE | ORGANIC MATTER | | |
| | Less than 2% | 2 to 4% | Over 4% |
| Coarse Soils (Sandy loam, loamy sand) | Do not use ³ | 1/2 | 2/3 |
| Medium Soils ¹ (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 to 2/3 | 2/3 to 5/6 | 5/6 to 1 |
| Fine Soils ¹ (Silty clay, silty clay loam ² , clay, clay loam) | 2/3 to 5/6 | 5/6 to 1 | 1 to 1 1/6 |
| Mississippi Delta Only | 1 | 1 1/6 | 1 1/3 |

¹ For control of lambsquarters, redroot pigweed, and wild mustard, and for suppression of green, yellow, and giant foxtails on alkaline (calcareous) soils in Nebraska, Minnesota, South Dakota, and North Dakota only, apply Metribuzin 75DF at rates of 1/3 lb./acre on medium soils and 1/3 to 1/2 lb./acre on fine soils regardless of soil organic matter percentage (use 1/2 lb. only where soil pH is less than 7.5 and weed pressure is heavy). The 1/3 lb./acre rate of Metribuzin 75DF alone can be applied regardless of soil pH. For control of other weeds listed on this label use Metribuzin 75DF at full rates recommended in the table above, but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.

² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

³ Refer to the appropriate section of this label for use of Metribuzin 75DF on soybeans in coarse soils with 0.5% or more organic matter in certain states.

USES OF METRIBUZIN 75DF HERBICIDE IN COMBINATION WITH OTHER HERBICIDES

SEQUENTIAL APPLICATION OF SCEPTER® FOLLOWING METRIBUZIN 75DF

If needed, application of Metribuzin 75DF alone or in a registered tank mix according to directions on this label, may be followed by an early postemergence application of Scepter herbicide (1.5 lb./gal liquid or 70 DG) for control of cocklebur. Apply 1/6 to 1/3 pint of Scepter (0.7 to 1.4 ounces of Scepter 70 DG) in a minimum of 20 gals. of water per acre. Use 1/6 pint of Scepter (0.7 ounce of Scepter 70DG) if cockleburs are less than 3 inches tall or have fewer than 3 leaves and are actively growing. For cockleburs less than 6 inches tall and actively growing, use 1/3 pint of Scepter (1.4 ounces of Scepter 70 DG) per acre. Do not use Scepter when soybeans or cockleburs have been subjected to stress conditions such as temperature or moisture extremes. Do not exceed a total of 2/3 pint of Scepter (2.8 ounces of Scepter 70 DG) per acre in one season. Wait at least 10 days after application of Scepter before cultivating.

When preparing the spray mixture with Scepter, add 2 pts. of nonionic surfactant approved for use on growing crops and containing at least 80% active ingredient per 100 gallons of mixture. Apply crop oil concentrate (COC) at the rate specified on the COC label.

Use Scepter only in the states where it is registered as listed on the product label.

Apply Scepter at least 90 days before harvest of soybeans. Do not graze or feed soybean forage, hay, or straw to livestock.

Refer to the Scepter label for additional cautions and precautions, directions, limitations, and information on environmental hazards and planting of rotational crops.

SPLIT-SHOT APPLICATION

A preplant incorporated application of Metribuzin 75DF tank mixed with either Treflan, Lasso, Dual, Prowl, or Sonalan and followed by a preemergence surface application of Metribuzin 75DF alone after planting but prior to soybean emergence will control more broadleaf and grass weeds in soybeans than when either herbicide is used alone.

Refer to the Treflan, Lasso, Dual, Prowl, or Sonalan labels, and to appropriate sections of this label for directions on soil preparation, herbicide application, incorporation techniques, herbicide rates, weed species controlled, and restrictions for using tank mix combinations of Metribuzin 75DF. Carefully observe the SPECIAL PRECAUTIONS section concerning the use of Metribuzin 75DF in tank mix combinations on soybeans.

When a Split-Shot application of Metribuzin 75DF with Prowl, Treflan, or Sonalan is used, the preplant incorporated tank mix may be applied up to 21 days prior to planting soybeans; with Dual or Lasso, the preplant incorporated tank mix may be applied up to 14 days prior to planting.

On medium- and fine-textured soils with greater than 2% organic matter, a rate range is recommended for the Metribuzin 75DF preemergence overlay application. The higher rate should be used (a) in fields with a history of severe broadleaf weed pressure, (b) when the time between preplant incorporated tank mix and preemergence overlay applications approaches the maximum stated above, and/or (c) when the organic matter content of the soil is at the upper end of the indicated range.

For black nightshade control, refer to the appropriate sections of the Lasso, Dual, or Sonalan labels for specific instructions.





| SPLIT-SHOT APPLICATION | | | | | | |
|--|--|------|---|---------------------------------|----------------------------------|--|
| Preplant Incorporated Tank mix Application | | | - FOLLOWED BY - | | Preemergence Overlay Application | |
| SOIL TEXTURE ¹ | Rate of Combination Product/Acre | Plus | Rate of Metribuzin 75DF lb/Acre | Rate of Metribuzin 75DF lb/Acre | | |
| | | | | ORGANIC MATTER | | |
| | | | | Less than 2% | 2% to 4% | Over 4% |
| Coarse (light) sand, loamy sand, sandy loam | Treflan 1 pt. -OR- Lasso 2 to 2 1/2 qt. -OR- Dual 1 1/4 to 1 1/2 pt. -OR- Prowl 1 1/2 pt. -OR- Sonalan 1 1/4 to 2 pt. | Plus | 1/3 - followed by | 1/6 | 1/6 | 1/6 to 1/3 |
| Medium Loam, silt loam, sandy clay loam, silt, sandy clay | Treflan 1 1/2 pt. -OR- Lasso 2 1/2 to 3 qt. -OR- Dual 1 1/2 pt. -OR- Prowl 1 1/2 pt. -OR- Sonalan 1 3/4 to 2 1/2 pt. | Plus | 1/2 - followed by -OR- 1/3 ² - followed by | 1/6 1/3 | 1/6 to 1/3 1/3 to 1/2 | 1/3 to 1/2 (1/2 to 2/3 ³) |
| Fine (heavy) Silty clay loam*, clay loam, silty clay, clay | Treflan 2 pt. -OR- Lasso 2 1/2 to 3 qt. -OR- Dual 2 to 2 1/2 pt. -OR- Prowl 1 1/2 to 2 pt. -OR- Sonalan 2 1/4 to 3 pt. | Plus | 2/3 - followed by -OR- 1/2 ² - followed by | 1/6 1/3 | 1/6 to 1/3 1/3 to 1/2 | 1/3 to 1/2 (1/2 to 2/3 ³) |

* Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

¹ On coarse-textured soils, do not use on sand soils with less than 1% organic matter or on loamy sand or sandy loam soils with less than 0.5% organic matter. However, on coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on sand soils with less than 2% organic matter or on loamy sand or sandy loam soils with less than 1% organic matter.

² Use this lower rate of Metribuzin 75DF in the preplant incorporated tank mix on soils having a calcareous surface area or a pH of 7.5 or higher and in those situations where soils within a field vary extremely in texture or organic matter content.

³ Reduce this preemergence overlay rate of Metribuzin 75DF by 1/6 lb./acre when using SPLIT-SHOT application on soils with over 4% organic matter and which have a calcareous surface area or a pH of 7.5 or higher.

| EXTENDED SPLIT-SHOT APPLICATION | | | | | | | | | |
|---|--|------|----------------------------------|----------------------------------|----------------------------------|------|----------------------------------|------------|------------|
| Early Preplant Tank Mix Application (Surface-Applied or Shallow Incorporated) | | | | Preemergence Overlay Application | | | | | |
| SOIL TEXTURE ¹ | Rate of Combination Product/Acre | Plus | Rate of Metribuzin 75DF lb./Acre | Followed By | Rate of Combination Product/Acre | Plus | Rate of Metribuzin 75DF lb./Acre | | |
| | | | | | | | ORGANIC MATTER | | |
| | | | | | | | 1/2 to 2% | 2% to 4% | over 4% |
| Coarse (light) sand, loamy sand, sandy loam | Dual 1 1/3 pt. OR Lasso 1 1/2 to 2 qt. | plus | 1/3 to 1/2 | Dual OR Lasso | 2/3 pt. 1 1/2 qt. | plus | 1/6 | 1/6 to 1/3 | 1/3 |
| Medium Loam, silt loam, sandy clay loam, silt, sandy clay | Dual 1 3/4 pt. OR Lasso 2 to 3 qt. | plus | 2 1/2 to 2/3 ² | Dual OR Lasso | 3/4 pt. 1 to 2 qt. | plus | 1/3 | 1/3 to 1/2 | 1/2 to 2/3 |
| Fine (heavy) Silty clay loam*, clay loam, silty clay, clay | Dual 2 pt. OR Lasso 2 to 3 qt. | plus | 2 2/3 to 5/6 ² | Dual OR Lasso | 1 pt. 1 to 2 qt. | plus | 1/3 | 1/3 to 1/2 | 1/2 to 2/3 |

* Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

¹ On coarse-textured soils, do not use on sand soils with less than 1% organic matter. However, on coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on sand soils with less than 2% organic matter or on loamy sand or sandy loam soils with less than 1% organic matter.

² Use this lower rate of Metribuzin 75DF in the early preplant tank mix on soils having a calcareous surface area or a pH of 7.5 or higher and in those rare situations where soils within a field vary extremely in texture or organic matter content.





EXTENDED SPLIT-SHOT APPLICATION

(Includes No-Till, Reduced-Till, Ridge-Till, Mulch-Till)

An early preplant (surface-applied or shallow incorporated) application of Metribuzin 75DF tank mixed with either Dual or Lasso followed by a preemergence surface application of Metribuzin 75DF tank mixed with Dual or Lasso after planting but prior to soybean emergence will control more broadleaf and grass weeds in soybeans than either herbicide used alone.

An Extended Split-Shot application will decrease the need for tillage and/or contact herbicides for the control of existing vegetation prior to planting, while providing residual control of weeds after planting.

When an Extended Split-Shot application of Metribuzin 75DF with Dual or Lasso is used, the preplant tank mix combination may be applied 15 to 30 days prior to planting soybeans. Follow directions on the label accompanying the product for Split-Shot applications from 0 to 14 days before planting.

Where a rate range is recommended, the higher rates should be used (a) in fields with a history of severe weed pressure, (b) when the time between early preplant tank mix and preemergence overlay applications approaches the maximum 30 days, (c) when the organic matter content of the soil is at the upper end of the indicated range, (d) when heavy crop residues are present on the soil surface, and/or (e) when the early preplant tank mix application is shallow incorporated (e.g. use 2 to 2 1/2 qt. Lasso in the early preplant tank mix when surface applied and use 2 1/2 to 3 qt. Lasso when the tank mix is to be lightly incorporated).

When weeds exceed 1 to 1 1/2 inches in height or diameter at application, use a contact herbicide such as Round-Up® or Gramoxone®.

Refer to the Dual or Lasso label and to appropriate sections of this label for additional information on soil preparation, herbicide application, weeds controlled, precautions, restrictions, limitations, and sprayer clean up.

METRIBUZIN 75DF PLUS SONALAN

Metribuzin 75DF plus Sonalan Overlay Application: Metribuzin 75DF may be applied as a preemergence overlay application following a preplant incorporated application of Sonalan 3 EC. Consult the Sonalan label for specific directions on use, recommendations, restrictions, and any additional weeds not specified on this label.

Metribuzin 75DF plus Sonalan Tank mix Application: Incorporate the tank mixture into the top 1 to 2 inches of soil within 21 days before planting according to label directions for Sonalan.

Apply Metribuzin 75DF plus Sonalan preplant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If soybeans are planted on beds, apply and incorporate the tank mixture after bed formation.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: Sonalan should be uniformly applied and thoroughly mixed into the soil within 2 days after application. For specific application information, refer to the APPLICATION under GENERAL INFORMATION section in the front of this label.

SPECIAL PRECAUTIONS (Metribuzin 75DF plus Sonalan): For additional precautions, restrictions, limitations, incorporation, and sprayer clean-up information, refer to the appropriate sections of this label and the Sonalan label.

For black nightshade control, refer to the Sonalan label for specific rates and application instructions.

BROADCAST RATES

| SOIL TEXTURE | Metribuzin 75DF | Sonalan 3EC |
|--|-----------------|----------------|
| | Lb./Acre | Pt./Acre |
| Coarse ¹ (Sandy loam, loamy sand) | 1/3 | 1 1/4 to 2 |
| Medium ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 | 1 3/4 to 2 1/2 |
| Fine ³ (Silty clay, silty clay loam ³ , clay, clay loam) | 2/3 | 2 1/4 to 3 |

¹ Do not use on coarse soils with less than 1% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.
³ For control of lambsquarters, redroot pigweed, wild mustard, and green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Metribuzin 75DF at rates of 1/3 lb./acre on medium soils and 1/3 to 1/2 lb./acre on fine soils regardless of soil organic matter percentage (use 1/2 lb. only where soil pH is less than 7.5 and weed pressure is heavy). The 1/3 lb. rate of Metribuzin 75DF in tank mix combination with Sonalan can be applied regardless of soil pH. For control of other weeds not listed on the label, use Metribuzin 75DF at full rates recommended in the table above, but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.

METRIBUZIN 75DF PLUS TREFLAN

Metribuzin 75DF and Treflan Overlay Application: Metribuzin 75DF may be applied as a preemergence broadcast or band overlay application following a preplant incorporated treatment of Treflan. Consult the Treflan label for specific directions for use, recommendations, restrictions, and any additional weeds not specified on the label.

Metribuzin 75DF plus Treflan Tank mix Application: A single application of a tank mix combination of Metribuzin 75DF and Treflan EC will control more broadleaf and grass weeds in soybeans than when either herbicide is used alone.

Prepare the soil surface by deep plowing, offset disking, or tandem disking prior to the application of the herbicide combination. The soil surface should be well prepared and free of clods and trash.

This Metribuzin 75DF plus Treflan tank mix combination may be applied and incorporated into the soil up to 10 days before planting.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: For specific application information, refer to the GENERAL INFORMATION section in the front of this label.

Apply Metribuzin 75DF plus Treflan to the soil surface and incorporate in the same operation if possible. Variable weed control may result from delayed incorporation if Metribuzin 75DF plus Treflan are applied to a wet, warm soil surface or if the wind velocity is 10 miles per hour or higher. Use machinery that mixes Metribuzin 75DF plus Treflan thoroughly with the soil. Incorporation may be delayed up to 24 hours after application. Shallow incorporation with implements set to cut less than 2 inches deep may result in erratic weed control. Do not use spike or spring-tooth harrows alone for incorporation.





| BROADCAST RATES | | |
|---|-----------------------------|------------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Treflan EC Pt./Acre |
| Coarse ¹ (Sandy loam, loamy sand) | 1/3 | 1 |
| Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 | 1 1/2 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) ³ | 2/3 | 2 |

¹ Do not use on coarse soils with less than 1% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.
³ For control of lambsquarters, redroot pigweed, wild mustard, and green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Metribuzin 75DF at rates of 1/3 lb./acre on medium soils and 1/3 to 1/2 lb./acre on fine soils regardless of soil organic matter percentage (use 1/2 lb. only where soil pH is less than 7.5 and weed pressure is heavy). The 1/3 lb. rate of Metribuzin 75DF in tank mix combination with Treflan can be applied regardless of soil pH. For control of other weeds listed on the label, use Metribuzin 75DF at full rates recommended in the table above but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.

Incorporation Equipment:

1. Set PTO-driven equipment (tillers, cultivators, hoes) to cut 2 to 3 inches deep and space rotors to provide a clean sweep of the soil. PTO equipment should not be operated at a speed greater than 4 miles per hour.
 2. Set disk to cut 4 to 6 inches deep and operate twice in different directions at 4 to 6 miles per hour.
 3. Set mulch treader and other similar disk-type implements to cut 3 to 4 inches deep and operate twice in different directions at 5 to 8 miles per hour.
- For Coarse- and Medium-Textured Soils Only:
4. Set rolling cultivator to cut 2 to 4 inches deep and operate twice at 6 to 8 miles per hour. Set bed conditioner (Do-All) to cut 2 to 4 inches deep and operate at 4 to 6 miles per hour.

SPECIAL PRECAUTIONS (Metribuzin 75DF plus Treflan): Seedling disease, cold weather, excessive moisture, high salt concentration, or drought may weaken soybean seedlings and increase possibility of damage from the tank mix. Do not plant soybeans deeper than 2 inches.

In the Central United States, do not plant sorghum or oats for 12 months where the tank mix has been applied unless 20 inches or more of irrigation and/or rainfall (total) was used to produce the crop. If less than 20 inches total water was used to produce the crop during the year, do not plant either crop for 18 months after the tank mix application. Cool, wet weather conditions during the early stage of growth may increase the possibility of injury to sorghum.

Do not rotate any crop not listed on this label for 18 months after tank mix application.

For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate section of this label. Do not use this tank mix combination on soils containing charcoal in Arkansas, Louisiana, and Mississippi.

METRIBUZIN 75DF PLUS DUAL

Metribuzin 75DF plus Dual Overlay Application: Apply a preplant incorporated treatment of Dual 8E as directed on that product label for use on soybeans. Follow with a preemergence treatment of Metribuzin 75DF as directed on this label for use on soybeans.

METRIBUZIN 75DF PLUS DUAL TANK MIX APPLICATIONS

Preplant Incorporated Application: Incorporate the tank mixture into the top 2 inches of soil within 14 days before planting using a disk, harrow, rolling cultivator, or similar implement.

Apply Metribuzin 75DF plus Dual preplant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If soybeans are planted on beds, apply and incorporate the tank mixture after bed formation.

Preemergence Application: Dry weather following preemergence application of Metribuzin 75DF plus Dual tank mixture may reduce effectiveness. If weeds develop, cultivate uniformly with shallow tillage equipment such as a rotary hoe that will not damage soybeans.

Mixing Instructions: Refer to the GENERAL INFORMATION section in the front of this label.





| BROADCAST RATES METRIBUZIN 75DF Plus DUAL Tank Mix Preemergence Applications | | |
|---|-----------------------------|---------------------|
| 0.5% TO 3% ORGANIC MATTER | | |
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Dual 8E Pt./Acre |
| Coarse ¹ (Loamy sand, sandy loam) | 1/3 | 1 1/4 |
| Medium (Loam, silt loam, silt) | 1/2 | 1 1/2 |
| Fine (Silty clay loam ² , sandy clay loam, silty clay, sandy clay, clay loam, clay) | 2/3 | 2 |
| Mississippi Delta Only (Silty clay, clay) | 1 | 2 |
| Over 3% ORGANIC MATTER | | |
| Coarse ¹ (Loamy sand, sandy loam) | 1/2 | 1 1/2 |
| Medium (Loam, silt loam, silt) | 2/3 | 2 |
| Fine (Silty clay loam ² , sandy clay loam, silty clay, sandy clay, clay loam, clay) | 2/3 to 5/6 | 2 to 2 1/2 |
| Mississippi Delta Only (Silty clay, clay) | 1 | 2 to 2 1/2 |

¹ Do not use on sand soils. Do not apply Metribuzin 75DF and Dual overlay or tank mix preemergence on loamy sand with less than 2% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

| BROADCAST RATES METRIBUZIN 75DF Plus DUAL Tank Mix Preplant Incorporated Applications | | |
|---|-----------------------------|---------------------|
| 0.5% to less than 3% ORGANIC MATTER | | |
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Dual 8E Pt./Acre |
| Coarse ¹ (Loamy sand, sandy loam) | 1/3 | 1 1/4 |
| Medium (Loam, silt loam, silt) | 1/2 | 1 1/2 |
| Fine (Silty clay loam ² , sandy clay loam, silty clay, sandy clay, clay loam, clay) | 2/3 | 2 |
| Mississippi Delta Only (Silty clay, clay) | 2/3 to 5/6 | 2 |
| 3% or Greater ORGANIC MATTER | | |
| Coarse ¹ (Loamy sand, sandy loam) | 1/3 | 1 1/2 |
| Medium (Loam, silt loam, silt) | 1/2 | 2 |
| Fine (Silty clay loam ² , sandy clay loam, silty clay, sandy clay, clay loam, clay) | 2/3 | 2 to 2 1/2 |
| Mississippi Delta Only (Silty clay, clay) | 2/3 to 5/6 | 2 to 2 1/2 |

¹ Do not use on sand soils. Do not apply Metribuzin 75DF plus Dual tank mix preplant incorporated on sand or loamy sand with less than 2% organic matter or crop injury may occur.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

SPECIAL PRECAUTIONS (Metribuzin 75DF and Dual): For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the Dual label.

METRIBUZIN 75DF PLUS PROWL

Metribuzin 75DF plus Prowl Overlay Application: Apply a preplant incorporated treatment of Prowl as directed on that product label for use on soybeans. Follow with a preemergence treatment of Metribuzin 75DF as directed on this label for use on soybeans.

Metribuzin 75DF plus Prowl Tank mix Application

Preplant Incorporated Application: Prepare the soil by plowing or disking to mix previous crop residues into the soil to a depth of 4 to 6 inches.

For specific application information refer to the GENERAL INFORMATION section in the front of this label.

Incorporate the tank mixture into the top 1 or 2 inches of soil within 7 days after application according to label directions for Prowl. Mechanical incorporation is not required if a rain of one-quarter inch or more occurs within 7 days after application. Soybeans must be planted no later than 7 days after application of the tank mixture.

Preemergence Application: Except for minimum and no-tillage systems, the seed bed should be firm and free of trash and clods.

For specific application information refer to the GENERAL INFORMATION section in the front of this label. **Do not apply Prowl preemergence north of Interstate 80.** This application must be made after planting and before crop emergence. Do not incorporate.

If cultivation is necessary because of soil crusting, soil compaction, or weed germination before rain or irrigation, use shallow tilling equipment such as a rotary hoe that does not damage soybeans.





Mixing Instructions: Refer to the GENERAL INFORMATION section in the front of this label.

For information on applying Metribuzin 75DF plus Prowl Overlay Application: in fluid or dry fertilizer refer to the APPLICATION OF METRIBUZIN 75DF IN FLUID FERTILIZERS or COMMERCIAL IMPREGNATION and APPLICATION OF METRIBUZIN 75DF ON DRY BULK FERTILIZER under the GENERAL INFORMATION section in the front of this label.

SOUTHERN STATES AND EASTERN COASTAL PLAINS

For use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, Southeastern Missouri "Bootheel" Region and Coastal Plains of Delaware, Maryland*, New Jersey*, and Virginia*.

*Metribuzin 75DF plus Prowl is not recommended for use on soils with less than 2% organic matter in the coastal plain of New Jersey or the Delmarva Peninsula.

| BROADCAST RATES Metribuzin 75DF Plus Prowl Tank mix Applications | | |
|--|-----------------------------|-------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Prowl Pt./Acre |
| Coarse ¹ (Sandy loam, loamy sand) | 1/3 | 1 1/2 |
| Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 | 1 1/2 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 2/3 | 1 1/2 to 2 |

¹ Do not use on sand soils. Do not use on loamy sand or sandy loam containing less than 1% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in certain regions of the U.S.

Do not use on muck or peat soils.

| NORTHEASTERN AND NORTH CENTRAL STATES For use only in Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New York, North Dakota, Ohio, Pennsylvania, South Dakota, Wisconsin, and Missouri (except the "Bootheel" Region). | | |
|--|-----------------------------|-------------------|
| BROADCAST RATES Metribuzin 75DF Plus Prowl Tank mix Applications | | |
| 1/2% to 3% ORGANIC MATTER | | |
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Prowl Pt./Acre |
| Coarse ¹ (Sandy loam, loamy sand) | 1/3 | 1 |
| Medium (Loam, silt loam, sandy clay, sandy clay loam) | 1/2 | 1 1/2 to 2 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 1/2 to 2/3 | 1 1/2 to 2 |
| Over 3% ORGANIC MATTER | | |
| Coarse ¹ (Sandy loam, loamy sand) | 1/2 | 1 1/2 |
| Medium (Loam, silt loam, sandy clay, sandy clay loam) | 1/2 to 2/3 | 1 1/2 to 2 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 2/3 to 5/6 | 2 to 2 1/2 |

¹ Do not use on sand soils. Do not use on loamy sand or sandy loam containing less than 1% organic matter. Where a range of rates is shown for medium and fine soils, use the higher rate if heavy weed infestations are anticipated.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in certain areas of the U.S.

Do not use on muck or peat soils.

SPECIAL PRECAUTIONS (Metribuzin 75DF and Prowl): Soil incorporation deeper than recommended will reduce weed control and can result in crop injury.

For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the Prowl label.

METRIBUZIN 75DF PLUS LASSO

Metribuzin 75DF Plus Lasso Tank Mix Application

PREEMERGENCE

Metribuzin 75DF may be used in a tank mix combination with Lasso as a preemergence band or broadcast application to soybeans in accordance with the specified soil types and dosages recommended.

For specific information regarding spray equipment, dilution rates, mixing, directions for use, methods of application, limitations, and restrictions, refer to the appropriate section of this label.

Refer to the Lasso label for pertinent recommendations, directions for use, restrictions, and any additional weeds not specified on this label.

Do not use on muck soils.





| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Lasso Tank mix Preemergence Application (Broadcast Rates) | | | |
|--|-----------------------------|------|-------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Plus | Lasso Qt./Acre |
| 1/2% to 3% Organic Matter | | | |
| Coarse ¹ (Sandy loam) | 1/3 | Plus | 1 1/2 to 2 |
| Medium ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 | Plus | 1 1/2 to 2 |
| Fine ² (Silty clay, silty clay loam ³ , clay, clay loam) | 2/3 | Plus | 2 |
| Mississippi Delta Only (Silty clay to heavy clay) | 1 1/3 | Plus | 2 to 2 1/2 |
| Greater than 3% Organic Matter | | | |
| Coarse ¹ (Sandy loam) | 1/2 | Plus | 1 1/2 to 2 |
| Medium ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 2/3 | Plus | 1 1/2 to 2 |
| Fine ² (Silty clay, silty clay loam ³ , clay, clay loam) | 2/3 to 5/6 | Plus | 2 to 2 1/2 |
| Mississippi Delta Only (Silty clay to heavy clay) | 1 1/3 | Plus | 2 to 2 1/2 |

¹ Do not use Metribuzin 75DF plus Lasso on sand or loamy sand soils with less than 2% organic matter.
² For control of lambsquarters, redroot pigweed, wild mustard, green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Metribuzin 75DF at rates of 1/3 lb./acre on medium soils and 1/3 to 1/2 lb./acre on fine soils regardless of soil organic matter percentage (use 1/2 lb. only where soil pH is less than 7.5 and weed pressure is heavy). The 1/3 lb./acre rate of Metribuzin 75DF in tank mix combination with Lasso can be applied regardless of soil pH. For control of other weeds use Metribuzin 75DF at full rates recommended in the table above, but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.
³ Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

PREPLANT INCORPORATED: For specific application information, refer to the GENERAL INFORMATION section in the front of this label.

Apply Metribuzin 75DF plus Lasso preplant incorporated if furrow irrigation is used or when a period of dry weather after application is expected. If

soybeans are planted on beds, apply and incorporate the tank mixture after bed formation. Apply within 7 days prior to planting and shallowly incorporate into the upper 1 to 2 inches of soil.

Do not use on muck soils.

| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Lasso Tank mix Preplant Incorporated Applications (Broadcast Rates) | | |
|---|-----------------------------|-------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Lasso Qt./Acre |
| Coarse ¹ (Loamy sand [over 2% organic matter], sandy loam) | 1/3 | 2 to 2 1/2 |
| Medium (Loam, silt loam, silt) | 1/2 | 2 1/2 to 3 |
| Fine (Silty clay loam ² , sandy clay loam, silty clay, sandy clay, clay loam, clay) | 2/3 | 2 1/2 to 3 |
| Mississippi Delta Only (Silty clay, clay) | 2/3 to 5/6 | 2 1/2 to 3 |

¹ Do not use Metribuzin 75DF plus Lasso on sand or loamy sand soils with less than 2% organic matter.
² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

SPECIAL PRECAUTIONS (Metribuzin 75DF Plus Lasso): For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the Lasso label.

METRIBUZIN 75DF PLUS COMMAND®

Metribuzin 75DF may be applied in combination with Command 4EC as a preplant or shallow incorporated application for the control of certain weeds in soybeans. Consult the Command 4EC label for specific directions on use, recommendations, restrictions, and any additional weeds not specified on this label.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: Metribuzin 75DF plus Command 4EC may only be applied with ground equipment as a preplant or shallow incorporated application. Metribuzin 75DF plus Command 4EC should be immediately incorporated into the top 1 to 3 inches after application unless surface is dry. On dry soils, incorporate into the top 1 to 3 inches within 3 hours of tank mix application.

Do not apply this tank mix within 1000 feet of towns and subdivisions, commercial vegetable, fruit, nurseries, or greenhouse operations.

A minimum of 15 gallons spray volume per acre is recommended with appropriate nozzle types and sizes to produce a coarse spray droplet. The use of an approved agricultural drift reducing additive is recommended for application volumes of 15 to 40 gallons per acre. The use of an approved agricultural drift reducing additive is required at spray volumes of 10 to 15 gallons per acre.

NOTE: Off-site movement of Command spray drift or vapors can cause foliar whitening or yellowing of some vegetation. Prior to application of Command, read and strictly follow all precautions and application instructions as set forth in that label.

For additional information on application, refer to the GENERAL INFORMATION section in the front of this label and the Command label.





| WEEDS CONTROLLED | | | |
|----------------------------|---|--------------------------|---------------|
| Annual Broadleaves: | | | |
| Bristly Starbur | Galinsoga | Prickly Sida/Teaweed | Smartweeds |
| Carpetweed | Jimsonweed | Purslane | Spurred Anoda |
| Copperleaf | Knotweed | Common Ragweed | Velvetleaf |
| Florida Beggarweed | Lambsquarters | Redweed | Venice Mallow |
| Florida Pusley | Pigweeds | Sesbania | Wild Mustard |
| Annual Grasses: | | | |
| Barnyardgrass* | Crabgrass* | Johnsongrass (seedling)* | Witchgrass |
| Bluegrass | Foxtails (Green, Giant, Yellow*, Robust Purple) | Fall Panicum* | |
| Broadleaf Signalgrass | Goosegrass | Texas Panicum | |

* Use 2 pt./Acre Command 4EC on coarse- and medium-textured soils with high populations of these weeds.

| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Command 4EC Tank mix Preplant Incorporated Application (Broadcast Rates) | | |
|---|-------------------------------|-------------------------|
| SOIL TEXTURE | Metribuzin 75DF (Lb./Acre) | Command 4EC Pt./Acre |
| 0.5% to 3% Organic Matter | | |
| Coarse ² (Sandy loam, loamy sand) | 1/3 | 1 1/2 to 2 |
| Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/3 to 1/2 | 1 1/2 to 2 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 1/3 to 1/2 | 1 1/2 to 2 |
| Over 3% Organic Matter | | |
| Coarse ² (Sandy loam, loamy sand) | 1/3 | 1 1/2 to 2 |
| Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/3 to 1/2 | 1 1/2 to 2 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 1/2 to 2/3 | 1 1/2 to 2 |
| ¹ Crop injury may occur on soils having a calcareous surface area or a pH of 7.1 or higher. ² Do not use on coarse soils with less than 1% organic matter. ³ Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. | | |

SPECIAL PRECAUTIONS (Metribuzin 75DF plus Command): Do not rotate to wheat, oats, barley, rye, alfalfa, or seed corn in the fall of the year of application or in the spring of the following year as crop injury may occur.

Do not apply when weather conditions favor drift. Do not use treated vines for feed or forage.

Observe all cautions and limitations on labeling of all products used in mixtures.

Do not apply aerially or through irrigation equipment.

METRIBUZIN 75DF PLUS COMMENCE®

Metribuzin 75DF Plus Commence Tank Mix Early Preplant Incorporated Application: Metribuzin 75DF in a tank mix with Commence 5.25 EC may be applied broadcast preplant incorporated up to 30 days before planting soybeans for the control of certain broadleaf weeds and grasses. Refer to the Commence herbicide label for additional directions for use, weeds controlled, recommendations, restrictions, and limitations not specified on this label.

Mixing: Refer to the GENERAL INFORMATION section on this label.

Application: For information on applying Metribuzin 75DF, refer to the GENERAL INFORMATION section on this label.





| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Commence 5.25 EC Tank mix Early Preplant Incorporated Application* | | |
|---|-----------------------------|------------------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Commence 5.25 EC Pt./Acre |
| 1/2% to 3% Organic Matter | | |
| Coarse* | 1/2 | 1 1/3 to 2 |
| Medium | 2/3 | 2 to 2 1/4 |
| Fine | 2/3 | 2 2/3 |
| Over 3% Organic Matter | | |
| Coarse* | 1/2 | 1 1/3 to 2 |
| Medium | 2/3 | 2 to 2 1/4 |
| Fine | 1 | 2 2/3 |
| ¹ For use on soils with a pH of 7.5 or lower. ² Do not use on coarse soils with less than 1% organic matter. | | |

Restrictions and Limitations: Do not apply aerially or through irrigation equipment.

Do not apply when weather conditions favor drift. Do not allow sprays to drift onto adjacent desirable plants. Do not use treated vines for feed or forage.

Do not rotate to wheat, barley, alfalfa, or seed corn in the fall of the year of application or in the spring of the following year as crop injury may occur.

Do not rotate to any crop not listed on this label for 18 months following application of Metribuzin 75DF.

Metribuzin 75DF plus Commence Tank Mix Preplant Incorporated: Metribuzin 75DF may be tank mixed with Commence 5.25 EC for preplant incorporated application to control certain weeds in soybeans. Refer to the GENERAL INFORMATION section of this label for information on mixing, application, restrictions, special precautions, and weeds controlled by Metribuzin 75DF. See appropriate sections of the Commence 5.25 EC herbicide label for additional precautionary statements, directions for use, recommendations, and additional weeds controlled.

| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Commence Tank Mix Preplant Incorporated Application (Broadcast Rates) | | |
|---|-----------------------------|-----------------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Commence 5.25EC Pt./Acre |
| 0.5% to 3% Organic Matter | | |
| Coarse² (Sandy loam, loamy sand) | 1/3 | 1 1/3 to 2 |
| Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/3 to 1/2 | 2 to 2 1/4 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 1/3 to 1/2 | 2 2/3 |
| Over 3% Organic Matter | | |
| Coarse² (Sandy loam, loamy sand) | 1/3 | 1 1/3 to 2 |
| Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/3 to 1/2 | 2 to 2 1/4 |
| Fine (Silty clay, silty clay loam ² , clay, clay loam) | 1/2 to 2/3 | 2 2/3 |
| ¹ Crop injury may occur on soils having a calcareous surface area or a pH of 7.1 or higher. ² Do not use on coarse soils with less than 1% organic matter. ³ Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. | | |

SPECIAL PRECAUTIONS (Metribuzin 75DF plus Commence): Do not rotate to wheat, barley, alfalfa, or seed corn in the fall of the year of application or in the spring of the following year as crop injury may occur.

Do not rotate to any crop not listed on this label for 18 months following application of Metribuzin 75DF.

Do not apply when weather conditions favor drift. Do not use treated vines for feed or forage.

Do not apply aerially or through irrigation equipment.

Do not allow sprays to drift onto adjacent desirable plants.

METRIBUZIN 75DF PLUS FREEDOM®

Metribuzin 75DF may be tank mixed with Freedom 3 EC for preplant incorporated application to control certain weeds in soybeans. Refer to the GENERAL INFORMATION section of this label for information on mixing, application, restrictions, special precautions, and weeds controlled by Metribuzin 75DF. See appropriate sections of the Freedom 3 EC herbicide label for additional precautionary statements, directions for use, recommendations, and additional weeds controlled.





| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Freedom Tank Mix Preplant Incorporated Application (Broadcast Rates) | | | |
|---|-----------------------------|------|--------------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Plus | Freedom 3 EC Qt./Acre |
| 1/2% to 3% Organic Matter | | | |
| Coarse ¹ (Sandy loam) | 1/3 | Plus | 2 3/4 to 3 1/2 |
| Medium ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 1/2 | Plus | 2 3/4 to 3 1/2 |
| Fine ² (Silty clay, silty clay loam ³ , clay, clay loam) | 2/3 | Plus | 3 1/2 to 4 |
| Mississippi Delta Only (Silty clay to heavy clay) | 1 1/3 | Plus | 3 1/2 to 4 1/2 |
| Greater than 3% Organic Matter | | | |
| Coarse ¹ (Sandy loam) | 1/2 | Plus | 3 to 3 1/2 |
| Medium ² (Loam, silt loam, silt, sandy clay, sandy clay loam) | 2/3 | Plus | 3 1/2 to 4 |
| Fine ² (Silty clay, silty clay loam ³ , clay, clay loam) | 2/3 to 5/6 | Plus | 3 1/2 to 4 1/2 |
| Mississippi Delta Only (Silty clay to heavy clay) | 1 1/3 | Plus | 3 1/2 to 4 1/2 |

¹ Do not use Metribuzin 75DF plus Freedom on sand or loamy sand soils with less than 2% organic matter.
² For control of lambsquarters, redroot pigweed, wild mustard, green and yellow foxtails on alkaline (calcareous) soils in Minnesota, Nebraska, South Dakota, and North Dakota only, apply Metribuzin 75DF at rates of 1/3 lb./acre on medium soils and 1/3 to 1/2 lb./acre on fine soils regardless of soil organic matter percentage (use 1/2 lb. only where soil pH is less than 7.5 and weed pressure is heavy). The 1/3 lb./acre rate of Metribuzin 75DF in tank mix combination with Freedom can be applied regardless of soil pH. For control of other weeds use Metribuzin 75DF at full rates recommended in the table above, but note that crop injury may occur on soils having a calcareous surface area or a pH of 7.5 or higher.
³ Silty clay loam soils are transitional soils and may be classified as medium - textured soils in some regions of the U.S.

Do not use on muck soils.

Do not allow sprays to drift onto adjacent desirable plants.

METRIBUZIN 75DF PLUS CANOPY® PLUS A GRASS HERBICIDE

A tank mix combination of Metribuzin 75DF plus Canopy 75 DF plus a registered and recommended grass herbicide (Dual, Lasso, Prowl, Sonalan, or Treflan) is recommended for control of the following weeds in soybeans:

| WEEDS CONTROLLED | | | |
|----------------------------|---------------|-------------------------|----------------|
| Annual Broadleaves: | | | |
| Bristly Starbur | Galinsoga | Prickly Sida/Teaweed | Shepherdspurse |
| Carpetweed | Jimsonweed | Purslane | Smartweed |
| Cocklebur | Knotweed | Ragweed, Common | Spurred Anoda |
| Copperleaf, Hophornbeam | Kochia | Redweed | Velvetleaf |
| Florida Beggarweed | Lambsquarters | Russian Thistle | Venice Mallow |
| Florida Pusley | Pigweed | Sesbania | Wild Mustard |
| Annual Grasses: | | | |
| Barnyardgrass | Crabgrass | Johnsongrass (seedling) | Sandbur |
| Bluegrass | Crowfootgrass | Junglerice | Sprangletop |
| Broadleaf Signalgrass | Foxtails | Panicum, Fall | Stinkgrass |
| Browntop Millet | Goosegrass | Panicum, Texas | |

Tank mix combinations which include Dual, Lasso, or Prowl can be applied broadcast or preplant incorporated broadcast. When Sonalan or Treflan are used in the tank mix, apply preplant incorporated broadcast. Refer to the table below for recommended rates of each product to be used in tank mix combinations:





| RECOMMENDED APPLICATIONS | | | |
|---|---------------------------------|-------------------------|-------------------------|
| Metribuzin 75DF Plus Canopy 75 DF Plus a Grass Herbicide (Broadcast Rates) | | | |
| PRODUCT | SOIL TEXTURE¹ | | |
| | Coarse² | Medium | Fine |
| Metribuzin 75DF (lb./acre) | 1/3 | 1/3 to 1/2 ³ | 1/2 to 2/3 ³ |
| Canopy DF (oz./acre) | 3 | 3 | 3 to 4 |
| Treflan (pt./acre) | 1 | 1 1/2 | 2 |
| Dual (pt./acre) | 1 1/4 to 1 1/2 | 1 1/2 to 2 | 2 to 2 1/2 |
| Prowl (pt./acre) | 1 1/2 | 1 1/2 to 2 | 1 1/2 to 2 1/2 |
| Lasso (qt./acre) | 2 to 2 1/2 | 2 1/2 to 3 | 2 1/2 to 3 |
| Sonalan (pt./acre) | 1 1/4 to 2 | 1 1/4 to 2 1/2 | 2 1/4 to 3 |

¹ Do not use on soils with a pH greater than 7.0.
² Refer to SOIL TEXTURE paragraph on this label for specific soil classification.
³ Use the lower rate of Metribuzin 75DF in preplant incorporated tank mix as in those situations where soils within a field vary extremely in texture or organic matter content.

IMPORTANT: If weeds escape in fields treated with these tank mix combinations, postemergence application of a registered and recommended herbicide will be needed for control.

SPECIAL PRECAUTIONS: For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of the labels for Metribuzin 75DF and Canopy 75 DF.

Do not use treated vines for feed or forage.

Refer to the GENERAL INFORMATION section of this label for mixing and application directions.

METRIBUZIN 75DF PLUS COMMAND PLUS A GRASS HERBICIDE

Metribuzin 75DF may be applied with Command 4EC and a grass herbicide (Treflan, Lasso, Dual, Prowl, or Sonalan) for the control of certain broadleaf weeds and grasses in soybeans. This combination will provide improved control of heavy infestations of velvetleaf, jimsonweed, and common ragweed. Metribuzin 75DF and Command 4EC plus a grass herbicide may be applied preplant incorporated broadcast. Consult the Command, Treflan, Lasso, Dual, Prowl, or Sonalan labels for specific directions for use, recommendations, restrictions, and additional weeds controlled not specified on this label.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: For specific application information, refer to the GENERAL INFORMATION section in the front of this label.

| WEEDS CONTROLLED | | | |
|----------------------------|----------------------|-------------------------|----------------|
| Annual Broadleaves: | | | |
| Bristly Starbur | Jimsonweed | Purslane | Sicklepod |
| Carpetweed | Knotweed | Ragweed, Common | Smartweeds |
| Copperleaf, Hophornbeam | Kochia | Redweed | Spotted spurge |
| Florida Beggarweed | Lambsquarters | Russian Thistle | Spurred Anoda |
| Florida Pusley | Pigweeds | Sesbania | Velvetleaf |
| Galinsoga | Prickly Sida/Teaweed | Shepherdspurse | Venice Mallow |
| | | | Wild Mustard |
| Annual Grasses: | | | |
| Barneyardgrass | Browntop Millet | Foxtails | Panicum, Fall |
| Bluegrass | Crabgrass | Goosegrass | Witchgrass |
| Broadleaf Signalgrass | Crowfootgrass | Johnsongrass (seedling) | |

Metribuzin 75DF and Command plus Treflan, Lasso, Dual, Prowl, or Sonalan will provide suppression (reduce the competition) of cocklebur and sunflower.





| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Command Plus a Grass Herbicide (Broadcast Rates) | | | |
|---|---------------------------|-------------------------|-------------------------|
| PRODUCT | SOIL TEXTURE ¹ | | |
| | Coarse | Medium | Fine |
| Metribuzin 75DF (lb/acre) | 1/3 | 1/3 to 1/2 ² | 1/2 to 2/3 ² |
| Command 4EC ³ (pt/acre) | 1/2 to 3/4 | 1/2 to 3/4 | 1/2 to 3/4 |
| Treflan (pt/acre) | 1 | 1 1/2 | 2 |
| Dual (pt/acre) | 1 1/4 to 1 1/2 | 1 1/2 to 2 | 2 to 2 1/2 |
| Prowl (pt/acre) | 1 1/2 | 1 1/2 to 2 | 1 1/2 to 2 1/2 |
| Lasso (qt/acre) | 2 to 2 1/2 | 2 1/2 to 3 | 2 1/2 to 3 |
| Sonalan (pt/acre) | 1 1/4 to 2 | 1 3/4 to 2 1/2 | 2 1/4 to 3 |

¹ Refer to SOIL TEXTURE paragraph on this label for specific soil classification. On coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on loamy sand or sandy loam soils with less than 1% organic matter.

² The higher rate of Metribuzin 75DF is recommended for the control of sicklepod and hemp sesbania. Use the lower rate of Metribuzin 75DF in the preplant incorporated tank mix on soils having a calcareous surface area or a pH of 7.5 or higher, and in those situations where soils within a field vary extremely in texture or organic matter content.

³ Higher rate is recommended under moderate to heavy weed infestations.

METRIBUZIN 75DF PLUS SCEPTER PLUS A GRASS HERBICIDE

Metribuzin 75DF may be applied with Scepter herbicide and a grass herbicide (Treflan, Lasso, Dual, Prowl, or Sonalan) for the control of certain broadleaf weeds and grasses in soybeans. Metribuzin 75DF and Scepter plus Treflan or Sonalan may be applied preplant incorporated broadcast. Metribuzin 75DF and Scepter plus Lasso, Dual, or Prowl may be applied preplant incorporated, preemergence broadcast, or in a band application.

Consult the Scepter, Treflan, Lasso, Dual, Prowl, or Sonalan labels for specific

directions for use, recommendations, restrictions, and additional weeds controlled not specified on this label.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: For specific application information, refer to the GENERAL INFORMATION section in the front of this label.

Weeds Controlled: Metribuzin 75DF plus Scepter plus Treflan, Lasso, Dual, Prowl, or Sonalan will control the following broadleaf weeds and grasses:

| WEEDS CONTROLLED | | | |
|----------------------------|---------------------------|-------------------------|----------------|
| Annual Broadleaves: | | | |
| Bristly Starbur | Galinsoga | Prickly Sida/Teaweed | Smartweeds |
| Buffalobur | Jimsonweed | Purslane | Spotted Spurge |
| Carpetweed | Knotweed | Ragweed, Common | Spurred Anoda |
| Cocklebur | Kochia | Redweed | Sunflower |
| Coffee Senna | Lambsquarters | Russian Thistle | Velvetleaf |
| Copperleaf, Hophornbeam | Morningglory, pitted | Sesbania | Venice Mallow |
| Florida Beggarweed | Morningglory, smallflower | Shepherdspurse | Wild Mustard |
| Florida Pusley | Pigweeds | Sicklepod | |
| Annual Grasses: | | | |
| Barnyardgrass | Browntop Millet | Foxtails | Panicum, Fall |
| Bluegrass | Crabgrass | Goosegrass | Witchgrass |
| Broadleaf Signalgrass | Crowfootgrass | Johnsongrass (seedling) | |

Metribuzin 75DF and Scepter plus Treflan, Lasso, Dual, Prowl, or Sonalan will suppress (reduce the competition) of ivyleaf and tall morningglory and red rice.





| Metribuzin 75DF Plus Scepter Plus a Grass Herbicide (Broadcast Rates) | | | |
|---|---------------------------|-------------------------|-------------------------|
| PRODUCT | SOIL TEXTURE ¹ | | |
| | Coarse | Medium | Fine |
| Metribuzin 75DF (lb./acre) | 1/3 | 1/3 to 1/2 ² | 1/2 to 2/3 ² |
| Scepter (1.5 lb./gal liquid ³ pt./acre) | 1/3 to 1/2 | 1/3 to 1/2 | 1/3 to 1/2 |
| -or- Scepter 70 DG ³ (oz./acre) | 1.4 to 2.1 | 1.4 to 2.1 | 1.4 to 2.1 |
| Treflan (pt./acre) | 1 | 1 1/2 | 2 |
| Dual (pt./acre) | 1 1/4 to 1 1/2 | 1 1/2 to 2 | 2 to 2 1/2 |
| Prowl (pt./acre) | 1 1/2 | 1 1/2 to 2 | 1 1/2 to 2 1/2 |
| Lasso (qt./acre) | 2 to 2 1/2 | 2 1/2 to 3 | 2 1/2 to 3 |
| Sonalan (pt./acre) | 1 1/4 to 2 | 1 3/4 to 2 1/2 | 2 1/4 to 3 |

¹ Refer to SOIL TEXTURE paragraph on this label for specific soil classification. On coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on loamy sand or sandy loam soils with less than 1% organic matter.

² The higher rate of Metribuzin 75DF is recommended for preemergence tank mix application and for the control of sicklepod and hemp sesbania. Use the lower rate of Metribuzin 75DF in the preplant incorporated tank mix on soils having a calcareous surface area or a pH of 7.5 or higher and in those situations where soils within a field vary extremely in texture or organic matter content.

³ Higher rate is recommended under moderate to heavy weed infestations.

METRIBUZIN 75DF PLUS PURSUIT® AND A GRASS HERBICIDE

Metribuzin 75DF may be tank mixed with Pursuit herbicide and a registered and recommended grass herbicide (Dual, Lasso, Prowl, Sonalan, or Treflan) for control of certain broadleaf and grass weeds in soybeans. Refer to the product labels for Pursuit, Dual, Lasso, Prowl, Sonalan, or Treflan for additional directions for use, recommendations, restrictions, and limitations not included on this label.

Tank mix combinations of Metribuzin 75DF, Pursuit and Dual, Lasso or Prowl can be applied broadcast preemergence or preplant incorporated. When the grass herbicide used is Sonalan or Treflan, apply the tank mix broadcast preplant incorporated.

Mixing and Application: Refer to the GENERAL INFORMATION section of this label for directions on mixing and application of Metribuzin 75DF.

| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Pursuit and a Grass Herbicide* | | |
|---|--------------------------|------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Pursuit Oz./Acre |
| Coarse | 1/3 | 4 |
| Medium | 2/5 to 1/2 | 4 |
| Fine | 1/2 to 2/3 | 4 |

* For control of grass weeds include Dual, Lasso, Prowl, Sonalan, or Treflan at label rates in the tank mix with Metribuzin 75DF and Pursuit herbicides.

Restrictions and Limitations: Do not apply this tank mix with aerial or irrigation equipment. Do not apply when weather conditions favor drift or allow sprays to drift onto adjacent desirable plants. Do not use treated vines for feed or forage. Refer to appropriate sections of the Pursuit herbicide label for restrictions on use area and rotational crops. Observe all cautions and limitations on the labeling of all products used in mixtures.

METRIBUZIN 75DF PLUS PURSUIT PLUS HERBICIDE

Metribuzin 75DF may be tank mixed with Pursuit Plus herbicide for broadcast preemergence or preplant incorporated application to soybeans for control of certain broadleaf and grass weeds. Refer to the Pursuit Plus herbicide label for additional directions for use, recommendations, restrictions, and limitations not included on this label.

Mixing and Application: Refer to the GENERAL INFORMATION section of this label for directions on mixing and application of Metribuzin 75DF.

| RECOMMENDED APPLICATIONS Metribuzin 75DF Plus Pursuit Plus Herbicide | | |
|---|--------------------------|------------------|
| SOIL TEXTURE | Metribuzin 75DF Lb./Acre | Pursuit Oz./Acre |
| Coarse | 1/3 | 2 1/2 |
| Medium | 2/5 to 1/2 | 2 1/2 |
| Fine | 1/2 to 2/3 | 2 1/2 |

Restrictions and Limitations: Do not apply this tank mix with aerial or irrigation equipment. Do not apply when weather conditions favor drift or allow sprays to drift onto desirable plants.

Do not use treated vines for feed or forage.

Refer to appropriate sections of the Pursuit Plus herbicide label for restrictions on use area and rotational crops.



**METRIBUZIN 75DF PLUS LINURON PLUS LASSO or DUAL**

Metribuzin 75DF plus Linuron plus Lasso or Dual Tank Mix Application: Metribuzin 75DF may be applied in combination with Linuron 50 DF or 4L and Lasso 4 or Dual 8 EC as a preemergence application for the control of certain weeds in soybeans. Consult the linuron, Lasso, or Dual labels for specific directions for use, recommendations, restrictions, and any additional weeds not specified on this label.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: Applications can be made only with ground spray equipment in accordance with specified soil types and dosage rates. For specific application information, refer to the GENERAL INFORMATION section in the front of this label.

| Metribuzin 75DF Plus Linuron Plus Lasso or Dual Broadcast Rates (0.5 to 3% Organic Matter Only) | | | |
|--|---|--|--|
| PRODUCT | SOIL TEXTURE | | |
| | Coarse¹ (Sandy, loamy sand, sandy loam) | Medium (Loam, silt loam, silt, sandy clay, sandy clay loam) | Fine (Silty clay, silty clay loam ² , clay, clay loam) |
| Metribuzin 75DF (lb./acre) | 1/6 to 1/4 | 1/4 to 1/3 | 1/3 to 1/2 |
| Linuron 50 DF (lb./acre) -or- Linuron 4L (pt./acre) | 1/3 to 1/2 | 1/2 to 3/4 | 3/4 to 1 1/2 |
| Lasso 4 (qt./acre) -or- Dual 8 EC (pt./acre) | 3/4 to 1 | 1 to 1 1/2 | 1 1/4 to 2 |
| | 1 to 1 1/4 | 1 1/4 to 1 1/2 | 1 1/2 to 2 |

¹ Do not use Metribuzin 75DF plus linuron plus Lasso or Dual on sand soils with less than 1% organic matter.

² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

SPECIAL PRECAUTIONS (Metribuzin 75DF plus linuron plus Lasso or Dual: For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the linuron label and the Lasso or Dual labels.

FOR USE IN COARSE (LIGHT) SOILS in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia.

Metribuzin 75DF herbicide is recommended alone or in combination with

Treflan, Lasso, or Dual for use in coarse-textured, low organic matter soils in the states listed above for the control of certain weeds in soybeans. Refer to the appropriate sections of this label and the Treflan, Lasso, or Dual label for specific directions for use, recommendations, restrictions, and any additional weeds not specified on this label.

Mixing: Refer to the GENERAL INFORMATION section in the front of this label.

Application: For specific application information, refer to the GENERAL INFORMATION section in the front of this label.

| Metribuzin 75DF (Alone) Preemergence Application (Broadcast Rates) | | |
|---|-----------------------|---------------------------------|
| SOIL TEXTURE | ORGANIC MATTER | METRIBUZIN 75DF lb./acre |
| Coarse (Light) Soils-Sand¹, Loamy Sand, Sandy Loam | 0.5% or Above | 1/3 to 1/2 ² |

¹ Not recommended for use on sand with less than 1% organic matter.

² Use the higher rate under heavy weed pressures and/or on soils higher in organic matter.

Metribuzin 75DF in Combination with Other Herbicides: Metribuzin 75DF is recommended in a tank mix combination with Treflan as a preplant incorporated application or as a preemergence overlay application following

a preplant incorporated application of Treflan. Metribuzin 75DF is also recommended for use as a preemergence application in combination with Lasso or Dual.

| FOR USE IN COARSE (Light) SOILS 0.5% or Above Organic Matter (Broadcast Rates) | | | |
|---|---|-------------|---------------------------------|
| Soil Texture | Combination Product/Acre | Plus | Metribuzin 75DF lb./acre |
| Coarse (Light) Soils- Sand ¹ , Loamy sand, Sandy loam | Preplant Incorporated Treflan 4EC 1 pt. | Plus | 1/3 to 1/2 ² |
| | Preemergence Lasso 4E 1 1/2 to 2 qt. Dual 8E 1 1/4 to 1 1/2 pt. | Plus | 1/3 to 1/2 ² |

¹ Not recommended for use on sand with less than 1% organic matter.

² Use the higher rate under heavy weed pressures and/or on soils higher in organic matter.

SPECIAL PRECAUTIONS: Do not use on sand soils with less than 1% organic matter or on sandy loam or loamy sand soils with less than 0.5% organic matter.

For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label and the Treflan, Lasso, Dual, Surlflan, or Amiben labels.





BURNDOWN WEED CONTROL-Field Corn and Soybeans

Metribuzin 75DF can be used as part of an herbicide program for burndown of existing vegetation prior to crop emergence in conservation tillage systems. Metribuzin 75DF may be tank mixed with 2,4-D low volatile ester (LVE), Gramoxone Extra, or Roundup/Roundup Ultra/Touchdown for control of emerged weeds prior to field corn or soybean emergence. Metribuzin 75DF tank mixes with 2,4-DB, Fusion, Poast Plus, or Select may also be used in soybeans for control of emerged weeds prior to crop emergence. Metribuzin 75DF burndown tank mixes can be applied before planting or prior to crop emergence in the following areas:

Field Corn: Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota, and Wisconsin.

Soybeans: All areas for all products except Fusion tank mixes-see Fusion section of this label for recommended states.

Application: Metribuzin 75DF may be applied up to 30 days prior to planting or preemergence. Apply only by ground equipment when Metribuzin 75DF is used for burndown of existing vegetation in conservation tillage systems. Metribuzin 75DF and tank mix partner burndown rates are listed in the following three tables.

| METRIBUZIN 75 DF BURNDOWN RATES Field Corn and Soybeans | | |
|---|--------------------------|------------------------------|
| Crops | Application Timing | Metribuzin 75DF Rate (oz./A) |
| Field Corn: Iowa, Kansas, Missouri, Nebraska, South Dakota | Preplant (0 to 30 days) | 2 to 5 1/3 |
| | Preemergence | |
| Field corn: Illinois, Indiana, Kentucky, Michigan, Minnesota, Ohio, Wisconsin | Preplant (10 to 30 days) | 2 to 5 1/3 |
| | Preplant (0 to 9 days) | 2 to 4 |
| | Preemergence | |
| Soybeans | Preplant (0 to 30 days) | 2 to 5 1/3 |
| | Preemergence | |

SPECIAL PRECAUTIONS: Do not apply these treatments after crop emergence. Observe all precautions and limitations on the labeling of all products used in tank mixtures. Refer to the GENERAL INFORMATION section of this label for additional information, precautions, and limitations.

Field Corn:

- Do not apply on coarse-textured soils with less than 1.5% organic matter.
- Do not apply more than 4 oz. of Metribuzin 75DF per acre on soils with less than 2% organic matter.
- Do not apply on soils having pH 7.0 or greater.
- Do not apply more than 5 1/3 oz. Metribuzin 75DF (0.25 pound active ingredient) per acre per growing season.
- Corn seed should be planted a minimum of 1 1/2 inches deep.
- Metribuzin 75DF may only be used in hybrid seed corn production fields if both inbred parents are known to be tolerant to Metribuzin 75DF.

Soybeans:

- Apply only 2,4-D low volatile ester formulations which are registered and recommended for preplant or burndown use in soybeans.
- Do not apply tank mixture containing 2,4-D LVE if wind is blowing toward desired susceptible plants (i.e. cotton, tobacco, tomato, etc.) or when wind speeds exceed 6 miles per hour.

Feeding Restrictions: Corn treated with Metribuzin 75DF may be harvested for silage or grain 60 days after treatment. Soybean vines or hay treated with Metribuzin 75DF may be grazed or fed to livestock 40 days after application. Do not feed hay, forage, fodder, or graze 2,4-D-, Select-, or Fusion-treated vegetation. Follow the most restrictive preharvest interval of all products used in a tank mixture.

| METRIBUZIN 75DF PLUS TANK MIX PARTNER BURNDOWN RATES-Field Corn or Soybeans | | |
|---|--|---|
| Product | Rate | Directions and Remarks |
| Metribuzin 75DF + 2,4-D LVE | 2 to 5 1/3 oz./A* + 1/4 to 1 lb. ai/A | In soybeans, apply at least 7 days preplant when using 2,4-D LVE at 1/4 to 1/2 lb. ai/A and at least 30 days preplant with rates greater than 1/2 lb. ai/A. Include crop oil concentrate (COC) at the rate of 1 gal./100 gal of spray solution (1% v/v). In corn, apply at least 7 days preplant or at least 3 days after planting but before corn emergence. |
| Metribuzin 75DF + Gramoxone Extra | 2 to 5 1/3 oz./A* + 24 to 48 fl. oz./A | Must be applied prior to crop emergence. Use 24 to 32 fluid ounces of Gramoxone Extra for weeds less than 4 inches in height and 32 to 48 fluid ounces when weeds are 4 to 6 inches in height. Apply in 20 to 60 gallons of water per acre. Include either nonionic surfactant at 1 quart per 100 gallons (0.25% v/v) or crop oil concentrate at 1 gallon per 100 gallons (1% v/v) of spray solution. |
| Metribuzin 75DF + Gramoxone Extra + 2,4-D LVE | 2 to 5 1/3 oz./A* + 24 to 48 fl. oz./A + 1/4 to 1 lb. ai/A | For this tank mix follow the Directions and Remarks section above for Metribuzin 75DF + 2,4-D LVE and Metribuzin 75DF + Gramoxone Extra, paying special attention to crop planting restrictions with 2,4-D LVE. Include either nonionic surfactant or crop oil concentrate in this tank mix. |





| Product | Rate | Directions and Remarks |
|---|---|--|
| Metribuzin 75DF + Roundup/Roundup Ultra or Touchdown | 2 to 5 1/3 oz./A* + 12 to 24 fl. oz./A or 8 to 16 fl. oz./A | Must be applied prior to crop emergence. Use the higher rates as weeds approach the maximum weed heights listed in the WEEDS CONTROLLED section below. Apply in 10 to 20 gallons of water per acre. With Roundup and Touchdown, include nonionic surfactant at 2 quarts per 100 gallons (0.5% v/v) and ammonium sulfate (spray grade) at 17 pounds per 100 gallons of spray solution. With Roundup Ultra, include ammonium sulfate (spray grade) at 17 pounds per 100 gallons of spray solution. Any glyphosate formulation registered and labeled for use in field corn or soybeans may be tank mixed with Metribuzin 75DF. |
| Metribuzin 75DF + Roundup/Roundup Ultra/ Touchdown + 2,4-D LVE | 2 to 5 1/3 oz./A* + 12 to 24 fl. oz./A or 8 to 16 fl. oz./A + 1/4 to 1 lb. ai/A | For this tank mix follow the Directions and Remarks section above for Metribuzin 75DF + 2,4-D LVE and Metribuzin 75DF + Roundup/Roundup Ultra/Touchdown, paying special attention to planting restrictions with 2,4-D LVE. Use the adjuvant recommendations under the Metribuzin 75DF + Roundup/Roundup Ultra/Touchdown tank mix. Do not use crop oil concentrate. |

* If applied to field corn grown in Illinois, Indiana, Kentucky, Michigan, Minnesota, Ohio, and Wisconsin, refer to Table 1 for correct Metribuzin 75DF rate based on application timing.

METRIBUZIN 75DF PLUS TANK MIX PARTNER BURNDOWN RATES- Soybeans Only

| Product | Rate | Directions and Remarks |
|--|---|--|
| Metribuzin 75DF + 2,4-D LVE | 2 to 5 1/3 oz./A* + 1/8 to 7/32 lb. ai/A | Apply preplant or before soybean emergence. Include nonionic surfactant at 2 quarts per 100 gallons (0.5% v/v) of spray solution. |
| Metribuzin 75DF + Fusion + 2,4-D LVE | 2 to 5 1/3 oz./A* + 4 to 8 fl. oz./A + 1/4 to 1 lb. ai/A | For use only in Delaware, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Pennsylvania, South Dakota, Virginia, West Virginia, and Wisconsin. For this tank mix, follow the planting restrictions under the Directions and Remarks section above for Metribuzin 75DF + 2,4-D LVE. Fusion rates of 4, 6 and 8 fl. ounces will control certain grasses up to 2, 4 and 6 inches in height, respectively. Include either crop oil concentrate at 1 gallon per 100 gallons (1.0% v/v) or nonionic surfactant at 1 to 2 quarts per 100 gallons (0.25 to 0.5% v/v) of spray solution. Refer to the Fusion label for additional information. |
| Metribuzin 75DF + Poast Plus + 2,4-D LVE | 2 to 5 1/3 oz./A* + 8 to 16 fl. oz./A + 1/4 to 1 lb. ai/A | For this tank mix follow the planting restrictions under the Directions and Remarks section above for Metribuzin 75DF + 2,4-D LVE. The 8 and 12 fl. oz. rate of Poast Plus will control certain grasses up to 2 and 3 inches in height, respectively. Include either crop oil concentrate at the rate of 1 gallon per 100 gallons of spray solution (1% v/v) or Dash HC at 1 pint per acre. Refer to the Poast Plus label for additional information. |
| Metribuzin 75DF + Select + 2,4-D LVE | 2 to 5 1/3 oz./A* + 3 to 4 fl. oz./A + 1/4 to 1 lb. ai/A | For this tank mix follow the planting restrictions under the Directions and Remarks section above for Metribuzin 75DF + 2,4-D LVE. The 3 and 4 fluid ounce rates of Metribuzin 75DF will control certain grasses up to 3 and 4 inches in height, respectively. Include crop oil concentrate at the rate of 1 quart per acre and 28% UAN (urea ammonium nitrate) at a rate of 1 to 2 quarts per acre. Refer to the Select label for additional information. |

Weeds controlled: Metribuzin 75DF in tank mixtures with the above herbicides will provide burndown control of the weeds listed below.





| WEEDS CONTROLLED BY BURNDOWN RATES OF METRIBUZIN 75DF | | | | | | | | | |
|---|--------------------------------|------------------------|--------------------|--------------------|-----------------------------------|---|--------------------|-----------------------------|--------------------------------|
| Weeds Controlled | METRIBUZIN 75DF | | | | | | | | |
| | 2,4-D LVE | Poast Plus + 2,4-D LVE | Select + 2,4-D LVE | Fusion + 2,4-D LVE | Roundup/ Roundup Ultra/ Touchdown | Roundup/ Roundup Ultra/ Touchdown + 2,4-D LVE | Gramoxone Extra | Gramoxone Extra + 2,4-D LVE | 2,4-DB |
| ANNUAL GRASSES | | | | | | | | | |
| MAXIMUM BURNDOWN HEIGHT (INCHES) | | | | | | | | | |
| Barley | | - | - | - | 8 | | 4 to 6 | | |
| Barnyardgrass | Does not control these species | 2 to 3 | 3 to 4 | - | 6 | | 4 to 6 | | Does not control these species |
| Crabgrass spp. | | 2 to 3 | - | - | 6 | | 4 to 6 | | |
| Foxtail spp. | | 2 to 3 | 3 to 4 | 2 to 6 | 8 | | 4 to 6 | | |
| Johnsongrass, seedling | | 2 to 3 | - | - | 8 | | 4 to 6 | | |
| Panicum, fall | | 2 to 3 | 3 | 2 to 6 | 6 | | 4 to 6 | | |
| Sandbur, field | | - | - | - | 8 | | 4 to 6 | | |
| Shattercane | | 2 to 3 | - | - | 8 | | 4 to 6 | | |
| Wheat, volunteer | | - | - | - | 6 | | 4 to 6 | | |
| Witchgrass | | 2 to 3 | - | - | 6 | | 4 to 6 | | |
| BROADLEAVES | | | | | | | | | |
| MAXIMUM BURNDOWN HEIGHT (INCHES) | | | | | | | | | |
| Buffalobur | | - | - | - | 6 | 6 | 4 to 6 | 4 to 6 | - |
| Chickweed, common | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 2 |
| Cocklebur, common | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 6 |
| Dandelion, common | | 6 dia ^a | | | 2 dia ^b | 6 dia ^a | 4 dia ^d | 6 dia ^a | 2 dia |
| Henbit | | 4 | | | 4 | 4 | 4 to 6 | 4 to 6 | - |
| Horseweed/marestail | | 6 ^{ac} | | | 4 ^b | 6 | 3 | 6 ^a | 2 ^c |
| Jimsonweed | | 6 | | | 6 | 6 | 4 to 6 | 4 to 6 | 2 |
| Kochia* | | 4 ^{ac} | | | 4 | 4 | 4 | 4 | - |
| Ladysthumb | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Lambsquarters, common | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 2 |
| Lettuce, prickly | | 6 | | | 4 | 6 | 4 to 6 | 4 to 6 | 2 |
| Mallow, Venice | | 6 | | | 6 | 6 | 4 to 6 | 4 to 6 | - |
| Morningglory spp. | | 6 | | | 2 | 4 | 2 | 4 | 4 |
| Mustard spp. | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 2 |
| Pennycress, field | | 6 | | | 6 | 6 | 4 to 6 | 4 to 6 | 2 |
| Pigweed, spp. (annual) | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Ragweed, common | | 6 | | | 6 ^b | 8 | 4 to 6 | 4 to 6 | 2 |
| Ragweed, giant | | 6 ^{ac} | | | 4 ^b | 6 | 4 | 6 | 2 |
| Shepherdspurse | | 6 | | | 6 | 6 | 4 to 6 | 4 to 6 | - |
| Sida, prickly | | 6 | | | 4 | 4 | 4 | 4 | 1 |
| Smartweed, Pennsylvania | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Sunflower, common | | 6 | | | 6 | 6 | 4 to 6 | 4 to 6 | 4 |
| Thistle, Russian | | 4 ^{ac} | | | 2 to 4 ^{bc} | 6 | 4 | 4 to 6 | 3 ^c |
| Velvetleaf | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |
| Waterhemp spp. | | 6 | | | 6 | 8 | 4 to 6 | 4 to 6 | 3 |

^a Use 2,4-D LVE at 0.5 pound active ingredient per acre.
^b Use a minimum Roundup/Roundup Ultra rate of 16 fl. oz./A and a minimum Touchdown rate of 10.6 fl. oz./A.
^c Use Metribuzin 75DF at 4 oz./A for optimum control.
^d Suppression only.
* Does not control triazine-resistant biotypes.



RESIDUAL WEED CONTROL

Metribuzin 75DF burndown programs can be used as part of a full season weed control program in both field corn and soybeans when, (1) applied as a tank mixture with residual herbicides, or (2) followed with a postemergence weed control program which is registered for use on the crop.

For residual control, Metribuzin 75DF burndown programs may include tank mixes with the following herbicides or combination of herbicides:

| Field Corn | | | |
|--------------------|--------------|---------------------------|-----------------|
| Alachlor | Bullet | Linex | Ramrod/Atrazine |
| Atrazine | Clarity | Linuron | Simazine |
| Banvel | Dual | Lorox | Surpass |
| Bicep | Dual II | Marksman | Surpass 100 |
| Bicep II | Frontier | Pentagon | Top Notch |
| Bicep Lite | Guardsman | Prowl | |
| Bladex | Harness | Pursuit ^a | |
| Broadstrike + Dual | Harness Xtra | Pursuit Plus ^a | |
| Broadstrike Plus | Lariat | Ramrod | |

^aUse only Pursuit resistant/tolerant corn hybrids.

| Soybeans | | | |
|--------------------|----------|----------------|---------------------|
| Alachlor | Dual | New Lorox Plus | Pursuit Plus |
| Broadstrike + Dual | Dual II | Pentagon | Scepter |
| Canopy | Frontier | Preview | Sencor ^b |
| Command | Gemini | Prowl | Squadron |
| Detail | Linuron | Pursuit | Turbo |

^b Metribuzin 75DF used (alone and in tank mixes) on soybeans at higher labeled rates than those listed for burndown weed control will also provide residual control of those weeds listed in the WEEDS CONTROLLED by Metribuzin 75DF and Metribuzin 75DF TANK MIX COMBINATIONS section of the Metribuzin 75DF label.

Refer to the individual product labels for additional information, precautions, and limitations.

SOUTHERN AND SOUTHEASTERN STATES ONLY**Postemergence Directed Spray Applications**

Metribuzin 75DF can be applied in postemergence directed sprays to soybeans for control of certain weeds which escape preplant or preemergence herbicide applications and for control of additional flushes of weeds that may occur after soybeans have emerged. Postemergence directed sprays of Metribuzin 75DF can be applied to soybeans in addition to a preemergence or preplant application of Metribuzin 75DF herbicide according to label directions.

Weeds Controlled: Metribuzin 75DF applied postemergence to soybeans as a directed spray according to directions on this label will control the following at rates shown (broadcast basis) when grasses and common ragweed are less than 1 inch tall and other broadleaves are less than 3 inches tall:

1/3 Lb./Acre

Florida beggarweed (*Desmodium tortuosum*)
Pigweeds (*Amaranthus* spp.)
Carpetweed (*Mollugo verticillata*)
Cocklebur (*Xanthium pensylvanicum*)
Dayflower (*Commelina* spp.)
Mexicanweed (*Caperonia castaniifolia*)
Purslane (*Portulaca oleracea*)
Sicklepod (*Dassia obtusifolia*)
Velvetleaf (*Abutilion theophrasti*)
Crabgrass (*Digitaria* spp.)

1/3 to 2/3 Lb./Acre

Sesbania (*Sesbania* spp.)
Prickly sida/Teaweed (*Sida spinosa*)

2/3 Lb./Acre

Ragweed, common (*Ambrosia artemisiifolia*)

At the rate of 2/3 lb./acre morningglory species (*Ipomoea* spp.), horsenettle (*Solanum* spp.), Florida pusley (*Richardia acabra*), spotted spurge (*Euphorbia maculata*), and wild poinsettia (*Euphorbia heterophylla*) are suppressed when Metribuzin 75DF is applied before these weeds are 3 inches tall. The 2/3 lb./acre rate will suppress broadleaf signal grass (*Brachifolia platyphylla*) up to 1 inch tall.



| METRIBUZIN 75DF POSTEMERGENCE DIRECTED SPRAY | |
|---|---------------------------------|
| RECOMMENDED APPLICATIONS | |
| CROP | Metribuzin 75DF Lb./Acre |
| Soybeans (Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, and Texas) | 1/3 to 2/3 (Broadcast Basis) |
| Apply proper dosage using 10 to 40 gallons of water per acre as a directed spray in a 6- to 8-inch band on each side of the row after soybeans are 8 inches tall and before broadleaf weeds are 3 inches tall and before grasses and common ragweed are 1 inch tall. For best results, the spray must cover weed foliage with minimum or no contact with soybean foliage. Add a nonionic surfactant such as Ortho X-77 to the spray mixture to obtain better wetting of weed leaf surfaces. To determine the correct dosage of Metribuzin 75DF for a band application, see BANDED APPLICATION under the GENERAL INFORMATION section in the front of this label. | |
| If necessary, a second postemergence directed spray application can be made after 7 days. | |
| Do not feed or graze green soybean vines. Do not harvest soybeans or use dry soybean vines for feed or forage within 70 days of last application. | |

SPECIAL PRECAUTIONS (Directed Postemergence): Do not apply directly to soybeans or serious crop injury will occur. Do not allow spray to contact more than the lower 1/4 to 1/3 of soybean plants. Soybean leaves contacted by the spray will be killed.

Do not apply Metribuzin 75DF postemergence to sensitive soybean varieties. See SPECIAL PRECAUTIONS in the front of this label.

To avoid injury to other crops or desirable plants from spray drift, sprayer pressure must not exceed 30 psi and the sprayer must be fitted with nozzles no smaller than 8002-T-Jet (or equivalent). Do not apply under weather conditions which favor drift.

POTATOES

Metribuzin 75DF herbicide is recommended for use in ground, aircraft, or specified chemigation equipment as a preemergence and/or postemergence application to potatoes. Early maturing smooth skinned white and all red skinned varieties may be injured with postemergence applications. The varieties Atlantic, Bellchip, Centennial, Chipbelle, and Shepody are sensitive to Metribuzin 75DF. Avoid postemergence applications on these varieties.

Preemergence applications on these varieties may cause crop injury under adverse weather conditions on coarse soils, under high soil pH, with higher rates per acre, and with mechanical incorporation.

Ground Application: Metribuzin 75DF is recommended for use with ground spray equipment applied as a preemergence and/or postemergence application for control of the listed grass and broadleaf weeds in potatoes. Apply as a uniform broadcast spray at 20 or more gallons per acre.

Aerial Application: Metribuzin 75DF may be applied in aerial spray equipment as a preemergence and/or postemergence application at 5 or more gallons per acre.

Chemigation: Metribuzin 75DF may be applied preemergence and/or early postemergence to potatoes using center pivot, solid set, and lateral roll systems. Apply specified dosage in 1/4 to 3/4 inch of water per acre (1/4 to 1/2 inch on sandy soil) as a continuous injection in self propelled systems or apply in the last 15 to 30 minutes of the set in other systems. Be sure all the Metribuzin 75DF has been flushed from the lines before shutting down the system.

WEEDS CONTROLLED

Metribuzin 75DF applied to potatoes according to directions will provide economic control of the following weeds. For optimum control, applications should be made before weeds are 1 inch tall. (See NOTE)

| Broadleaves: | | | |
|--------------------------------------|----------------------------------|-------------------------------------|--|
| Carpentweed, common ¹ | Mustard, Indian ¹ | Pigweed, redroot ^{1,2} | Sicklepod ¹ |
| Cocklebur, common ^{1,2} | Mustard, tansy ¹ | Pigweed, smooth ^{1,2} | Smartweed, Pennsylvania ^{1,2} |
| Jimsonweed ¹ | Mustard, tumble ¹ | Ragweed, common ^{1,2} | Sunflower, common ³ |
| Kochia ¹ | Mustard, wild ¹ | Shepherdspurse ¹ | Thistle, Russian ² |
| Lambsquarters, common ^{1,2} | Pennycress, field ^{1,2} | | |
| Grasses: | | | |
| Barnyardgrass ³ | Foxtail, giant ¹ | Johnsongrass, seedling ¹ | Signalgrass, broadleaf ¹ |
| Crabgrass, large ¹ | Foxtail, green ¹ | Panicum, fall ¹ | |
| Crabgrass, smooth ¹ | Foxtail, yellow ¹ | | |

¹ Weeds controlled with preemergence applications.

² Weeds controlled with postemergence applications.

³ Weeds requiring two applications for control.



HARD TO CONTROL WEEDS

Although Metribuzin 75DF may not provide commercially acceptable control in every instance, it will suppress growth of the following weeds and reduce their competition with potato plants.

| Broadleaves | Grasses |
|-------------------|------------------|
| Kochia | Barnyardgrass |
| Nightshade, hairy | Nutsedge, yellow |
| Purslane, common | |
| Sunflower, common | |

NOTE: Where triazine-resistant weeds are present, Metribuzin 75DF alone may not provide adequate control.

RECOMMENDED BROADCAST APPLICATIONS FOR POTATOES

| Directions/Remarks | Metribuzin 75DF Lb./Acre |
|--|--------------------------|
| Apply specified dosage as a broadcast spray. Do not mechanically incorporate into soil. Use the 1/3 to 2/3 lb./acre rate for control of wild mustard (<i>Brassica</i> spp.) only. On sand soils or sensitive varieties, do not exceed 2/3 lb./acre. | 1/3 to 1 1/3 |
| Postemergence (except early maturing smooth skinned, red skinned, and other specified varieties): Apply specified dosage as a broadcast spray over the tops of potato plants*. Use rates of 1/3 to 2/3 lb./acre for control of redroot pigweed and common lambsquarters only. Apply the 2/3 lb./acre rate for control of other weeds listed on this label. | 1/3 to 2/3 |
| Split Applications: This product may be applied once preemergence and once postemergence as directed above.* Do not exceed 1 1/3 lb. total per acre per season. | |
| Idaho, Oregon, and Washington Only: Two postemergence applications can be made as broadcast sprays over the tops of potato plants if Metribuzin 75DF is applied preemergence. Use 1/3 to 2/3 lb./acre for control of redroot pigweed and lambsquarters only. On coarse (sandy) soils with low organic matter, do not exceed 1/2 lb./acre per application. On medium and heavy soils only, use 2/3 lb./acre per application for control of other weeds listed on this label and for suppression of hairy nightshade. Make the first application early in the season while weeds are still small. Allow at least 14 days before the second application. Do not apply after June 30 if treated land is to be planted to crops other than potatoes. | |
| *Refer to SPECIAL PRECAUTIONS-(Potatoes) below | |

TANK MIXES

Metribuzin 75DF may be tank mixed with the following herbicides: Dual/Dual II, Eptam, Prowl 3.3 EC, and Matrix. In addition, three-way tank mix combinations may be used for Metribuzin 75DF plus Dual/Dual II, Eptam, or Prowl 3.3 EC plus Matrix when applied preemergence. Refer to each product's label for precautionary statements, restrictions, application information, and weeds controlled.

Dual/Dual II: Metribuzin 75DF may be applied in a tank mix combination with Dual/Dual II as a preemergence broadcast application. Apply Metribuzin 75DF at 1/2 to 1 1/3 lbs. and Dual/Dual II at 1.5 to 3 pints per acre according to the respective labels for use of each product alone on potatoes.

Eptam: Metribuzin 75DF may be tank mixed with Eptam at rates and uses permitted on each product's label.

Prowl 3.3 EC: Metribuzin 75DF may be applied in tank mix combination with Prowl as a preemergence or early postemergence broadcast application. As a preemergence mix, apply Metribuzin 75DF at 2/3 to 1 1/3 lbs. and Prowl at 1.2 to 3.6 pints per acre. As an early postemergence spray, apply Metribuzin 75DF at 1/3 to 2/3 lb. and Prowl at 1.2 to 3.6 pints per acre before the crop is in the 6-inch growth stage.

Matrix (except the following counties in Colorado: Almosa, Conejos, Costilla, Rio Grande, and Saguache): Metribuzin 75DF may be applied in tank mix combination with Matrix as a preemergence and/or early postemergence application for improved control on weeds such as Russian thistle, kochia, and common lambsquarters. As a preemergence mix, apply Metribuzin 75DF at 1/3 to 3/4 lb. and Matrix at 1 to 1 1/2 oz. product per acre. As an early postemergence spray, apply Metribuzin 75DF at 1/3 to 2/3 lb. and Matrix at 1 to 1 1/2 oz. product per acre. Use a nonionic surfactant at a rate of 0.125% v/v (1 pt./100 gallons of water). Apply before the crop exceeds 14 inches in height.

Postemergence applications of Matrix treatments should be made prior to June 30.

SPECIAL PRECAUTIONS (Potatoes): Do not use Metribuzin 75DF on potatoes in Kern County, California.

Do not apply more than a total of 1 1/3 lbs. Metribuzin 75DF per acre in a single crop season regardless of the method of application.

Do not make postemergence applications prior to rainfall or irrigation on recently cultivated potatoes, nor within 3 days after periods of cool, wet cloudy weather or injury may occur.

Postemergence applications may cause some chlorosis or minor necrosis. These symptoms may be more severe if seed-piece decay is occurring or if growing conditions favor crop stress.

Postemergence applications may be made only on russet or white skinned varieties that are not early maturing.

Potato varieties may vary in their response to herbicide applications. When using Metribuzin 75DF for the first time on a particular variety, always determine crop tolerance before using on a field scale.

Do not apply Metribuzin 75DF within 60 days of harvest.

Do not use air blast sprayers.

Do not apply to sweet potatoes or yams.

Do not plant sensitive crops such as onions, lettuce, cole crops, and cucurbits during the next growing season following Metribuzin 75DF application.

Do not rotate any crop not listed on this label for 18 months following application of Metribuzin 75DF.



Certain cereal varieties are sensitive to Metribuzin 75DF (see cereal section of this label for sensitive varieties) and should not be planted during the next growing season unless the following cultural practices occur:

1. Potato vines left in rows as a result of harvest must be uniformly distributed over the soil surface prior to plowing, and
2. Plow with a moldboard plow to a depth sufficient to mix the upper 8 inches of soil.

ALFALFA AND SAINFOIN

Metribuzin 75DF herbicide is labeled for use in alfalfa and sainfoin in the following areas:

1. Alfalfa and sainfoin (including mixed stands with grasses) (all areas except California).
2. Alfalfa and sainfoin (including mixed stands with grasses) (California only).
3. Alfalfa-Tank mix combination with Gramoxone (Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming, and the following California counties: Del Norte, Lassen, Modoc, Nevada, Plumas, Shasta, Sierra, and Siskiyou).
4. Alfalfa-Post Dormant Application of Metribuzin 75DF Impregnated on Dry Fertilizer Only (Connecticut, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New Mexico, New York, Ohio, Oklahoma, Pennsylvania, South Dakota, Tennessee, Texas, and Wisconsin).
5. Alfalfa-Non-Dormant, Non-Winter Hardy varieties (Arizona only).

Metribuzin 75DF is recommended for use in aerial or ground spray equipment as a broadcast surface application to established crops of alfalfa and sainfoin for the control of certain grass and broadleaf weeds.

APPLICATION: Refer to GENERAL INFORMATION in the front of this label for detailed information on the application of Metribuzin 75DF. For information on applying Metribuzin 75DF in fluid or on dry fertilizer refer to the APPLICATION OF METRIBUZIN 75DF IN FLUID FERTILIZERS or COMMERCIAL IMPREGNATION and APPLICATION OF METRIBUZIN 75DF ON DRY BULK FERTILIZER under the GENERAL INFORMATION section of this label.

SPECIAL PRECAUTIONS: Use Metribuzin 75DF only on established alfalfa and sainfoin. Do not apply Metribuzin 75DF after growth begins in the spring or before growth ceases in the fall except as specified on this label.

Do not graze or harvest within 28 days after application.

For best weed control, apply Metribuzin 75DF when weeds are less than 2 inches tall or before weed foliage is 2 inches in diameter.

Reduced weed control may occur when extended dry conditions follow application of Metribuzin 75DF.

Crop injury may occur when:

1. Crop is under stress conditions such as diseases, insect infestation, poorly drained soils, drought or winter injury at time of application;
2. Crop is treated within 12 months after seeding;
3. There is excessive irrigation or rainfall immediately after application. Do not apply more than 1/2 inch of water in the first irrigation after Metribuzin 75DF is applied.

ALFALFA AND SAINFOIN (All Areas Except California)

| RECOMMENDED BROADCAST APPLICATIONS | |
|---|--------------------------|
| CROP | Metribuzin 75DF Lb./Acre |
| Alfalfa and Sainfoin (Except California) | 1/3 to 1 1/3 |
| Select the proper dosage according to weeds known to be and present in field to be treated. On loamy sand soils in Oregon and Washington, do not apply more than 2/3 lb. of Metribuzin 75DF per acre. | |

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES: Rates of 2/3 to 1 lb. of Metribuzin 75DF per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa. Higher rates will severely reduce forage grass stands.

Metribuzin 75DF should not be used on sand soils. In areas West of the Rocky Mountains, avoid using Metribuzin 75DF on soils with calcareous surface area, high levels of lime or sodium, or a pH greater than 8.2.

| WEEDS CONTROLLED (Except California) | |
|--|---|
| 1/3 to 1/2 lb. Metribuzin 75DF/Acre | |
| Chickweed, common (<i>Stellaria media</i>) | |
| 1/2 to 2/3 lb. Metribuzin 75DF/Acre | |
| Cheat (<i>Bromus secalinus</i>) | Pennycress (<i>Thlaspi arvense</i>) |
| Deadnettle, Purple (<i>Lamium purpureum</i>) | Rescuegrass (<i>Bromox catharticus</i>) |
| Downy brome (<i>Promus tectorum</i>) | Shepherdspurse (<i>Capsella bursa pastoris</i>) |
| Japanese brome (<i>Bromus japonicus</i>) | |
| 2/3 to 1 1/3 lb. Metribuzin 75DF/Acre | |
| Broadleaves | Grasses |
| Fleabane, Rough (<i>Erigeron striosus</i>) | Mustard, Jim Hill (tumble) (<i>Sisymbrium altissimum</i>) |
| Flixweed (<i>Descurainia sophia</i>) | Mustard, Tansy (<i>Descurainia pinnata</i>) |
| Henbit (<i>Lamium amplexicaule</i>) | Pepperweed (<i>Lepidium virginicum</i>) |
| Kochia (<i>Kochia scoparia</i>) | Pigweed, Redroot (<i>Amaranthus retroflexus</i>) |
| Lambsquarters, common (<i>Chenopodium album</i>) | Prickly Lettuce (<i>Lactuca serriola</i>) |
| Marestail (Horseweed) (<i>Hippuris vulgaris</i>) | White Cockle (<i>Melandrium album</i>) |
| Meadow Salsify (<i>Tragopogon pratensis</i>) | Wild Buckwheat (<i>Polygonum convolvulus</i>) |
| Mustard, Blue (<i>Chorispora tenella</i>) | Yellow Rocket (<i>Barbarea vulgaris</i>) |



**ALFALFA AND SAINFOIN (Continued)**

| WEEDS CONTROLLED (Except California) | |
|--|--|
| 2/3 to 1 1/3 lb. Metribuzin 75DF/Acre | |
| Grasses Foxtail, Green (<i>Setaria viridis</i>) Little Barley (<i>Hordeum pusillum</i>) | Smooth Brome (<i>Bromus inermis</i>) Wild Oats (<i>Avana fatua</i>) |
| 1 1/3 lb. Metribuzin 75DF/Acre | |
| Broadleaves Chickweed, Mouseear (<i>Cerastium vulgatum</i>) Dandelion (<i>Taraxacum officinale</i>) Ragweed, common (<i>Ambrosia artemisiifolia</i>) | Barnyardgrass (<i>Echinochloa crus-galli</i>) Bluegrass (<i>Poa annua</i>) Foxtail Barley (<i>Hordeum jubatum</i>) |

Weeds Partially Controlled: At the rate of 1 1/3 lb./acre, Metribuzin 75DF may be used to reduce the competition from curly dock (*Rumex crispus*).

At 2/3 to 1 1/3 lb./acre, Metribuzin 75DF may be used to reduce the competition of German Moss or knawel (*Scleanthus annus*).

ALFALFA AND SAINFOIN (California Only)-(Including Mixed Stands with Grasses)

Metribuzin 75DF is recommended for use in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin.

APPLICATION: Metribuzin 75DF is recommended for use in aerial or ground spray equipment as a broadcast surface application to dormant established crops of alfalfa and sainfoin for control of certain grass and broadleaf weeds. Do not apply Metribuzin 75DF after growth begins in the spring or before growth ceases in the fall. Do not apply to either alfalfa or sainfoin during the first growing season after seeding.

For information on applying Metribuzin 75DF in fluid fertilizer solutions to alfalfa, refer to the appropriate section of this label.

For information on commercial impregnation and application of Metribuzin 75DF on dry bulk fertilizer, refer to the appropriate section of this label.

| WEEDS CONTROLLED | |
|---|---|
| 1/2 to 2/3 lb. Metribuzin 75DF/Acre | |
| Cheatgrass (downy brome) (<i>Bromus secalinus</i>) | |
| 2/3 to 1 1/3 lb. Metribuzin 75DF/Acre | |
| Broadleaves Chickweed, common (<i>Stellaria media</i>) Flaxweed (<i>Descurainia sophia</i>) Henbit (<i>Lamium amplexicaule</i>) Kochia (<i>Kochia scoparia</i>) Meadow Salsify (<i>Tragopogon pratensis</i>) Mustard, Blue (<i>Chorispora tenella</i>) | Mustard, Tansy (<i>Descurainia pinnata</i>) Pepperweed, Virginia (<i>Lepidium virginicum</i>) Shepherdspurse (<i>Capsella bursa pastoris</i>) White Cockle (<i>Melandrium album</i>) Wild Buckwheat (<i>Polygonium convolvulus</i>) Yellow Rocket (<i>Barbarea vulgaris</i>) |
| Grasses Smooth Brome (<i>Bromus inermis</i>) Wild Oats (<i>Avana fatua</i>) | |
| 1 1/3 lb. Metribuzin 75DF/Acre | |
| Broadleaves Dandelion (<i>Taraxacum officinale</i>) | Grasses Barnyardgrass (<i>Echinochloa crus-galli</i>) Bluegrass (<i>Poa annua</i>) Foxtail Barley (<i>Hordeum jubatum</i>) |

| RECOMMENDED BROADCAST APPLICATIONS | |
|--|---------------------------------|
| CROP | Metribuzin 75DF Lb./Acre |
| Alfalfa and Sainfoin (California Only) | 1/2 to 1 1/3 |
| Select the proper dosage according to weeds known to be and present in field to be treated. Apply specified dosage in 20 to 40 gallons of water per acre with ground spray equipment or 3 to 10 gallons of water per acre with aerial spray equipment fitted with nozzles suitable for broadcast applications of herbicides. Treat only dormant established crops of alfalfa and sainfoin. Injury may occur to alfalfa if Metribuzin 75DF is applied earlier than 12 months after seeding. Do not apply after Spring growth begins or before growth ceases in the fall. Do not graze or harvest within 28 days after application. At the 1 1/3 lb./acre rate, Metribuzin 75DF may be used for suppression of curly dock. | |





FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES: Rates of 2/3 to 1 1/3 lb. of Metribuzin 75DF per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa. Higher rates will severely reduce forage grass stands.

Do not apply with aerial spray equipment when wind speed is greater than 10 mph. Do not apply when weather conditions favor spray drift and/or when sensitive cool season crops, such as cole crops, onions, peas, or strawberries, are present in adjacent fields. Applications should not be made when weather conditions favor spray drift, especially in areas where wheat is growing on coarse-textured soils in adjacent fields, or injury may occur.

ALFALFA

METRIBUZIN 75DF PLUS GRAMOXONE EXTRA TANK MIX

Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming, and the following California counties: Del Norte, Lassen, Modoc, Nevada, Plumas, Shasta, Sierra, and Siskiyou.

APPLICATION: Metribuzin 75DF plus Gramoxone Extra tank mix application is recommended for use, during the dormant season, in aerial or ground spray equipment as a broadcast surface application to established (at least 1 year old) alfalfa for the control of certain grass and broadleaf weeds. Do not apply Metribuzin 75DF/ Gramoxone Extra tank mix to regrowth (after grazing or cutting) that is more than 2 inches tall. Apply once per season. Do not apply following cuttings during growing season. Use a minimum of 10 gallons of water per acre with aerial spray equipment and a minimum of 20 gallons of water per acre with ground spray equipment. Add a non-ionic spreader at label rates to the spray solution.

| WEEDS CONTROLLED | |
|--|--|
| Metribuzin 75DF plus Gramoxone Extra (1 1/2 to 2 1/2 pts./acre) tank mix application will control established weeds. Gramoxone controls weeds by contact activity. | |
| 1/3 to 1/2 lb. of Metribuzin 75DF/Acre | |
| Common Chickweed | |
| 1/2 to 1 lb. of Metribuzin 75DF/Acre | |
| Bluegrass Cheat Downy brome Field pennycress | Henbit Japanese brome Rescuegrass Shepherdspurse |
| Use Metribuzin 75DF at 2/3 to 1 lb./Acre for control of the following weeds: | |
| Blue mustard Common lambsquarters Flixweed Green foxtail Groundsel Jim Hill mustard Kochia Little barley Marestail (Horseweed) Meadow salsify Pepperweed | Prickly lettuce Redroot pigweed Rough fleabane Ryegrass Smooth brome Sowthistle Tansy mustard White cockle Wild oats Wild buckwheat Yellow rocket |
| RECOMMENDED APPLICATIONS | |
| Dosage/Acre | Directions/Remarks |
| Metribuzin 75DF 1/3 to 1 lb. plus Gramoxone Extra 1 1/2 to 2 1/2 pt. | Apply specified dosages of Metribuzin 75DF and Gramoxone Extra in at least 10 gallons of water per acre with aerial equipment or at least 20 gallons of water per acre with ground equipment. Do not apply this tank mix to alfalfa growth if more than 2 inches tall. For best weed control, apply when broadleaf weeds and grasses are 1-6 inches tall and are actively growing. Care should be taken to avoid overlaps. Do not apply more than 2/3 lb. of Metribuzin 75DF per acre on loamy sand soils. Reduced weed control may occur when extended dry conditions follow application of Metribuzin 75DF. Crop injury may occur if alfalfa is under stress conditions such as diseases, insect infestations, drought, or winter injury or if Metribuzin 75DF is applied to alfalfa earlier than 12 months after seeding. |

FOR USE ON MIXED STANDS OF ALFALFA AND GRASSES: Rates of 2/3 to 1 lb. of Metribuzin 75DF per acre will provide partial reduction of forage grass stands. These rates may be used to reduce forage grass stands to prevent crowding out of alfalfa.

Do not graze or harvest within 42 days after application.

In areas west of the Rockies, avoid the use of Metribuzin 75DF on soils with calcareous surface, soils with high levels of lime or sodium, and with a pH greater than 8.2.

Do not apply when weather conditions favor spray drift. Aerial application should not be made when wind speed is greater than 10 mph.

Do not use on sand soil.

Refer to the Gramoxone Extra label for additional directions, weed species controlled, and precautions.





POST DORMANT APPLICATION OF METRIBUZIN 75DF IMPREGNATED ON DRY FERTILIZER ONLY

Metribuzin 75DF may be applied after dormancy has broken, but prior to three inches of new alfalfa shoot growth, only when impregnated on dry fertilizer in Connecticut, Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Nebraska, New Mexico, New York, Ohio, Oklahoma, Pennsylvania, South Dakota, Tennessee, Texas, and Wisconsin. Apply at rates of 1 to 1 1/3 lb. per acre as directed on this label for application during dormancy. Apply only when alfalfa foliage is dry or crop injury may occur. When using this application method do not harvest or graze treated alfalfa for 60 days after application.

ALFALFA (Arizona Only)

Non-dormant, Non-Winter Hardy Varieties

Metribuzin 75DF is recommended as a broadcast surface application to established crops of non-dormant alfalfa varieties for preemergence and postemergence control of certain winter annual weeds following either a fall or winter sheep grazing/green-chop harvest.

| WEEDS CONTROLLED | |
|---|---|
| Field pepperweed Lambsquarters Little mallow (cheeseweed) Littleseed canarygrass London rocket (mustard) Prickly lettuce | Mouse barley Nettleleaf Shepherdspurse Silversheath knotweed Spiny sowthistle |

| RECOMMENDED APPLICATIONS (Arizona only) | |
|---|--------------------------|
| CROP | METRIBUZIN 75DF Lb./ACRE |
| Alfalfa Non-dormant Non-winter Hardy varieties | 1/2 to 2/3 |

Apply specified dosage by aerial or ground spray equipment in 7 to 40 gallons of water per acre. Treat established alfalfa stubble after fall or winter sheep grazing or green-chop harvest and prior to the time regrowth is 2" tall. Alfalfa foliage present at time of application can exhibit yellowing. Injury may occur to alfalfa in areas of high salt concentration where the crop is stunted and/or has a poorly developed root system, or if alfalfa is under stressed growing conditions such as diseases, insect infestations, or drought. For most effective postemergence weed control, treatment should be made before weeds are 2" tall or before leaf rosettes are 2" wide. For maximum control, rainfall (1/4" or more) or irrigation is necessary within 30 days of treatment, however, do not flood irrigate within 2 days after treatment. Use 1/2 lb. Metribuzin 75DF on sand soil when only mustard, goosefoot, lambsquarters, or canary grass are the weeds to be controlled. Do not apply earlier than 6 months after seeding. Do not graze or harvest within 28 days after application.

SPECIAL PRECAUTIONS: Maintain continuous mechanical agitation in the spray tank to insure a uniform spray mixture.

Do not apply with aerial spray equipment when wind speed is greater than 10 mph. Do not apply when weather conditions favor spray drift and/or when sensitive cool season crops such as cole crops, onions, peas, or strawberries are present in adjacent fields. Applications should not be made when weather conditions favor drift especially in areas where wheat is growing on coarse-textured soils in adjacent fields or injury may occur.

ASPARAGUS (Established)

Metribuzin 75DF is recommended for use in ground spray equipment or sprinkler irrigation (center pivot, lateral move, or solid set) systems as a single preemergence broadcast application or as a split application consisting of a preemergence broadcast application followed by a post harvest broadcast application.

Aerial application is prohibited.

Refer to the GENERAL INFORMATION section of this label for directions.

| WEEDS CONTROLLED | |
|---|---|
| Metribuzin 75DF applied to established asparagus according to directions will effectively control: | |
| Broadleaves Chickweed, Common (<i>Stellaria media</i>) Jimsonweed (<i>Datura stramonium</i>) Lambsquarters (<i>Chenopodium album</i>) Pigweed, Redroot (<i>Amaranthus retroflexus</i>) | Ragweed, common (<i>Ambrosia artemisiifolia</i>) Smartweed, Pennsylvania (<i>Polygonum pensylvanicum</i>) Sorrel, Red (<i>Rumex acetosella</i>) Velvetleaf (<i>Abutilon theophrasti</i>) |
| Grasses Crabgrass (<i>Digitaria</i> spp.) Foxtails (<i>Setaria</i> spp.) | Sandbur, Field (<i>Cenchrus pauciflorus</i>) |





| RECOMMENDED BROADCAST APPLICATIONS (Asparagus) | |
|---|--|
| Directions/Remarks | Metribuzin 75DF Lb./Acre |
| Preemergence Application Only: Make a single surface application in early spring before asparagus spears or ferns emerge. If the field is to be disked, apply Metribuzin 75DF after disking but before the crop emerges. Use the lower rate for control of the broadleaf weeds listed above. Use the higher rate in fields with a history of severe infestations of grasses and for maximum residual control. Do not apply within 14 days of harvest. | 1 1/3 to 2 2/3 |
| Split Application- Preemergence and Post Harvest: Preemergence Application: Apply before asparagus spears or ferns emerge. If the field is to be disked, apply after disking but prior to crop emergence. Do not apply within 14 days of harvest. Post Harvest Application: Apply after last harvest of the season but prior to emergence. The lower combination rates may be used for control of common ragweed, lambsquarters, redroot pigweed, and red sorrel. Use the higher combination rates for other weeds listed or in fields with severe post harvest grass infestations or for maximum post harvest control of emerged weeds. | 2/3 to 1 1/3 preemergence + 1 1/3 to 2 |
| IMPORTANT: The total amount of Metribuzin 75DF applied in one crop season may not exceed 2 2/3 lb. per acre. | |

SPECIAL PRECAUTIONS (Asparagus): Do not use on newly seeded asparagus or on young plants during the first growing season after setting crowns.

DO NOT APPLY POST HARVEST APPLICATIONS UNTIL AFTER THE LAST HARVEST OF SPEARS.

CARROTS

Special Conditions of Sale Provision for Use on Carrots: The following directions for use were developed under the direction of IR-4 (government minor crops use program). As such the testing was done independently from the testing program of Makhteshim Agan of North America, Inc. Buyer is advised that Makhteshim Agan makes no assurances regarding satisfaction with the product and all risks of crop injury or product performance are assumed by the Buyer.

Apply Metribuzin 75DF herbicide with ground equipment as specified below under RECOMMENDED APPLICATIONS. For effective control of broadleaf

weeds with postemergence applications, apply Metribuzin 75DF before weeds are 1 inch in height or diameter. Thorough spray coverage is essential for adequate weed control.

Do not use air blast or other high pressure spray equipment to make postemergence applications of Metribuzin 75DF. Refer to the appropriate section of this label for additional information regarding spray equipment, dilution rates, mixing, sprayer cleanup, restrictions, container disposal, and cautions.

Refer to MIXING under the GENERAL INFORMATION section in the front of this label.

For specific application information see GENERAL INFORMATION and APPLICATION sections at the front of this label.

| WEEDS CONTROLLED | |
|--|--|
| Metribuzin 75DF applied to carrots according to directions will effectively control: | |
| Carpetweed (<i>Mullugo verticillata</i>) | Pigweed, Redroot (<i>Amaranthus retroflexus</i>) |
| Galinsoga (<i>Galinsoga parviflora</i>) | Pigweed, Smooth (<i>Amaranthus hybridus</i>) |
| Horseweed (<i>Conyza Canadensis</i>) | Pineappleweed (<i>Matricaria matricarioides</i>) |
| Lambsquarters, Common (<i>Chenopodium album</i>) | Prickly Lettuce (<i>Lactuca serriola</i>) |
| Mustard, Wild (<i>Sinapis arvensis</i>) | Shepherdspurse (<i>Capsella bursa-pastoris</i>) |

| RECOMMENDED APPLICATIONS (Carrots) | |
|---|--------------------------|
| Directions/Remarks | Metribuzin 75DF lb./Acre |
| Apply specified dosage per acre as a broadcast spray over the tops of carrot plants. Application should be made after carrots have formed 5 to 6 true leaves but before weeds are 1 inch in height or diameter. If needed, a second application may be made after an interval of at least 3 weeks. Applications may be made up to 60 days of harvest. | 1/3 |
| IMPORTANT: The total amount of Metribuzin 75DF applied in one crop season must not exceed 2/3 lb. per acre. | |

SPECIAL PRECAUTIONS:

Do not apply to carrots grown for seed.

Do not apply within 3 days after periods of cool, wet, or cloudy weather or crop injury will occur.

Do not apply Metribuzin 75DF within 3 days of any other chemical unless specified on this label.

Do not apply on very hot days or excessive crop injury will result.

Do not apply until carrots have at least 5 to 6 true leaves. Earlier applications will result in excessive crop damage.

Crop injury or delayed maturity may result from applications of Metribuzin 75DF if carrots are growing under stress conditions such as periods of drought or cool, wet, and cloudy weather preceding application.

Following an application of Metribuzin 75DF, chlorosis (yellowing) and burning of the leaf tissue may occur.

For newly introduced varieties of carrots with unknown tolerance to Metribuzin 75DF, treat only a small area to determine if Metribuzin 75DF can be used without injury to the crop.





FIELD CORN

POSTEMERGENCE APPLICATION

Metribuzin 75DF is recommended for control of selected broadleaf weeds when applied as a tank mix combination with certain broadleaf herbicides presently registered and recommended for postemergence use in field corn. Herbicides which may be tank mixed with Metribuzin 75DF include:

| | | |
|---------------------|-----------------------------|--------------|
| 2,4-D | Buctril + atrazine (Premix) | Resource |
| Atrazine | Clarity | Scorpion III |
| Banvel | Laddok S-12 | Tough |
| Basagran | Marksman | |
| Buctril/Buctril Gel | Pursuit* | |

* Use only on Pursuit resistant/tolerant corn hybrids (IMI-Corn)

APPLICATION: Metribuzin 75DF may be applied to field corn after crop emergence until just prior to tasseling. Broadcast applications may be made with ground or aerial equipment. For optimum weed control, apply treatments when weeds are small and actively growing, but before reaching the maximum heights listed in the WEEDS CONTROLLED table.

POSTEMERGENCE BROADCAST APPLICATION

Ground Application: Adjust nozzle height above crop and weed canopy to ensure uniform spray coverage. Gallonage should be increased with increasing weed size and population density.

For tank mixes of Metribuzin 75DF plus atrazine, Basagran, Laddok S-12, Buctril, Buctril + atrazine, Pursuit, Resource, Tough, or 2,4-D amine formulations, use flat fan nozzles spaced a maximum of 20 inches apart. Best results are achieved using a minimum spray volume of 10 gallons per acre and spray pressure from 20 to 40 psi.

For Metribuzin 75DF tank mixes with Banvel, Clarity, Marksman, or 2,4-D low volatile ester formulations, use drift-reducing nozzles which are specifically designed to produce coarse sprays and reduce the amount of driftable fines. Additional measures which will help avoid potential drift to sensitive crops and plants include using a minimum spray volume of 20 gallons per acre and keeping spray pressures at or below 20 psi unless otherwise specified by the nozzle manufacturer.

For further precautions and additional instructions and recommendations, consult the tank mix partner's label.

Aerial Application: Apply in a minimum spray volume of 3 gallons per acre. For optimum spray coverage and distribution, use a minimum of 5 gallons per acre and a maximum pressure of 40 psi. Use a boom and nozzle configuration which will provide a uniform deposition pattern and coverage with low drift potential. Avoid overlaps to prevent potential crop injury. Do not apply near sensitive crops or sensitive plants growing near the treated area. Do not apply when wind speed is greater than 10 mph or when winds are moving toward sensitive crops or plants. To avoid drift hazards, applicator must follow the most restrictive labeling of the products used in a tank mix. Refer to the appropriate tank mix partner's label for further precautions and recommendations.

POSTDIRECTED APPLICATION

Metribuzin 75DF in tank mix combinations with Banvel, 2,4-D, Buctril, or Scorpion III may be applied postdirected to field corn. Use drop nozzles and appropriate spacing to direct spray below the corn whorl and upper leaves. The top of the target weed canopy must be sufficiently below the whorl and upper leaves of the crop to permit this application and provide adequate spray coverage. The height differential required between the crop and weed canopy will depend on the specific equipment used. Apply before tassel emergence. For further precautions and additional recommendations, refer to the appropriate tank mix partner's label.

ADJUVANTS

The adjuvant types listed below may be utilized with certain Metribuzin 75DF tank mix combinations. Consult the tank mix recommendations section for the

appropriate adjuvant and rate. Use of non-recommended adjuvants or rates may result in severe leaf burn, crop stunting, and/or stand reduction. Use only adjuvants which are exempt from tolerance requirements under 40 CFR 180.1001.

UAN (urea ammonium nitrate) is commonly referred to as 28, 30, or 32% N.

Ammonium sulfate (spray grade) may be used as alternative to UAN with certain tank mix combinations.

Non-ionic surfactants should contain at least 80% active ingredient.

DO NOT USE crop oil concentrate (COC) or any adjuvant containing vegetable or petroleum oils with any Metribuzin 75DF tank mixtures as severe leaf burn, crop stunting, and/or stand reduction may occur.

RAINFASTNESS

Metribuzin 75DF will not reduce rainfastness of the recommended tank mix partners. Refer to the individual product labels for rainfastness recommendations.

SPRAYER CLEANUP

Refer to each tank mix partner's label and the Sprayer Cleanup section of the Metribuzin 75DF label for specific instructions on cleaning spray equipment. Special attention should be given to the required cleanup procedures for 2,4-D, Banvel, Clarity, and Marksman.

SPECIAL PRECAUTIONS

1. Do not use on corn grown for seed, sweet corn, popcorn, or white corn.
2. Do not apply more than 0.25 pounds a.i. metribuzin (5 1/3 ounces Metribuzin 75DF) per acre per use season.
3. Do not apply when field corn is under stress (see Stress statement below).
4. Do not use aerial applications if sensitive crops or plants are growing in the vicinity of the area to be treated.
5. Do not allow spray drift onto sensitive crops or plants.
6. Do not use on sand, loamy sand, or sandy loam soils that have less than 0.5% organic matter.
7. Do not use on sand or loamy sand soils in Washington, Oregon, or Idaho or crop injury may occur.
8. Observe all precautions and limitations on labeling of all products used in the tank mixtures.

Stress is any condition or combination of conditions which impairs normal crop growth. Weather, disease, insect damage, fertility, or other factors may cause stress. Applications made before or after the corn is under stress from these factors or from periods of prolonged cool, wet, and cloudy weather or widely fluctuating day and nighttime temperatures, may result in temporary leaf burn, yellowing, and/or stunting of the crop. Recovery from damage is generally rapid with no lasting effects on new growth. Under extreme stress, stand reductions may occur.

Feeding Restrictions: Field corn treated with Metribuzin 75DF may be grazed or harvested for silage or grain 60 days after treatment. Follow the most restrictive preharvest interval on the labels of the products used in the tank mixtures.



TANK MIX COMBINATIONS

The Metribuzin 75DF tank mixtures listed below can be utilized for control of certain annual broadleaf weeds.

| METRIBUZIN 75DF POSTEMERGENCE BROADCAST RECOMMENDATIONS | | |
|--|--|---|
| PRODUCT | RATE | DIRECTIONS AND REMARKS* |
| Metribuzin 75DF + 2,4-D Amine or 2,4-D LVE | 2 oz./A + 1/2 to 1 pt./A' or 1/3 to 1/2 pt./A' | Apply as a broadcast spray during the interval from corn emergence until corn is 8 inches tall. Apply only to varieties known to be tolerant to 2,4-D. DO NOT USE ADJUVANTS. 2,4-D may cause injury to nearby sensitive crops. 2,4-D applications may result in brittle corn stalks and winds or cultivation may cause stalk breakage. To reduce damage, delay cultivation 8 to 10 days after application. |
| Metribuzin 75DF + Atrazine | 2 oz./A + 1/2 to 1 1/2 lb. ai/A | Apply as a broadcast spray during the interval from corn emergence until corn is 12 inches tall. A non-ionic surfactant (1 qt./100 gal of spray solution) may be added to improve weed control. Atrazine is a restricted use herbicide. Follow all state and federal label recommendations and restrictions pertaining to atrazine applications. |
| Metribuzin 75DF + Banvel or Clarity | 2 oz./A + 1/2 to 1 pt./A or 1/2 to 1 pt./A | Apply as a broadcast spray during the interval from corn emergence through the five-leaf stage or when corn is 8 inches tall, whichever occurs first. For Banvel applications to corn greater than 8 inches in height, consult the Banvel label for use rates and restrictions. If growing conditions are dry and plants are stressed, addition of a non-ionic surfactant (1 qt./100 gal of spray solution) may improve weed control. For corn grown on coarse-textured soils, apply Banvel or Clarity at 0.5 pt./A, regardless of application method. Application may cause injury to nearby sensitive crops or plants. Application may result in temporary leaning of corn plants. Delay cultivation until plants return to normal growth patterns to avoid stalk breakage. |
| Metribuzin 75DF + Basagran | 2 oz./A + 1 pt./A | Apply as a broadcast spray after corn emergence but before corn exceeds 30 inches in height and the crop canopy closes the row. Adjuvants such as UAN (0.5 to 1 gal./A), ammonium sulfate (17 lbs./100 gal of spray solution), or non-ionic surfactant (1 qt./100 gal of spray solution) may improve weed control. |
| Metribuzin 75DF + Buctril or Buctril Gel | 1.6 to 2 oz./A + 1 pt./A or 1/2 pt./A | Apply as a broadcast spray when corn is in the fourth true leaf stage or later but before the crop canopy closes the row. DO NOT USE ADJUVANTS. Occasional temporary corn leaf burn may occur and is similar to that observed from liquid fertilizers. Recovery is generally rapid with no lasting effect. To reduce potential for crop damage, application should be made to dry corn foliage when weather conditions are not extreme. |
| Metribuzin 75DF + Buctril + atrazine (Premix) | 1.6 to 2 oz./A + 1 1/2 to 2 pt./A | Apply as a broadcast spray during the interval from corn emergence until corn is 12 inches tall. DO NOT USE ADJUVANTS. Occasional temporary corn leaf burn may occur and is similar to that observed from liquid fertilizers. Recovery is generally rapid with no lasting effect. To reduce potential for crop damage, application should be made to dry corn foliage when weather conditions are not extreme. |
| Metribuzin 75DF + Laddok S-12 | 2 oz./A + 1.33 to 1.66 pt./A | Apply as a broadcast spray after corn emergence until the corn is 12 inches tall. Adjuvants such as UAN (0.5 to 1 gal./A) may increase weed control. Laddok S-12 contains atrazine and is a restricted use product. Follow all state and federal label recommendations and restrictions pertaining to atrazine. |
| Metribuzin 75DF + Marksman | 2 oz./A + 1 1/2 to 2 pt./A | Apply as a broadcast spray during the interval from corn emergence through the five-leaf stage or when corn is 8 inches tall, whichever occurs first. DO NOT USE ADJUVANTS. Application may cause injury to nearby sensitive crops or plants. Application may result in temporary leaning of corn plants. Delay cultivation until plants return to normal growth patterns to avoid stalk breakage. Marksman contains atrazine and is a restricted use product. Follow all state and federal label recommendations and restrictions pertaining to atrazine. |
| Metribuzin 75DF + Pursuit | 2 oz./A + 2 to 4 oz./A | Use only on designated IMI-Corn hybrids (hybrids which are resistant/tolerant to Pursuit). Apply the 4-ounce rate of Pursuit if grasses are present or broadleaf weeds are near the maximum heights shown. Apply in combination with a non-ionic surfactant (1 qt./100 gal of spray solution) and UAN (1 to 2 qt./A). |
| Metribuzin 75DF + Resource | 3 fl. oz./A + 4 to 6 fl. oz./A | Apply as a broadcast spray to field corn from 2-leaf through 10-leaf (visible leaf collars) stage. Adjuvants such as nonionic surfactant (0.25% v/v), UAN (2% v/v) or ammonium sulfate (2.5 lbs./A) may increase weed control. |

| PRODUCT | RATE | DIRECTIONS AND REMARKS* |
|---|------------------------------|--|
| Metribuzin 75DF + Tough | 2 oz./A + 1 to 2 pt./A | Apply as a broadcast spray after corn emergence but before corn height exceeds 30 inches and the crop canopy closes the row. A non-ionic surfactant (1 qt./100 gal of spray solution) may be added to improve weed control. Use the higher rates of Tough as weeds approach the maximum height listed or are found in high density. Tough may improve control on triazine/ALS - resistant weeds. |
| * Consult the appropriate tank mix partner's label for additional recommendations or restrictions. The most restrictive labeling applies to tank mixes with Metribuzin 75DF. ¹ Application rate is based on, but not restricted to 4 pounds active ingredient per gallon of 2,4-D. | | |

| WEEDS CONTROLLED-POSTEMERGENCE BROADCAST APPLICATION | | | | | | | | | | |
|---|-------------------|--------------------|----------------|--------------------------------------|----------------|----------|----------------|-----------------|----------|-------|
| These tank mixtures with Metribuzin 75DF will control the following annual weeds up to the maximum weed heights listed: | | | | | | | | | | |
| Common weed Name | METRIBUZIN 75DF + | | | | | | | | | |
| | Atrazine | Banvel/ Clarity | Basagran | Buctril/ Buctril + atrazine | 2,4-D | Marksman | Pursuit | Laddok S-112 | Resource | Tough |
| MAXIMUM WEED HEIGHT IN INCHES* | | | | | | | | | | |
| Amaranth, Palmer | 4 ^a | 4 | 2 ^a | 4 ^a | 4 | 4 | 8 ^b | 6 | 4 | 4 |
| Buckwheat, wild | 3 | 3 | 3 | 3 | 2 | 3 | 2 | 3 | 4 | |
| Buffalobur | 4 | 4 | | 4 | | 4 | 1 | | | |
| Burcucumber | | 4 | | 4 | 2 | 4 | | | | 4 |
| Carpetweed | 2 | 2 | 2 | 2 | 2 | 2 | | 2 | 3 | 4 |
| Cocklebur, common | 8 | 8 | 8 | 8 | 8 | 8 | 8 ^b | 8 | 3 | 6 |
| Eclipta | 3 | 3 | 3 | 3 | 3 | 3 | | 3 | | 3 |
| Henbit | 3 | 3 | 2 | 2 | 2 | 4 | 3 | 3 | | 4 |
| Horseweed/marestail | 3 | 4 | 1 | 1 | 3 | 6 | | 2 | 3 | |
| Jimsonweed | 5 | 5 | 6 | 5 | 5 | 5 | 5 | 6 | 3 | 5 |
| Knotweed | 6 | 6 | 6 | 4 | 2 | 6 | 4 | 6 | | 4 |
| Kochia | 2 ^a | 2 | 1 ^a | 2 ^a | 2 ^a | 2 | 2 | 2 ^a | | 4 |
| Ladysthumb | 6 | 6 | 6 | 6 | 4 | 6 | 4 | 6 | 4 | 6 |
| Lambsquarters, common | 6 ^a | 6 | 1 | 6 | 6 | 6 | 4 | 5 | 4 | 4 |
| Lettuce, prickly | 4 | 4 | | 3 | 4 | 5 | | 3 | | |
| Mallow, Venice | 2 | 2 | 2 | 2 | 2 | 2 | 2 | 4 | | |
| Morningglory, entire leaf | 3 | 3 | 1 | 3 | 3 | 3 | 2 | 2 | | |
| Morningglory, ivyleaf | 3 | 3 | 1 | 3 | 3 | 3 | 2 | 2 | | |
| Morningglory, pitted | 3 | 3 | 1 | 3 | 3 | 3 | 2 | 2 | | |
| Morningglory, tall | 3 | 3 | 1 | 3 | 3 | 3 | 2 | 2 | | |
| Mustard, tansy | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | | |
| Mustard, wild | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | | 4 |
| Nightshade, black | 6 | 6 | | 6 | 1 | 6 | 3 | 1 | | 4 |
| Nightshade, eastern black | 6 | 6 | | 6 | 1 | 6 | 3 | 1 | | 4 |
| Pigweed, redroot | 6 ^a | 6 | 2 ^a | 6 ^a | 6 | 6 | 8 ^b | 6 ^a | 4 | 6 |
| Pigweed, smooth | 6 ^a | 6 | 2 ^a | 6 ^a | 6 | 6 | 8 ^b | 6 ^a | 4 | 6 |
| Poorjoe | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | | |
| Purslane, common | 1 | 3 | | | | 4 | 1 | | | 3 |

(continued)



| WEEDS CONTROLLED-POSTEMERGENCE BROADCAST APPLICATION (continued) | | | | | | | | | | |
|---|-------------------|--------------------|----------------|--------------------------------------|-------|----------|----------------|-----------------|----------|-------|
| These tank mixtures with Metribuzin 75DF will control the following annual weeds up to the maximum weed heights listed: | | | | | | | | | | |
| Common weed Name | METRIBUZIN 75DF + | | | | | | | | | |
| | Atrazine | Banvel/ Clarity | Basagran | Buctril/ Buctril + atrazine | 2,4-D | Marksman | Pursuit | Laddok S-112 | Resource | Tough |
| MAXIMUM WEED HEIGHT IN INCHES* | | | | | | | | | | |
| Pusley, Florida | 3 | 3 | 3 | 3 | 3 | 3 | | 3 | 3 | |
| Ragweed, common | 5 | 5 | 3 | 5 | 5 | 6 | 3 | 4 | 3 | |
| Ragweed, giant | 4 | 5 | 2 | 4 | 3 | 6 | 4 | 4 | | |
| Sicklepod | 3 | 3 | 3 | 3 | 3 | 3 | 3 | 3 | | 2 |
| Sida, prickly | 1 | 1 | 3 | 1 | 1 | 2 | 1 | 2 | 2 | 1 |
| Smartweed, Pennsylvania | 6 | 6 | 6 | 6 | 4 | 6 | 4 | 6 | 4 | |
| Sunflower, common | 6 | 6 | 6 | 6 | 6 | 6 | 5 | 6 | | 5 |
| Thistle, Russian | 1 | 3 | | 3 | 1 | 3 | 1 | 1 | | 3 |
| Velvetleaf | 6 ^a | 6 | 6 | 6 | 4 | 6 | 5 | 6 | 6 | |
| Waterhemp, spp. | 5 ^a | 5 | 2 ^a | 5 ^a | 5 | 5 | 4 ^b | 2 ^a | 4 | 5 |

* When weeds are approaching the maximum height listed or found in high densities, use the higher rate of Metribuzin 75DF and the selected tank mix partners.

^a These treatments will not control triazine-resistant biotypes.

^b These treatments will not control ALS-resistant biotypes.

| METRIBUZIN 75DF POSTDIRECTED RECOMMENDATIONS | | |
|--|---|--|
| PRODUCT | RATE | DIRECTIONS AND REMARKS* |
| Metribuzin 75DF + 2,4-D Amine or 2,4-D LVE | 2 to 3 oz./A + 3/4 to 1 1/2 pt./A ¹ or 1/2 to 3/4 pt./A ¹ | For corn greater than 8-inches tall, apply as a directed spray with drop nozzles before tassel emergence. Apply only to varieties known to be tolerant to 2,4-D. DO NOT USE ADJUVANTS. 2,4-D may cause injury to nearby sensitive crops. 2,4-D applications may result in brittle corn stalks and winds or cultivation may cause stalk breakage. To reduce damage, delay cultivation 8 to 10 days after application. |
| Metribuzin 75DF + Banvel | 2 to 3 oz./A + 1/2 pt./A | For corn 8- to 36-inches tall, apply as a directed spray with drop nozzles. Application may be made up to 15 days prior to corn tasseling. If growing conditions are dry and plants are stressed, addition of a non-ionic surfactant (1 qt./100 gal of spray solution) may improve weed control. For corn grown on coarse-textured soils, apply Banvel at 0.5 pt./acre, regardless of application method. Application may cause injury to nearby sensitive crops or plants. Application may result in temporary leaning of corn plants. Delay cultivation until plants return to normal growth patterns to avoid stalk breakage. |
| Metribuzin 75DF + Buctril or Buctril Gel | 2 to 3 oz./A + 1 to 1 1/2 pt./A or 1/2 to 3/4 pt./A | Apply as a directed spray with drop nozzles before tassel emergence. DO NOT USE ADJUVANTS. Occasional temporary corn leaf burn may occur and is similar to that observed from liquid fertilizers. Recovery is generally rapid with no lasting effect. To reduce potential for crop damage, application should be made to dry corn foliage when weather conditions are not extreme. |
| Metribuzin 75DF + Scorpion III | 3 to 4 1/2 fl. oz./A + 4 oz./A | For corn 8 - to 24 - inches tall, apply as a directed spray with drop nozzles. Include nonionic surfactant (1 qt./100 gal) plus UAN (2.5 gal./100 gal) for optimum weed control. |

* Consult the appropriate tank mix partner's label for additional recommendations or restrictions. The most restrictive labeling applies to tank mixes with Metribuzin 75DF.

¹ Application rate is based on but not restricted to 4 pounds active ingredient per gallon of 2,4-D.





| WEEDS CONTROLLED-POSTDIRECTED APPLICATION | | | | |
|---|--------------------------------|--------|---------|--------------|
| These tank mixtures with Metribuzin 75DF will control the following annual weeds up to the maximum weed heights listed: | | | | |
| COMMON WEED NAME | METRIBUZIN 75DF + | | | |
| | 2,4-D | Banvel | Buctril | Scorpion III |
| | MAXIMUM WEED HEIGHT IN INCHES* | | | |
| Amaranth, Palmer | 12 | 12 | 6 | 8 |
| Cocklebur, common | 12 | 12 | 12 | 15 |
| Jimsonweed | 12 | 10 | 10 | 8 |
| Ladythumb | 6 | 8 | 6 | 6 |
| Lambsquarters, common | 12 | 12 | 10 | 12 |
| Morningglory, entire leaf | 18 | 18 | 6 | 12 |
| Morningglory, ivyleaf | 18 | 18 | 6 | 12 |
| Morningglory, pitted | 18 | 18 | 6 | 12 |
| Morningglory, tall | 18 | 18 | 6 | 12 |
| Nightshade, black | 10 | 8 | 8 | 6 |
| Nightshade, eastern black | 10 | 8 | 8 | 6 |
| Pigweed, redroot | 12 | 12 | 6 | 8 |
| Pigweed, smooth | 12 | 12 | 6 | 8 |
| Ragweed, common | 8 | 8 | 8 | 10 |
| Ragweed, giant | 12 | 12 | 8 | 15 |
| Smartweed, Pennsylvania | 6 | 8 | 6 | 6 |
| Sunflower, common | 12 | 12 | 12 | 12 |
| Velvetleaf | 10 | 8 | 8 | 8 |
| Waterhemp tall, spp. | 12 | 12 | 6 | 8 |

* When weeds are approaching the maximum height listed or found in high densities, use the higher rate of Metribuzin 75DF and the selected tank mix partners.

PERENNIAL WEED SUPPRESSION

The following Metribuzin 75DF tank mixtures will provide top growth burndown and in season suppression of the following perennial weeds; however, regrowth may occur. For the best performance on these weeds, use the maximum rates of Metribuzin 75DF, Banvel, Buctril, Buctril + atrazine, Clarity, Marksman, 2,4-D LVE, or Pursuit recommended for these tank mixtures.

Metribuzin 75DF + Banvel or Clarity

Bindweed, field; Dandelion, common; Dock, curly; Smartweed, swamp; Thistle, Canada

Metribuzin 75DF + Buctril or Buctril + atrazine

Thistle, Canada

Metribuzin 75DF + 2,4-D LVE

Bindweed, field; Dandelion, common; Dock, curly; Smartweed, swamp; Thistle, Canada.

Metribuzin 75DF + Marksman

Bindweed, field; Dandelion, common; Dock, curly; Smartweed, swamp; Thistle, Canada.

Metribuzin 75DF + Pursuit

Thistle, Canada



**PREPLANT AND PREEMERGENCE**

Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota, and Wisconsin

Metribuzin 75DF is recommended for additional residual control of certain broadleaf weed species in corn when applied as a tank mix combination with both grass and broadleaf herbicides registered and recommended for use in field corn. Metribuzin 75DF can be tank mixed with recommended rates of the following herbicides:

| | | |
|--------------------|---------------|-----------------|
| Alachlor | Dual | Prowl |
| Atrazine | Dual II | Pursuit Plus* |
| Banvel | Frontier | Pursuit* |
| Bicep | Guardman | Ramrod |
| Bicep II | Harness Extra | Ramrod/atrazine |
| Bicep Lite | Lariat | Simazine |
| Bladex | Linex | Surpass |
| Broadstrike + Dual | Linuron | Surpass 100 |
| Broadstrike Plus | Lorox | Topnotch |
| Bullet | Marksman | |
| Clarity | Pentagon | |

* Use only on Pursuit resistant/tolerant corn hybrids (IMI corn).

APPLICATION: Metribuzin 75DF may be applied to field corn preplant without incorporation up to 30 days prior to planting or preemergence. Applications may be made by either ground or aerial equipment. For tank mixes, follow the most restrictive application methods of all products used.

SPECIAL PRECAUTIONS:

- Do not apply more than 5 1/3 ounces Metribuzin 75DF (0.25 pound active ingredient) per acre per growing season.
- Do not apply on soils having pH 7.0 or greater.
- Corn seed should be planted a minimum of 1 1/2 inches deep.

- Metribuzin 75DF may only be used in hybrid seed corn production fields if both inbred parents are known to be tolerant to Metribuzin 75DF.
- Not recommended for use on muck soils as reduced weed control may result.
- Observe all precautions and limitations on labeling of all products used in tank mixes.

Feeding Restrictions: Corn treated with Metribuzin 75DF may be harvested for silage or grain 60 days after treatment. For tank mixes, follow the most restrictive preharvest interval of all products used.

Weeds Controlled: Metribuzin 75DF will aid in the residual preemergence control of the following weed species when tank mixed with other registered grass and/or broadleaf corn herbicides:

| | | |
|-----------------------|-------------------------|-----------------|
| Horseweed/marestail | Pigweed spp. | Sunflower |
| Ladythumb | Ragweed, common | Velvetleaf |
| Lambsquarters, common | Smartweed, Pennsylvania | Waterhemp, tall |

* For control of emerged weeds refer to the BURNDOWN WEED CONTROL section of the Metribuzin 75DF label.

| METRIBUZIN 75DF FIELD CORN RATE RECOMMENDATIONS | | |
|---|--|-----------------------|
| States | Application Timing | Metribuzin 75DF Oz./A |
| Iowa Kansas Missouri | Preplant (0 to 30 days) Preemergence | 2 to 5 1/3 |
| Illinois Indiana Kentucky Minnesota | Michigan Ohio Wisconsin | 2 to 5 1/3 |
| | Preplant (0 to 9 days) Preemergence | 2 to 4 |
| REMARKS: Apply as a broadcast spray prior to corn emergence from the soil. Do not apply Metribuzin 75DF on coarse textured soils with less than 1.5% organic matter. Do not apply more than 4 oz. Metribuzin 75DF per acre on soils with less than 2.0% organic matter. For heavy weed infestations and/or early preplant applications, use the higher rates of Metribuzin 75DF. | | |
| Consult the label of herbicide tank mix partners to determine proper use rates for the other product(s). | | |





SWEET CORN

PREPLANT AND PREEMERGENCE APPLICATIONS

Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, Ohio, South Dakota, and Wisconsin

Metribuzin 75DF is recommended for additional residual weed control of certain broadleaf weed species when applied in combination with other broadleaf and/or grass herbicides as a tank mixture. All products used must be labeled for use on sweet corn. The most restrictive restrictions and precautions of all the products used must be observed. Use only labeled rates and methods of applications.

Tank Mixtures: Metribuzin 75DF can be tank mixed with the products containing one or more of the following herbicides: 2,4-D, alachlor, atrazine, glyphosate, linuron, metolachlor, metribuzin, paraquat, and pendimethalin.

Weeds Controlled: Refer to the PREPLANT AND PREEMERGENCE APPLICATION- FIELD CORN section of this label for a list of weeds controlled by Metribuzin 75DF when applied before weed emergence. Use recommended adjuvants when emerged weeds are present. Refer to the BURNDOWN WEED CONTROL- FIELD CORN section for a list of weeds controlled and weed height restrictions.

Sequential Applications: Sequential applications of all herbicides containing metribuzin (the active ingredient in Metribuzin 75DF) are subject to a limitation of not more than 0.25 pounds a.i. of metribuzin (5 1/3 ounces of Metribuzin 75DF) per acre of corn per use season. There are no other specific restrictions on sequential applications due to the application of Metribuzin 75DF.

SPECIAL PRECAUTIONS:

- Do not apply more than a total of 5 1/3 ounces Metribuzin 75DF (0.25 pounds metribuzin) per acre per growing season.

- Do not apply preplant or preemergence on soils having a pH of 7.0 or greater.
- Corn seed should be planted a minimum of 1 1/2 inches deep.
- Metribuzin 75DF may only be used in hybrid seed production fields, if both inbred parents are known to be tolerant to Metribuzin 75DF.
- Reduced residual weed control may result when used on organic soils. For this reason, residual weed control is not claimed on organic soils.
- Observe all precautions and limitations on labeling of all products used in tank mixtures.

Feeding Restrictions: Grain, forage, and processing waste may be fed to livestock if harvested at least 60 days after the last application of Metribuzin 75DF.

Sensitive Sweet corn Hybrids: Make applications only to hybrids that have established tolerance to the application planned.

Application Methods and Timing: Metribuzin 75DF can be applied preplant surface or preemergence as a broadcast or band application in water, fluid fertilizer, or impregnated on dry fertilizer. Ground or aerial equipment may be used. See DIRECTIONS FOR USE section of this label for directions.

Application Rate Recommendations: Refer to the DIRECTIONS FOR USE section of this label for definitions of SOIL TEXTURE GROUP and other information that applies to all applications. Use the lowest rate of the recommended rate range on soils with the lowest percent clay and organic matter for the group and progressively higher rate for increased clay and organic matter content. The clay content is at least twice as important as organic matter when adjusting rates. Rates will vary based on local conditions.

| SOIL TEXTURE GROUP | SOIL ORGANIC MATTER CONTENT | |
|--------------------|-----------------------------|------------------|
| | 1.6% to 2.9% | 3.0% or More |
| All Sand Soils | DO NOT USE | |
| Coarse | 1.6 to 2.4 oz./A | 2.5 to 2.8 oz./A |
| Medium | 3 to 3.3 oz./A | 3.2 to 3.7 oz./A |
| Fine | 3.6 to 4.0 oz./A | 3.6 to 4.4 oz./A |

For early preplant application more than 9 days before planting and fields with at least 30% crop residue on the soil surface at application, the application rate may be increased 1 oz./A but not to exceed 5 1/3 oz./A.

For band applications use proportionally less per planted acre.

See DIRECTIONS FOR USE section of this label.

GARBANZO BEANS (Chickpeas)

California, Idaho, Oregon, and Washington

Special Conditions of Sale for Use on Garbanzo Beans (Chickpeas): The following directions for use were developed under the direction of IR-4 (government minor crops use program). As such the testing was done independently from the testing program of Makhteshim Agan of North

America, Inc. Buyer is advised that Makhteshim Agan makes no assurances regarding satisfaction with the product and that all risks of crop injury or product performance are assumed by the Buyer.

Metribuzin 75DF herbicide is recommended as a preemergence application for the suppression of certain broadleaf weeds in garbanzo beans.

| WEEDS SUPPRESSED* | |
|----------------------|----------------|
| Common Chickweed | Henbit |
| Common Lambsquarters | Pigweed |
| Dog Fennel (Mayweed) | Shepherdspurse |
| Field Pennycress | Wild Mustard |

* Suppression is a reduction in weed size and growth compared to a non-treated area in the same field. Metribuzin 75DF used alone will not control triazine-resistant weed species.





| RECOMMENDED APPLICATIONS | |
|--------------------------|--|
| Crop | Metribuzin 75DF Lb./Acre |
| Garbanzo beans | 1/3 to 1/2 |
| | Apply specified dosage in a single preemergence application using 10 to 40 gallons of water per acre with ground spray equipment. Apply before or after planting but before crop emergence. Thorough incorporation, either by rainfall or by mechanical means, is essential for weed suppression. Under dry conditions, incorporate Metribuzin 75DF into the top 1 to 2 inches of soil with spike harrows or similar shallow incorporation equipment, then cross harrow to insure uniform soil incorporation. Where soil surface is moist at the time of application and rain follows before weed emergence, a broadcast application should provide adequate weed suppression. Use on coarse-textured soils, sandy soils, or any soil with less than 1.5% organic matter will likely cause crop injury. Use the higher rate on fine-textured soils (high in clay or organic matter) and in fields with a history of high weed populations. |

SPECIAL PRECAUTIONS: Crop injury may result if crop is under stress conditions caused by cold weather, poor soil fertility, disease, or insect damage.

Crop injury may result if application is followed by heavy rain. Avoid application of more than 1/2 inch of irrigation within one month after application of Metribuzin 75DF or crop injury may occur.

Do not use on clay knobs of poorly covered subsoils.

Do not apply preemergence on shallow seedings less than 2 inches deep.

Do not graze or feed treated vines to livestock within 40 days after application.

Maintain continuous spray tank agitation to keep material in suspension. Avoid overlapping of spray swaths and shut off spray booms while turning, slowing, or stopping, or crop injury will occur.

NOTE: This treatment may cause some chlorosis or minor necrosis. Because garbanzo bean varieties may vary in their susceptibility to Metribuzin 75DF, determine crop tolerance prior to adoption as a field scale practice to prevent possible injury.

LENTILS AND PEAS

Idaho, Oregon, Washington, Montana, and North Dakota

Metribuzin 75DF herbicide is recommended as a preemergence and postemergence application for the suppression of certain broadleaf weeds in lentils and peas.

| WEEDS SUPPRESSED* | |
|---|--|
| Common Chickweed** Lambsquarters Dog Fennel Shepherds-purse** Field Pennygrass Wild Mustard | Henbit** Corn Spurry Redroot Pigweed Pennsylvania Smartweed Pineapple Weed Prostrate Knotweed |
| * Suppression is a reduction in weed size and growth compared to a non-treated area in the same field. Metribuzin 75DF used alone will not control triazine-resistant weed species. | |
| ** Preemergence application only. | |

PREEMERGENCE APPLICATION: Make a single preemergence application of Metribuzin 75DF at 1/4 to 1/2 lb. per acre per crop year. Apply in 10 or more gallons of water per acre with ground spray equipment or 5 or more gallons of water per acre with aerial spray equipment. Apply Metribuzin 75DF before or after planting. Thorough incorporation either by rainfall or by mechanical means is essential for weed suppression. Under dry conditions, incorporate Metribuzin 75DF into the top 1 to 2 inches of soil with spike harrows or similar shallow incorporation equipment, then cross harrow to insure uniform soil incorporation. Where soil surface is moist at the time of application and rain follows before weed emergence, a broadcast application should provide adequate weed suppression.

Use the higher rate on fine-textured soils (high in clay or organic matter) and in fields with a history of high weed populations.

Metribuzin 75DF may be applied pre- or post-plant incorporated as a tank mix combination with Fargo 4 EC. Follow the DIRECTIONS FOR USE statements on both product labels.

POSTEMERGENCE APPLICATION: One postemergence application may be made per season. Use 1/6 to 1/3 lb. of Metribuzin 75DF per acre on lentils and spring peas. On winter peas, use 1/4 to 1/3 lb. of Metribuzin 75DF per acre. For suppression of dog fennel, use 1/3 lb. Metribuzin 75DF per acre. Apply

specified dosage in 20 or more gallons of water per acre with ground spray equipment or 5 or more gallons of water per acre with aerial spray equipment. Do not exceed 40 psi with ground spray equipment. Apply as a broadcast spray when weeds are small (less than 2 inches in height or diameter) and before crop is 6 inches tall.

Temporary chlorosis of the crop may occur. There is an added risk of crop injury if a postemergence application is made following a previous preemergence or post plant incorporated Metribuzin 75DF application.

Do not apply over very moist soils or wet crop foliage. Do not apply postemergence application within 3 days after periods of cool, wet, or cloudy weather or crop injury may occur.

Do not apply within 24 hours of treatment with other pesticides.

SPECIAL PRECAUTIONS (All Applications): Do not apply more than 2/3 lb. Metribuzin 75DF per acre per year. Crop injury may result if crop is under stress conditions caused by cold weather, low fertility, disease, or insect damage.

Crop injury may also result if application is followed by heavy rain.

Do not use on coarse-textured soils, sandy soils, or soils with less than 1.5% organic matter.





Do not apply to ESTIN lentils.

Do not use on clay knobs or poorly covered subsoils.

Do not apply on shallow seedings less than 2 inches deep (preemergence only).

Do not apply within 50 days of harvest of peas or within 75 days of harvest of lentils. Do not graze or feed treated vines to livestock within 40 days after application.

Do not rotate any crop not listed on this label for 18 months following application of Metribuzin 75DF. Refer to Crop Rotation Restrictions section of this label for more information.

Maintain continuous spray tank agitation to keep material in suspension. Avoid overlapping and shut off spray booms while turning, slowing, or stopping, or crop injury will occur.

NOTE: This treatment may cause some chlorosis or minor necrosis. Because lentil and pea varieties may vary in their susceptibility to Metribuzin 75DF, determining crop tolerance prior to adoption as a field scale practice is suggested to prevent possible injury.

For additional precautions, restrictions, limitations, and sprayer clean-up information, refer to the appropriate sections of this label.

SUGARCANE-Hawaii Only

Metribuzin 75DF, a selective herbicide, is effective as a preemergence and an early postemergence broadcast application for control of certain grass and broadleaf weeds. When applied as a spot treatment, it also provides excellent control of perennial grasses and broadleaves.

Ground Application: Metribuzin 75DF should be mixed by filling the spray tank half full of clean water. Then add the recommended amount of Metribuzin 75DF to suit the total tank capacity and the rate of application per acre (preferably 25 to 35 gallons per acre). Complete filling the tank and maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture.

Aerial Application: Metribuzin 75DF is recommended for use in aerial spray equipment as a preemergence or postemergence application to irrigated sugarcane. Aerial spray equipment should be calibrated to apply the proper amount of Metribuzin 75DF in 5 to 10 gallons of spray mixture per acre.

For aerial and chemigation application methods on sugarcane the maximum application rate is 2 2/3 lb. Metribuzin 75DF/acre.

To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

Metribuzin 75DF applied preemergence or postemergence to the sugarcane as a broadcast spray or spot treatment will effectively control the following when weeds are less than 3 inches in height.

| WEEDS CONTROLLED IN IRRIGATED AND NON-IRRIGATED SUGARCANE | |
|---|--|
| Broadleaves | Grasses |
| Amaranth, spiny (<i>Amaranthus spinosus</i>) | Crabgrass (<i>Digitaria</i> spp.) |
| Euphorbia, wild (<i>Euphorbia</i> spp.) | Guineagrass (<i>Panicum maximum</i>) |
| Fireweed (<i>Erechtites hieracifolius</i>) | Plushgrass (<i>Chloris radiata</i>) |
| Floras paintbrush (<i>Emilis sonochifolia</i>) | Ricegrass (<i>Oryzopsis humenoides</i>) |
| Spurge, Garden (<i>Euphorbia hirta</i>) | Wiregrass (<i>Eleusine indica</i>) |
| Spurge, Graceful (<i>Euphorbia glomerifera</i>) | |
| WEEDS CONTROLLED IN IRRIGATED SUGARCANE ONLY | |
| Broadleaves | Grasses |
| Amaranth, Spleen (<i>Amaranthus dubius</i>) | Alexandergrass (<i>Brachiaria plantaginea</i>) |
| Haole Koa (<i>Leucaena leucocephata</i>) | Bristly foxtail (<i>Setaria verticillata</i>) |
| Hialoa (<i>Waltheris Americana</i>) | |
| Hilahila (<i>Mimosa pudica</i>) | |
| Purslane, common (<i>Portulaca oleracea</i>) | |
| Rattlepod (<i>Crotalaria spectabilis</i>) | |
| WEEDS CONTROLLED IN NON-IRRIGATED SUGARCANE ONLY | |
| Broadleaves | |
| Ageratum (<i>Ageratum conyzoides</i>) | |
| Richardia (<i>Richardia brasiliensis</i>) | |
| Tarweed (<i>Cuphea carthagenesis</i>) | |





| SUGARCANE (HAWAII ONLY) | |
|---|---|
| RECOMMENDED BROADCAST APPLICATIONS | |
| Metribuzin 75DF Lb./Acre | Remarks |
| 2 2/3 to 5 1/3 (Non-irrigated) | Preemergence (Irrigated and non-irrigated sugarcane): Apply specified dosage per acre as a broadcast spray to the soil surface. Applications should be made within two weeks after planting prior to cane emergence or shortly after emergence (spike stage). |
| 5 1/3 to 8 (Irrigated) | OR Early Postemergence (Irrigated and non-irrigated sugarcane): Apply specified dosage per acre as a broadcast spray over the cane. Application may be delayed as long as 4 to 6 weeks after planting provided weeds are less than 3 inches in height. |
| 2 2/3 to 5 1/3 | OR Postemergence: Apply specified dosage per acre as a broadcast spray to control weeds prior to "close in" time when cane shades out the weed growth. |
| 3 1/3 to 6 2/3 | Spot Treatment: Apply specified dosage in 30 to 50 gallons of finished spray per acre. Spot treatments may be used to control weeds in missed areas, corners of fields, or areas of hard to control weeds. |
| NOTE: Do not apply more than 10 2/3 lb. of Metribuzin 75DF (8 lb. active ingredient) per acre per crop cycle regardless of the method of application. The last application may be made up to 17 months of harvest. | |

SPECIAL PRECAUTIONS: Do not use treated foliage for feed or forage.

| SUGARCANE-Louisiana and Texas Only | |
|---|--|
| Preemergence and postemergence applications of Metribuzin 75DF with aerial or ground spray equipment are recommended for control of the following weeds in sugarcane in Louisiana and Texas: | |
| Broadleaves Amaranth, spiny (<i>Amaranthus spinosus</i>) Bindweed, Field (<i>Convolvulus arvensis</i>) Chickweed (<i>Cerastium vulgatum</i>) Henbit (<i>Lamium amplexicaule</i>) Lambsquarters (<i>Chenopodium album</i>) London rocket (<i>Sisymbrium irio</i>) Marestalk (<i>Conyza Canadensis</i>) Mustard, Wild (<i>Brassica kaber</i>) Pigweeds (<i>Amaranthus</i> spp.) Purslane (<i>Portulaca oleracea</i>) Sowthistle (<i>Sonchus</i> spp.) | Grasses Broadleaf Signalgrass (<i>Brachiaria platyphylla</i>) Crabgrass (<i>Digitaria</i> spp.) Foxtails (<i>Setaria</i> spp.) Johnsongrass, Seedling (<i>Sorghum halepense</i>) Oats, Winter (<i>Avena</i> spp.) |

| SUGARCANE-RECOMMENDED APPLICATIONS-Louisiana and Texas Only | |
|---|--|
| Metribuzin 75DF Lb./Acre | Remarks |
| 2 to 4 | Broadcast: Apply specified dosage per acre using 20 to 30 gallons of water with ground equipment or 5 gallons of water with aircraft spray equipment. Apply as a broadcast spray during the fall after planting or to the stubble after harvest. Make a second application early in the spring. |
| 1 to 2 | Band: Apply specified dosage in 10 to 20 gallons of water per acre in a 30 to 36 inch band over the row during the fall after planting or to the stubble after harvest. Make a second application early in the spring. |

SPECIAL PRECAUTIONS (Louisiana and Texas only): Do not use treated foliage for feed or forage. Use the higher rate on heavy clay soil and soil with a high percentage of organic matter. If necessary, a third application may be made in late spring at lay-by. Do not apply within 60 days of harvest.

| SUGARCANE-Florida Only | |
|--|--|
| Postemergence over-the-top or directed spray applications of Metribuzin 75DF are recommended for the control of the following weeds in sugarcane in Florida. | |
| Broadleaves Amaranth, spiny (seedling) (<i>Amaranthus spinosus</i>) Butlerweed (Cressleaf groundsel)(<i>Senecio glabellus</i>) Cudweed (<i>Gnaphalium</i> spp.) Purslane (<i>Portulaca oleracea</i>) | Grasses *Crabgrass, large (<i>Digitaria sanguinalis</i>) Foxtail, bristlegrass (<i>Setaria magna</i>) Goosegrass (<i>Eleusine indica</i>) Panicum, broadleaf (<i>Panicum adspersum</i>) Signalgrass, broadleaf (<i>Brachiaria platyphylla</i>) |





| RECOMMENDED APPLICATIONS | |
|--------------------------|---|
| Metribuzin 75DF Lb./Acre | Remarks |
| 1 1/3 to 2 2/3 | <p>Ground Application: Metribuzin 75DF may be used in one or two applications with a minimum of 14 days between each application. Apply when weeds are less than 6 inches tall in 10 to 40 gallons of spray mixture per acre.</p> <p>Postemergence Broadcast or Band: Apply over-the-top of stubble or plant cane while sugarcane is less than 14 inches tall.</p> <p>Postemergence Directed Spray: Apply to sugarcane that is a minimum of 14 inches tall and before row closing.</p> |
| 1 1/3 to 2 | <p>Aerial Application: Apply when weeds are less than 4 inches tall in 5 to 10 gallons of spray mixture per acre. Apply to stubble or plant cane while the sugarcane is less than 14 inches tall.</p> |

METRIBUZIN 75DF PLUS ATRAZINE TANK MIX: Metribuzin 75DF may be used with atrazine as a preemergence or postemergence (before row closing) application to sugarcane. Rates for Metribuzin 75DF are 1 to 2 2/3 lb./acre and atrazine 80% WP (4L) are 2 1/2 to 5 lb./acre (2 to 4 qt./acre). For additional information on precautions, instructions, limitations, application, and weeds controlled, refer to this label and the atrazine label.

SPECIAL PRECAUTIONS (Florida Only): Do not use more than 2 2/3 lb. per acre in a single growing season. Do not use on sand soils.

Spray contact with sugarcane foliage may result in minor leaf margin chlorosis and/or necrosis.

Do not apply within 60 days of harvest. Do not use treated crop for feed or forage.

Avoid spray overlaps or variations in application speed that may result in insufficient or excessive rates of application.

TOMATOES

Apply Metribuzin 75DF herbicide with ground equipment to seeded and transplanted tomatoes as specified below under RECOMMENDED APPLICATIONS.

Aerial application is prohibited.

For effective control of grasses and broadleaf weeds with postemergence applications, apply Metribuzin 75DF before weeds are 1-inch tall. Thorough spray coverage on weed foliage is essential for adequate control with postemergence applications.

Do not use air blast or other high pressure spray equipment to make postemergence applications of Metribuzin 75DF. Refer to the appropriate section of the label for additional information regarding spray equipment, dilution rates, mixing, sprayer cleanup, restrictions, container disposal, and cautions.

For specific application information see the GENERAL INFORMATION section in the front of this label.

| WEEDS CONTROLLED | |
|---|--|
| PREPLANT INCORPORATED APPLICATIONS TRANSPLANT TOMATOES ONLY Broadcast Sprays-1/3 to 2/3 lb. Metribuzin 75DF/Acre | |
| <p>Broadleaves Galinsoga (<i>Galinsoga</i> spp.) Lambsquarters (<i>Chenopodium album</i>) *Pigweed, Redroot (<i>Amaranthus retroflexus</i>) *Purslane, Common (<i>Portulaca oleracea</i>)</p> | <p>Grasses *Goosegrass (<i>Elyusine indica</i>)</p> |
| <p>Preplant incorporated applications applied as directed will suppress foxtails, panicums, and barnyardgrass.</p> <p>Metribuzin 75DF/Trifluralin Tank Mix: This tank mix combination applied preplant incorporated as directed on this label will control the weeds listed above plus those weeds listed on the trifluralin label.</p> | |
| <p>* For optimum control of these weeds, use the highest rate recommended on the label for the type of application to be made. Repeat postemergence applications may be needed for best control. Postemergence applications as directed on this label will suppress barnyardgrass and crabgrass when these weeds are less than 1-inch tall.</p> | |





| WEEDS CONTROLLED | |
|---|---|
| POSTEMERGENCE APPLICATIONS- ESTABLISHED TOMATOES | |
| For effective control of weeds with postemergence applications, apply Metribuzin 75DF before weeds are 1-inch tall. | |
| Broadcast Sprays- 1/3 to 2/3 lb. Metribuzin 75DF/Acre Broadleaves Carpetweed (<i>Mollugo verticillata</i>) Fumitory (<i>Fumaria officinalis</i>) Galinsoga (<i>Galinsoga</i> spp.) *Jimsonweed (<i>Datura stramonium</i>) *Ladysthumb (<i>Polygonum persicaria</i>) Lambsquarters (<i>Chenopodium album</i>) Mustard, Wild (<i>Brasica kaber</i>) Pigweeds (<i>Amaranthus</i> spp.) Purslane (<i>Portulaca oleracea</i>) *Ragweed, common (<i>Ambrosia artemisiifolia</i>) *Smartweed, Pennsylvania (<i>Polygonum pennsylvanicum</i>) Toadflax (<i>Linaria</i> spp.) *Velvetleaf (<i>Abutilon theophrasti</i>) | Directed Sprays- 2/3 to 1 1/3 lb. Metribuzin 75DF/Acre Grasses *Foxtail, Yellow (<i>Setaria glauca</i>) Goosegrass (<i>Eleusine indica</i>) Plus Weeds Listed Under Broadcast Sprays |
| * For optimum control of these weeds, use the highest rate recommended on the label for the type of application to be made. Repeat postemergence applications may be needed for best control. Postemergence applications as directed on this label will suppress barnyardgrass and crabgrass when these weeds are less than 1-inch tall. | |

| RECOMMENDED BROADCAST APPLICATIONS FOR TOMATOES | |
|---|--|
| Metribuzin 75DF *Lb./Acre | Remarks |
| 1/3 to 2/3 | Preplant Incorporated-Transplant Tomatoes Only: Apply specified dosage in 10 or more gallons of water per acre as a broadcast spray to the soil surface immediately before transplanting. Incorporate to a depth of 2 to 4 inches with equipment capable of uniformly mixing the chemical into the soil. This application may be made alone or in a tank mix combination with trifluralin. When transplanting tomatoes, place the root system of the plants below the herbicide incorporation zone or injury may occur. Refer to the trifluralin label for specific rate of application and for additional precautions and restrictions for tomatoes. |
| 1/3 to 2/3 | Postemergence Broadcast Spray-Established Tomatoes: Apply specified dosage in 20 or more gallons of water per acre as a broadcast spray or apply in 1/4 to 3/4 inch of water (use 1/4 to 1/2 inch of water on sandy soils) per acre as a continuous injection in center pivot and lateral move systems or apply in the last 15 to 30 minutes of set in permanent solid set sprinkler systems. One or more applications may be applied per use season. Allow at least 14 days between applications or severe crop injury may occur. For transplanted tomatoes, do not apply until transplants have recovered from transplant shock and new growth is evident. Do not apply to tomatoes within 24 hours of application of other pesticides. Do not tank mix with other pesticides. (See SPECIAL PRECAUTIONS below.) |
| 2/3 to 1 1/3 | Postemergence Directed Spray-Established tomatoes: Apply specified dosage in 20 or more gallons of water per acre as a directed spray. One or more applications may be applied per use season. Allow at least 14 days between applications or severe crop injury may occur. Avoid contacting tomato foliage with spray. This method of treatment is recommended for use in fields with a history of severe weed pressure or in fields infested with hard-to-control weeds. For transplanted tomatoes, do not apply until transplants have recovered from transplant shock and new growth is evident. Do not apply to tomatoes within 24 hours of application of other pesticides. (See SPECIAL PRECAUTIONS below.) When banding, see the appropriate section in the front of this label. |
| * Use the higher rate in fields with a history of severe weed pressure and for maximum residual weed control. | |

SPECIAL PRECAUTIONS (Tomatoes): Do not apply more than a total of 1 1/3 lb. Metribuzin 75DF per crop season. Do not apply the total amount of 1 1/3 lb. Metribuzin 75DF within a time span of less than 35 days except in the case of directed sprays.

Allow at least 14 days between applications regardless of dosage or method of application or severe crop injury may occur.

Do not apply within 7 days of harvest.

Do not apply within 3 days after periods of cool, wet, or cloudy weather or crop injury will occur.

Do not use hot caps on tomatoes within 7 days before or at any time after

application of Metribuzin 75DF. Do not treat seeded tomatoes until plants have reached the 5- to 6-leaf stage or severe crop injury may occur.

Crop injury or delayed maturity may result from broadcast or directed spray applications if tomatoes are growing under stress conditions such as periods of drought or cool, wet, and cloudy weather preceding application.

For newly introduced tomato varieties with unknown tolerance to Metribuzin 75DF, treat only a small area to determine if Metribuzin 75DF can be used without injury to the crop.

DO NOT USE METRIBUZIN 75DF ON TOMATOES IN KERN COUNTY, CALIFORNIA.



TOMATOES (CALIFORNIA ONLY)

Apply Metribuzin 75DF as a ground application for chemical fallow on fields to be planted with tomatoes. Refer to the appropriate section of this product label for additional information regarding spray equipment, mixing, sprayer cleanup, restrictions, container disposal and cautions. Do not apply this product through any type of irrigation system. Aerial application is prohibited.

WEEDS CONTROLLED: Common chickweed, Lambsquarters, London rocket, Purslane, Redroot pigweed, Shepherdspurse, Wild mustard

| APPLICATION INSTRUCTIONS | |
|---|-------------|
| DOSAGE | 2/3 to 1 lb |
| BROADCAST SPRAY: Use lower rate on soils with 1/2 to 2% organic matter and higher rates on soils with more than 2% organic matter. Apply specified dosage per acre by ground equipment using a minimum of 10 gallons of water per acre. One fallow application may be made up to 90 days prior to planting. At least 2 inches of rain or equivalent irrigation are required between application and planting. Additional cultivation may be required for weed control and seed bed preparation prior to planting. | |

SPECIAL PRECAUTIONS:

Applications should not be made when weather conditions favor spray drift especially in areas where sensitive crops are growing in adjacent fields, or injury may occur. Light soils, alkaline soils or water, soil high in soluble salts, cold germination and/or growing conditions and interaction with other herbicides may harm tomatoes and require lower rates or longer preplant intervals.

Apply with properly calibrated equipment to assure uniform coverage of soil. Spray booms must be shut off while starting, turning, slowing or stopping, or crop injury may result. Do not overlap spray applications. Continuous agitation is required in the spray tank to keep material in suspension. Avoid land-planing, leveling etc., between Metribuzin 75DF application and planting as concentrating treated soil may cause injury.

Do not apply more than a total of 2/3 pound of Metribuzin 75DF or 1 pint of METRI 4F as a broadcast or directed spray on established tomatoes following the application of Metribuzin 75DF as a chemical fallow spray. Observe all precautions contained in the EPA registered label for use on tomatoes when making such applications.

DO NOT USE METRIBUZIN 75DF ON TOMATOES IN KERN COUNTY, CALIFORNIA.

CEREALS-Spring and Winter Barley and Winter Wheat

Metribuzin 75DF herbicide is recommended for control or suppression of certain grasses and broadleaf weeds when applied postemergence to spring and winter barley or winter wheat. Metribuzin 75DF alone and several tank mixture treatments are recommended for use in the following states: AR, GA, ID, IL, IN, KS, KY, LA, MS, MO, MT, NV, OH, OK, OR, TN, TX, UT, WA.

Mixing: See the GENERAL INFORMATION section of this label for specific mixing procedures. When tank mixing, carefully follow the instructions on this label. Refer to the other product labels registered for use in barley and winter wheat for additional use directions, rates, weeds controlled, and restrictions.

Application: Metribuzin 75DF may be applied by aerial or ground application equipment. Use a minimum spray volume of 2 GPA by air and 10 GPA by ground. Uniform spray coverage is necessary to obtain optimum weed control and to minimize potential for crop injury. Do not exceed rate specified on this label. Do not apply Metribuzin 75DF through any type of irrigation equipment. Apply Metribuzin 75DF when the crop is healthy and actively growing.

Metribuzin 75DF may be applied more than once per crop season. Allow a minimum of 21 days between applications if wheat is actively growing or allow 45 days between applications if wheat is growing in adverse conditions, has entered dormancy, or is stressed due to frost damage, disease, drought, or excessive moisture. Do not use on soils containing less than 0.75% organic matter. Do not apply more than a total of 10.66 ounces Metribuzin 75DF (8 ounces active ingredient) per acre per year. On irrigated cereals, do not apply more than 0.5 inch of water for the first irrigation. The maximum amount for each additional irrigation should not exceed 1 inch. Allow a minimum of 14 days between the first irrigation and subsequent irrigations.

Performance Factors: Weed control may not be observed for 2 to 4 weeks under normal growth conditions and for 4 to 6 weeks under very dry conditions. Moisture (at least 1/2 inch) is required within 2 to 3 weeks after application to move Metribuzin 75DF into the weed root zone. Lack of adequate moisture after application may result in poor or erratic weed control. Control or suppression of listed weeds is dependent on weed size at time of application. Control or suppression may be reduced if broadleaf weeds are taller than 1 inch or grasses have more than 2 leaves.

Tank mixtures: Metribuzin 75DF may be tank mixed with Ally, Amber, Finesse, Glean FC, Harmony Extra, 2,4-D, MCPA, Igran, Banvel/Banvel SGF, Bronate, or Butril herbicides. A nonionic surfactant containing at least 80% active ingredient may be used in Metribuzin 75DF tank mixes with sulfonyleurea herbicides (Ally, Amber, Finesse, Glean FC, Harmony Extra). Do not use a crop oil concentrate or any adjuvant containing vegetable or petroleum oils with any Metribuzin 75DF mix as crop injury may result. Additional pesticides may also be tank mixed with Metribuzin 75DF unless specifically prohibited on the mix products' label. In some instances, combinations with organophosphate insecticides may cause temporary leaf yellowing and/or crop injury, especially when widely fluctuating day/night temperatures occur near application. Always refer to the other product labels registered for use on cereals for additional directions, rates, and weed species controlled. Observe all precautions and limitations on labeling of all products used in mixtures.

Feeding Restrictions: Do not graze wheat within 14 days of Metribuzin 75DF application or, harvest grain within 21 days after last application. Do not graze or harvest barley before crop maturity. For tank mix combinations, follow the most restrictive label.

SPECIAL PRECAUTIONS: Cereal Injury-Crop injury may occur if Metribuzin 75DF is applied:

1. When the crop is under stress such as winterkill, frost damage, disease, drought or excessive moisture, severe grazing, or when these conditions follow the application.
2. In combination with fluid fertilizer especially with the addition of surfactant.
3. Prior to the growth stage specified on this label.
4. To soils high in lime or sodium, a pH greater than 7.7, calcareous, gravelly, thinly covered, or exposed subsoil areas.
5. To fields where cereal seeds have been planted less than 1 inch deep.
6. To a non-winter hardy wheat or barley variety.
7. To a sensitive wheat or barley variety as listed below.
8. To frozen soil or crop still in winter dormancy.

Cereal Rotations Following Potatoes Treated with Metribuzin 75DF: If planting a sensitive cereal variety (listed under the wheat and barley variety tolerance portion of this label), following potatoes treated with Metribuzin 75DF or metribuzin-containing products, refer to the potato section of the Metribuzin 75DF label for special cultural practices to follow.



APPLICATION RECOMMENDATIONS

Metribuzin 75DF alone or in a tank mix with labeled broadleaf herbicides may be applied by aerial or ground spray equipment as a broadcast postemergence spray.

| RECOMMENDED POSTEMERGENCE BROADCAST APPLICATIONS OF METRIBUZIN 75DF | | | |
|---|--------------|--|-------------|
| CROP GROWTH STAGE | SOIL TEXTURE | METRIBUZIN 75DF RATE (oz./A) % ORGANIC MATTER | |
| | | 0.75 TO 2.0 | OVER 2.0 |
| 2 Leaf To 2 Tiller | Coarse | 1 to 2 | 1 to 3 |
| | Medium | 1 to 3 | 2 to 3 |
| | Fine | 2 to 3 | 2 to 4 |
| Use these rates on crops with secondary roots smaller than 1 inch. For dryland winter wheat (nonirrigated), apply the highest recommended rate to achieve maximum weed suppression/control. | | | |
| 3 Tiller To 4 Tiller | Coarse | 3 to 4 | 4 to 5 |
| | Medium | 4 to 5 | 5 to 6 |
| | Fine | 5 to 6 | 5 to 6 |
| Do not apply within 2 weeks after grazing or breaking of winter dormancy. Apply after the crop is at or beyond the 3-tiller growth stage but before jointing. Secondary roots should be developed and larger than 1-inch long. Do not apply before 75 days after planting. For dryland winter wheat (nonirrigated), apply the highest recommended rate to achieve maximum weed suppression/control. GEORGIA ONLY: Wheat must be planted before November 15 in the Piedmont area and Northern part of the state and before December 1 in the Coastal Plain area. | | | |
| Over 4 Tillers | Coarse | 4 to 6 | 5 to 8 |
| | Medium | 4 to 8 | 5 to 8 |
| | Fine | 5 to 8 | 8 to 10 2/3 |
| Do not apply within 2 weeks after grazing or breaking of winter dormancy. Apply after the crop is at or beyond the 3-tiller growth stage but before jointing. Secondary roots should be developed and larger than 1-inch long. Do not apply before 75 days after planting. For dryland winter wheat (nonirrigated), apply the highest recommended rate to achieve maximum weed suppression/control. GEORGIA ONLY: Wheat must be planted before November 15 in the Piedmont area and Northern part of the state and before December 1 in the Coastal Plain area. | | | |

WHEAT AND BARLEY VARIETAL TOLERANCE*

Wheat and barley varieties vary in their tolerance to Metribuzin 75DF. Varieties below are **tolerant** to and are **recommended** for use with Metribuzin 75DF:

Winter Wheat: Abe, AgriPro Mason, AgriPro Shiloh, Arthur, AS 7846, AS 7853, Baker Seed 32, Barbie VI, Basin, Batur, Bayles, Becker, Bintee V, Buchshot DS 2368, Caldwell, Cardinal, Cashup, Centurk, Cherokee, Cheyenne, Clark, Coker 747, Coker 762, Coker 797, Coker 68-15, Coker 9134, Coker 9543, Coker 9904, Coker 9907, Daws, DB 533W, DB 562W, DB 580W, Delta King 502, Delta King 9027, Dixie 952, Doublecrop, Dusty, Dyna-gro 426, Dynasty, Excel, Faro, FFR 525W, Florida 302, FS 432, FS 433, FS 435, Gains, Garst 64, Georgia 100, Genie V, Hatton, Hawk, Hill 81, Howell, Hunter, Hyak, Hyslop, Katie VI, KY 16-2, Larned, Lewis 833, Lewjain, Lisa, Longhorn, Luke, Madsen, Magnum, Malcom, McDermid, McNair 1003, McNair 1813, Molly, Moro, Neely, Nelson, Newton, Norstar, Norwin, Nugaines, Oasis, Omega 78, Paha, Peck, Pike, PI 2157, PI 2180, PI 2510, PI 2545, PI 2548, PI 2550, PI 2552, PI 2555, PI 2566, PI 2571, PI 2580, PI 2684, Quantum 577, Redwin, Rocky, Saluda, Sawyer, SC 104, Siouxland, Sprague, Southern Belle, Stacy, Stallion, Stephens, TAM W101,

TAM 105, TE 877, TE 2548, TE SR204, Timber, Tomahawk, TR 8555, TR 8557, TR 8768, Traveler, Tres, Tye, Tyler, Verne, Victory, Wakefield, Wanser, Weston, Winalta, Wrangler.

Barley: Advance, Boyer, Clark, Compana, Hannchen, Hector, Hesk, Hudson, Lud, Luther, Kamiak, Klages, Olympic, Piroline, Steptoe, and Triumph.

The following cereal varieties are sensitive to Metribuzin 75DF and are not recommended for use:

Winter Wheat: AgriPro Clemens, AT 90W, AT 91W, Arapaho, Baker Seed 33, Century, Cimarron, Coker 833, Coker 916, Coker 983, Coker 9024, Coker 9105, Coker 9323, Coker 9474, Coker 9663, Coker 9835, Coker 9766, Coker 9877, EK 102, EK 114, FFR 555, Florida 304, Freedom, FS 417, FS 423, FS 425, FS 430, Gore, Hazen, Hickory, Jackson, Julie III, KY 49-25, Linden, Madison, Mesa, Mustang, Pacer, PI XW 522, PI 2551, PI 2163, Pioneer 2691, Princeton 733, PSRW 71, PSR 226, PSR 278, Rosen, Savannah, Sierra, TAM 107, TR 101, TR 1011, TR 8822, Triumph 64, Vona, Wings, Winridge, Yamhill.

Spring/Durum Wheat: Avoid use on Spring wheat and Durum wheat varieties.





Barley: Glenn, Morex, Moravian 3, Larker, Summit, Bracken, Anheuser Busch B2601, and varieties with Morex parentage.

Varieties Not Listed: To avoid possible crop injury on any variety not mentioned in this label, contact a Makhteshim Agan representative or herbicide expert for a variety recommendation prior to treatment or treat a

small strip of the unlisted variety with the recommended Metribuzin 75DF rate to ascertain crop tolerance before treating an entire field.

***Abbreviated names of vendors:** AS (Agseco), AT (Agratech), DB (Diener Bros.), FS (Growmark FS), PI (Pioneer), PSR (Hybritech), SC (J.M. Schultz), TE (Terra), and TR (Terral).

WEEDS CONTROLLED

Used at recommended rates, Metribuzin 75DF will control many annual broadleaf weeds. Control is best when applied to young, actively growing weeds. Weeds controlled by Metribuzin 75DF include:

| | |
|--|---|
| Bittercress Catchfly, Conical (Sand) Catchweed (Madwort) Chickweed, Common Chickweed, Mouseear Comcockle Dogfennel (Mayweed) Evening Primrose, Cutleaf Falseflax, Smallseed Fiddleneck, Tarweed Filaree, Redstem Geranium, Carolina Gromwell, spp. Henbit | Knotweed, Prostrate Lambsquarters, Common Lettuce, Miners Mustard, Blue Mustard, Wild Pennycress, Field Pepperweed, Virginia Pigweed, spp. Pineappleweed Polemonium, Annual (Jacob's Ladder) Radish, Wild Shepherdspurse Speedwell, Ivyleaf Turnip, Wild |
|--|---|

WEEDS SUPPRESSED

Metribuzin 75DF control of the following weeds varies from poor to excellent depending on time of application, stage of growth at application, temperatures, and soil moisture conditions following treatment. For maximum effect on these weeds, apply the highest recommended rate at the earliest growth stage timing for each particular soil type and organic matter. Suppression is a reduction in weed size and growth as compared to a non-treated area in the same field.

| | |
|---|--|
| Broadleaves Buckwheat, Wild* Buttercup, spp. Cowcockle Kochia* Lettuce, Prickly | Mustard, Tumble (Jim Hill)* Tansy mustard Thistle, Russian Vetch, Winter |
| Grasses Barley, Hare (Wild) Barley, Little Blackgrass Bluegrass, Annual Bluegrass, Bulbous Brome, Downy* Brome, Japanese* | Brome, Ripgut* Cheat* Foxtail, spp.* Oat, Wild* Rescuegrass* Whitlowgrass, Spring (Vernal) Windgrass |

* Use the highest recommended Metribuzin 75DF rate for maximum weed suppression.

FOR WEED CONTROL IN A WHEAT/FALLOW/WHEAT ROTATION

Idaho, Oregon, Utah, and Washington Only

Metribuzin 75DF may be applied to provide weed control during the fallow period after wheat harvest or in the spring before winter wheat is planted. Winter wheat can be seeded 4 months (120 days) after spring application. Mechanical tillage or the application of a contact herbicide may be required

to control weeds germinating prior to seeding of winter wheat. Best results will be obtained where straw and chaff are evenly distributed across the field.

For specific application information see the GENERAL INFORMATION section in the front of this label.

Where weed growth is present at application time, Metribuzin 75DF should be applied with Gramoxone or other contact herbicide. Refer to the other product label registered for additional directions, rates, and weed species controlled.





| WEEDS CONTROLLED | |
|--|--|
| Broadleaves Chickweed, Common (<i>Stellaria media</i>) Henbit (<i>Lamium amplexicaule</i>) *Kochia (<i>Kochia scoparia</i>) Lambsquarters (<i>Chenopodium album</i>) Mustard, Blue or Purple (<i>Chorispora tenella</i>) Mustard, Jim Hill (<i>Sisymbrium altissimum</i>) | Mustard, Tansy (<i>Descurainia pinnata</i>) Mustard, Treacle (<i>Erysimum repandum</i>) Mustard, Wild (<i>Brassica kaber</i>) Pennycress, Field (Fanweed) (<i>Thlaspi arvense</i>) Pigweeds (<i>Amaranthus</i> spp.) Russian Thistle (<i>Salsola iberica</i>) Sunflower (<i>Helianthus</i> spp.) |
| Grasses Cheatgrass (<i>Bromus secalinus</i>) Downy brome (<i>Bromus tectorum</i>) *Wheat, Volunteer (<i>Triticum</i> spp.) | |

***NOTE:** Since control of these weeds may be variable depending on moisture following application, the higher labeled rate is recommended.

After Harvest Application (Fall Fallow): Metribuzin 75DF may be applied to wheat stubble after harvest in the fall. Apply 2/3 to 5/6 lb. per acre broadcast before weeds emerge. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

Do not plant crops in treated areas for at least 10 months following fall applications.

Metribuzin 75DF may be applied at 2/3 to 5/6 lb. per acre as directed above for a fall application. If other vegetation is present at the time of application use a contact herbicide.

Spring Application (Summer Fallow): Metribuzin 75DF may be applied to wheat stubble in the spring. Apply 1/2 to 2/3 lb. per acre broadcast before weeds emerge in the spring. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

Precautions and Restrictions: Do not graze treated fields.

Do not plant spring-seeded cereals following fall fallow applications of Metribuzin 75DF.

Where Metribuzin 75DF was applied in the fall, do not apply Metribuzin 75DF in the spring.

Do not rotate any crop not listed on this label for 18 months following application of Metribuzin 75DF.

FOR WEED CONTROL IN A FALLOW ROTATION WITH BARLEY AND WHEAT Colorado, Kansas, Montana, Nebraska, and Wyoming Only

Metribuzin 75DF may be applied to provide weed control during the fallow period after wheat or barley harvest or in the spring before planting of winter wheat or barley. Mechanical tillage or the application of a contact herbicide may be required to control weeds germinating prior to seeding of winter wheat or barley.

For specific application information see the GENERAL INFORMATION section in the front of this label.

Where weed growth is present at application time, Metribuzin 75DF should be applied with Gramoxone, Roundup, or other contact herbicide. Refer to the other product label registered for additional directions, rates, and weed species controlled. Do not plant crops in treated areas earlier than 10 months following fall applications.

Do not rotate any crop not listed on this label for 18 months following application of Metribuzin 75DF.

| WEEDS CONTROLLED | |
|--|--|
| Broadleaves Chickweed, Common (<i>Stellaria media</i>) Cowcockle (<i>Vaccaria pyramidata</i>) Henbit (<i>Lamium amplexicaule</i>) *Kochia (<i>Kochia scoparia</i>) Lambsquarters (<i>Chenopodium album</i>) Mustard, Blue or Purple (<i>Chorispora tenella</i>) Mustard, Jim Hill (<i>Sisymbrium altissimum</i>) | Mustard, Tansy (<i>Descurainia pinnata</i>) Mustard, Treacle (<i>Erysimum repandum</i>) Mustard, Wild (<i>Brassica kaber</i>) Pennycress, Field (Fanweed) (<i>Thlaspi arvense</i>) Pigweeds (<i>Amaranthus</i> spp.) Russian Thistle (<i>Salsola iberica</i>) Sunflower (<i>Helianthus</i> spp.) |
| Grasses Cheatgrass (<i>Bromus secalinus</i>) Downy Brome (<i>Bromus tectorum</i>) *Foxtail, Green (<i>Setaria viridis</i>) | *Wheat, Volunteer (<i>Triticum</i> spp.) *Wild Oats (<i>Avena fatua</i>) |

***NOTE:** Since control of these weeds may vary depending on moisture following application, use the higher rate recommended below.

After Harvest Application (Fall Fallow): Metribuzin 75DF may be applied to stubble after harvest in the fall. Apply 5/6 to 1 lb. per acre broadcast before weeds emerge. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

Spring Application (Summer Fallow): Metribuzin 75DF may be applied to wheat stubble in the spring. Apply 1/2 to 2/3 lb. per acre broadcast before

weeds emerge in the spring. Use the higher rate for longer weed control or weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation. Wheat or barley can be seeded 120 days after spring application.

Precautions and Restrictions: Do not graze treated fields.

Do not plant spring-seeded cereals following fall applications for fallow.

Where Metribuzin 75DF was applied in the fall, do not apply Metribuzin 75DF in the spring.



CROP ROTATION RESTRICTIONS

Waiting Period After Metribuzin 75DF Herbicide Application¹

| | | |
|--|---|---|
| 4 Months Alfalfa Asparagus Barley ² Corn Forage Grasses Sainfoin Soybeans Sugarcane ³ Tomatoes Wheat ² | 8 Months Barley Lentils Peas Wheat | 12 Months Potatoes Rice 18 Months Sugar Beets Onions And Other Root Crops not listed on this label and all other crops not listed on this label |
|--|---|---|

¹ Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed. Stand reductions may occur in some areas.

² Following peas, lentils, or soybeans.

³ Do not rotate any crop not listed on this label after application of Metribuzin 75DF to sugarcane.

FOR USE ON BENTGRASS GROWN FOR SEED AND FOR WEED CONTROL IN ESTABLISHED¹ PERENNIAL GRASSES GROWN FOR SEED

¹Established grasses are those which have been harvested at least once for seed or were planted one year or more prior to application.

For Weed Control in Established Perennial Bentgrass Grown For Seed in

| Crop | Metribuzin 75DF lb/acre | Remarks |
|--------------------------|-------------------------|---|
| Bentgrass grown for seed | 0.38 to 0.5 | Apply Metribuzin 75DF as a broadcast spray in at least 15 gallons of spray solution per acre when the volunteer grasses are in the 1- to 2-leaf growth stage following fall rainfall or irrigation and before active spring growth. Excessive crop injury and/or failure to control weeds may result if application is made after mid-February. Allow at least 120 days between application and harvest for seed. |

Application Restrictions:

- Do not apply more than once per year.
- Do not apply to a crop that is under stress, for example, from disease, severe insect damage, nutrient deficiency, cool to cold temperatures, or deficient or excessive moisture.
- Apply only to Colonial and Creeping Bentgrass.
- Apply only to established bentgrass that is at least one year old and has been harvested for seed at least once.
- Do not tank mix with other herbicides.

Feeding Restrictions: Do not use the crop or crop residues as feed or livestock bedding for at least 28 days following the last application.

For weed control in Established Perennial Grasses Grown For Seed in Oregon West of the Cascade Mountains and in Crook, Deschutes, Jefferson, and Wasco Counties WEEDS CONTROLLED

When used as directed below, Metribuzin 75DF will reduce competition from volunteer seedlings of the indicated crop, annual *Bromus* species, annual ryegrass, and annual bluegrass. Metribuzin 75DF will control rattail fescue, henbit, ivyleaf speedwell, chickweed, mustards, and shepherdspurse. The addition of wetting agents containing crop oil may enhance control of the

Oregon West of the Cascade Mountains and in Crook, Deschutes, and Wasco Counties
WEEDS CONTROLLED

When used as directed below, Metribuzin 75DF will reduce competition from seedlings of annual *Bromus* species, annual ryegrass, and annual bluegrass. Metribuzin 75DF will control rattail fescue, henbit, ivyleaf speedwell, chickweed, mustards, and shepherdspurse.

Crop Tolerance: Crop tolerance is marginal and crop injury and yield reduction are possible. To minimize crop injury, apply when the crop is not under stress. Use of adjuvants will reduce crop tolerance. Making the application after three consecutive sunny days will reduce the potential for crop injury.

volunteer crop and grassy weeds. When adding wetting agents, follow the directions for use and recommended rates on the wetting agent label.

Metribuzin 75DF is compatible with most fertilizers, fungicides, and insecticides. Metribuzin 75DF may be combined with other herbicides for enhanced weed control. Prior to tank mixing with another herbicide, refer to the **GENERAL INFORMATION** section of the Metribuzin 75DF label booklet and a knowledgeable authority or a Makhateshim Agan representative.



APPLICATION INSTRUCTIONS

| Crop | Metribuzin 75DF lb/acre | Remarks |
|--|----------------------------|--|
| Perennial Ryegrass Tall Fescue | 1/3 to 3/4 | Apply specified dosage as a broadcast spray in at least 15 gallons of spray solution per acre when the volunteer grasses are in the 1- to 2-leaf stage following fall rainfall or irrigation but prior to active spring growth. Excessive crop injury and/or failure to control weeds may result if application is made after mid-February. Allow at least 120 days between application and harvest. |
| Bluegrass Fine Fescue Orchardgrass | 1/3 to 1/2 | |

Application Restrictions:

- Do not apply more than once per year.
- Do not apply Metribuzin 75DF through any type of irrigation system.
- Crop and crop residues may be fed to livestock or used as bedding. If the seed crop is terminated and grazed or cut for forage, allow at least 28 days between application and use as animal feed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Non-refillable Container (flexible-bag-all weights): Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available.

Non-refillable Container (rigid-fifty lbs. or less): Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **Conditions, Disclaimer of Warranties and Limitations of Liability**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.





