

SUPPLEMENTAL LABELING

DuPont Crop Protection

"..... A Growing Partnership With Nature"

DUPONT™ ASSURE® II HERBICIDE FOR USE ON FLAX AND SUNFLOWER

DUPONT™ ASSURE® II HERBICIDE

EPA REG. NO. 352-541

FOR USE ON FLAX AND SUNFLOWER

GENERAL INFORMATION

DuPont™ ASSURE® II herbicide can be used for weed control in flax and sunflower. ASSURE® II is a systemic herbicide that is rapidly absorbed by treated foliage and translocated to the roots and other growing points of the plant. In general, effects are first observed within 7 to 14 days after application depending on the grass species treated and the environmental conditions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations. Use of ASSURE® II on flax and sunflower is subject to the use precautions and limitations provided here and on the complete ASSURE® II label.

ASSURE® II herbicide may be applied postemergence to flax and sunflower. Refer to the ASSURE® II herbicide label for information regarding use rates, weeds controlled, use restrictions, rotational cropping recommendations, sprayer cleanup, use precautions and other important information.

Other suitable herbicides, fungicides, and insecticides registered for use on flax and sunflower may be tank mixed or used sequentially with ASSURE® II, provided that the recommended application timing is the same. Read and follow all manufacturer's label recommendations for the tank mix partner prior to use. The most restrictive provisions on either label apply.

IMPORTANT PRECAUTIONS

SEASONAL USE LIMITS AND PRE-HARVEST INTERVALS

Flax

- Do not apply ASSURE® II within 70 days of harvest.
- The maximum use rate of ASSURE® II is 24 fluid ounces per acre per season.
- Application intervals should be greater than 7 days.

Sunflowers

- Do not apply ASSURE® II within 60 days of harvest.
- The maximum use rate of ASSURE® II is 18 fluid ounces per acre per season.
- Application intervals should be greater than 7 days.
- Unacceptable crop injury may occur if ASSURE® II is applied to Sunflowers under stress (see Environmental Conditions and Biological Activity Section of the ASSURE® II label). Delay application until the stress passes and sunflowers have resumed growth.
- Unacceptable crop injury may occur if petroleum-based crop oil concentrates, or methylated seed oils, are used with ASSURE® II. Nonionic surfactants at 1 quart of product per 100 gallons of spray solution (0.25% v/v) are the preferred adjuvant in sunflowers.

IMPORTANT

BEFORE USING ASSURE® II AND ANY TANK-MIX OR SEQUENTIALLY APPLIED PESTICIDES, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABELS.

This label contains supplemental instructions for use of the referenced DuPont product which do not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

R-255 092606 111606

© 2006 E. I. Du Pont de Nemours and Company, DuPont Crop Protection, Wilmington, Delaware 19898