



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

ADMIRE® PRO Systemic Protectant

**For Use on Containerized Citrus
to control pests**

EPA Reg. No. 264-827

PRODUCT BULLETIN

ADMIRE® PRO Systemic Protectant

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This 2(ee) Recommendation must be in the possession of the user at the time of pesticide application.

CITRUS (Containerized)

Crops of Crop Group 10 Including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, White sapote (*Casimiroa* spp), and other cultivars and/or hybrids of these

Pests Controlled	Rate mL / "citra pot" (0.1 ft³ container media)
Aphids Blackfly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies Citrus root weevil (larval complex)	0.11

Pests Suppressed	
Citrus thrips (foliage feeding thrips only)	0.11

Notes and Restrictions

Pre-Harvest Interval (PHI): **0 day**

Maximum ADMIRE PRO allowed per application: **0.11 mLs / 0.1 ft³ container media.**

Maximum ADMIRE PRO allowed per crop season: **3.0 mLs / plant.**

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Applications

For commercial nursery production in standard "citra pot" of 0.1 ft³ volume

Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. Basal, soil drench in a minimum of 30 milliliters (mLs) total solution per "citra pot".

Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, treatment should be made at planting/transplanting prior to insect infestation. Retreat if necessary but do not apply more than 3.0 mLs per plant per season. For control of larvae of the citrus root weevil complex, application must be made prior to neonate larvae entering potting media.

Application - For citrus production with other container volumes:

Determine volume of container and calculate required dosage based on 0.50 mLs / 0.1 ft³ potting media. Apply calculated dosage per container as described above. Do not exceed rate of 3.0 mLs / plant per crop season regardless of container size.

Important

Before using this product, read and carefully observe directions, cautionary statements and other information appearing on the product packaging label. This product is sold subject to the Conditions of Sale set forth on the container label.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

Admire is a registered trademark of Bayer