

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban[®] 15G

EPA Reg. No. 62719-34

2(ee) Recommendation[†]

For Distribution and Use in the States of Alabama, Florida, Georgia, Mississippi, North Carolina, South Carolina and Virginia

Sequential Applications for Control of Southern Corn Rootworm and Lesser Cornstalk Borer in Peanuts

ATTENTION

[†] This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Lorsban[®] 15G insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of Lorsban 15G is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban 15G.

Directions for Use

Follow all applicable use directions, precautions and restrictions for peanut on the label for Lorsban 15G.

Southern Corn Rootworm Larvae

Apply **two** split applications of Lorsban 15G at 2 lb ai per acre (e.g., 15 oz per 1000 row feet on 36-inch rows), 10 to 30 days apart, to maintain southern corn rootworm (SCRW) control when **any** of the following conditions exist:

- Soil organic matter is 2% or higher
- Multiple generations of SCRW are expected
- Extended periods of infestation by SCRW are expected

To achieve maximum crop protection, make applications on a preventative basis prior to infestation. The first application may be made at-plant or postplant. The second application should generally be made 30 days following the first application and/or timed to coincide with the period when significant SCRW populations are expected. Make the second application postplant no sooner than 10 days after the first application. Do not make more than two applications of Lorsban 15G or apply more than 4 lb ai per acre per season. Refer to the product label for pest-specific application instructions.

Lesser Cornstalk Borer

Apply **two** split applications of Lorsban 15G at 2 lb ai per acre (e.g., 15 oz per 1000 row feet on 36-inch rows), 10 to 30 days apart, to maintain lesser cornstalk borer (LCB) control when **any** of the following conditions exist:

- Extended periods of infestation by LCB are expected (particularly in drought conditions)
- Multiple generations of LCB are expected

To achieve maximum crop protection, make applications on a preventative basis prior to infestation. The first application should usually be made postplant just prior to or at pegging or as a rescue treatment when the insects first appear. The second application should generally be made 30 days following the first application and/or timed to coincide with the period when significant LCB populations are expected. Make the second application no sooner than 10 days after the first application. Do not make more than two applications of Lorsban 15G or apply more than 4 lb ai per acre per season. Refer to the product label for pest-specific application instructions.

®Trademark of Dow AgroSciences LLC

R061-011

Issued: 06/27/07

Initial printing.