

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

BIFENTHRIN | **GROUP 3A** | **INSECTICIDE**

Supplemental Labeling

This supplemental label expires on December 31, 2024 and must not be used or distributed after this date.



For At-Plant use on Sunflower

For mixing directly with liquid fertilizer to control listed soil insect pests.

EPA Reg. No. 279-3302

Active Ingredient:	By Wt.
Bifenthrin: *	17.15%
Other Ingredients:	82.85%
	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum
This product contains 1.5 lbs active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

This label must be in the possession of the user at the time of application.
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH ITS LABELING.
ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA
REGISTERED LABEL ARE TO BE FOLLOWED.**

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for CAPTURE LFR Insecticide before applying. Use of CAPTURE LFR Insecticide according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container of CAPTURE LFR Insecticide. Carefully follow all precautionary statements and application use directions.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

SUNFLOWER (Crop Subgroup 20B)

Calendula, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Safflower, Stokes Aster, Tallowwood, Tea Oil Plant, Vernonia, cultivars, varieties, and/or hybrids of these

At-Plant

PEST	USE RATES			DIRECTIONS
	fl oz/A	fl oz/1000 Linear ft.	lb ai/A	
Army cutworm Cutworm spp. Grape colaspis Root aphids Seedcorn maggot White grub spp. Wireworm	3.4 – 17.0	0.2 - 0.98	0.04 - 0.2	Apply as a 5 to 7-inch band (T-band) over an open furrow, or in-furrow with the seed. For Army cutworm or Cutworm species, apply as a 5 to 7-inch band over the row on the soil surface, a 5 to 7-inch band over the open furrow (T-band), in-furrow with the seed, or broadcast to the soil surface.
At-Plant Restrictions: <ul style="list-style-type: none"> • Do not apply more than 0.2 lb ai/A per year as an at-plant application. • Do not make more than 1 application per year. 				

FMC, Capture, LFR - Trademarks of FMC Corporation.
 © 2022 FMC Corporation. All rights reserved.

R-4654 020722 02-04-22