



RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

LAMBDA-CYHALOTHRIN GROUP **3A** INSECTICIDE

FIFRA Section 24(c) Special Local Need (SLN) Label
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Warrior II with Zeon Technology®
EPA Reg. No. 100-1295
EPA SLN No. WA-260014
For Control of Lygus Bug on Dill, Carrot, Parsley, and Parsnip Grown for Seed

This label for Warrior II with Zeon Technology expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA-registered label.

Dill, Carrot, Parsley, Parsnip Grown for Seed

Target Pest	Rate (fl oz/A) (lb ai/A)	Comments
Lygus Bug spp. ¹	1.28-1.92 (0.02-0.03)	<ul style="list-style-type: none">• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gallons per acre by air or 10 gallons per acre by ground. When foliage is dense and/or pest populations are high 5-10 gallons per acre by air or 20 gallons per acre by ground and higher use rates are recommended. Use higher rates for increased residual control, such as prior to crop blooming. If application is made during bloom, use the lower rate of application.• This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or broadleaf weeds. Do not apply the 1.92 oz/acre (0.03 lb ai/acre) rate of this product to blooming seed crops. Apply the 1.92 oz/acre (0.03 lb ai/acre) rate as a pre-bloom or post-bloom spray only. Applications of the 1.28 oz/acre (0.02 lb ai/acre) rate of this product to blooming seed crops must be timed to coincide with periods of minimum bee activity between late evening and midnight. Be aware of bee hazard resulting from a cool evening and/or morning dew. Do not apply directly to bee shelters/hives. It may be advisable to remove bee shelters/hives during and for 2-3 days following application.• If Warrior II with Zeon Technology is used as a prebloom spray it is not advisable to use during bloom to reduce potential for the development of insecticide resistance.

¹See resistance statement under General Directions for Use on the EPA-registered label.



Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Restrictions/Precautions

- The Restricted Entry Interval (REI) is 24 hours.
- Do not apply more than 0.12 lb ai (0.48 pints) per acre per season.
- Chemigation: For use in accordance with this SLN label do not apply this product through any type of irrigation system.

Crops Grown for Seed Restrictions/Precautions

1. All dill, carrot, parsley and parsnip seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
2. No portion of the dill, carrot, parsley, parsnip seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
3. Dill, carrot, parsley, parsnip seed shall bear a tag or container label, which forbids use of the seed for human consumption or animal feed.
4. Dill, carrot, parsley, parsnip seed may not be distributed for human consumption or animal feed.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling a cleaned container is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington

©2025 Syngenta

Warrior II with Zeon Technology® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: WA1295099DA0526