

---

**SUPPLEMENTAL LABEL**

---



**Syngenta Crop Protection, Inc.**  
P.O.Box 18300  
Greensboro, NC 27419-8300

**SUPPLEMENTAL LABEL FOR DURIVO® TO REVISE SOIL APPLICATION METHODS FOR BRASSICA (COLE) LEAFY VEGETABLES, CUCURBIT VEGETABLES, FRUITING VEGETABLES AND LEAFY VEGETABLES.**

Active Ingredient:	
Thiamethoxam <sup>1</sup> .....	17.5%
Chlorantraniliprole <sup>2</sup> .....	8.8%
Other Ingredients:	73.7%
Total:	100.0%

<sup>1</sup>CAS No. 153719-23-4  
<sup>2</sup>CAS No. 500008-45-7

Durivo is a soluble concentrate containing 1.67 lb. of thiamethoxam and 0.835 lb. chlorantraniliprole per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

EPA Reg. No. 100-1318

SCP 1318A-S1 0709

This label expires January 16, 2011.

**All applicable directions, restrictions and precautions on the EPA-registered label must be followed.**

**Before using Durivo, as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

## DIRECTIONS FOR USE

### APPLICATION PROCEDURES

#### SOIL APPLICATION

Select spray nozzles or metering orifices which will provide accurate and uniform deposition. Durivo can be applied during planting or transplanting as an in-furrow spray, transplant hole drench or narrow surface band above the seedline; after planting as a post-seeding, post-transplanting, hill drench, drip chemigation, root zone and shank application. These application methods allow the insecticide to be absorbed by plant roots. Although Durivo is rapidly taken up by plant roots and rapidly moves throughout the plant, the use of sufficient water volume and root zone placement will ensure that the chemical contacts the roots, resulting in optimal uptake and performance.

When making post-emergence applications, direct the application at the base of the plant for optimum root uptake. To help insure accuracy, calibrate sprayer before each use. For information on spray equipment and calibration, consult sprayer manufacturers and/or State Extension Service specialists.

**Do not apply Durivo as a broadcast foliar spray with ground or aerial equipment.**

### CROP USE DIRECTIONS

Crop	Pest	Rate Per Acre Per Application
<b>Brassica (Cole) Leafy Vegetables</b>		
<b>Head &amp; Stem Brassica</b> Broccoli Broccoli, Chinese Brussels sprouts Cabbage Cabbage, Chinese (napa) Cabbage, Chinese mustard Cauliflower Cavalo broccolo Kohlrabi  <b>Leafy Brassica Greens</b> Broccoli raab Cabbage, Chinese (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens	Aphids Beet Armyworm Cabbage Looper Cabbage Webworm Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Southern Cabbageworm Thrips Whiteflies Yellowstriped Armyworm	10.0 – 13.0 fl. oz./A

#### Use Restrictions

- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per growing season.
- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet
- **Application Method:** Apply specified dosage in sufficient water volume to ensure uniform application and incorporation into the soil using one of the following methods:
  1. In-furrow spray at the seeding or transplant depth, or a narrow surface band above the seedline during planting. For surface-banded applications, incorporate to the seeding depth with sufficient sprinkle or drip irrigation within 24 hours.

2. Post seeding, transplant, or hill drench using sufficient water volume to ensure incorporation into the root zone.
3. In drip (trickle) chemigation.
4. Shanked into the root zone after establishment or transplanting using fertilizer knives or other similar equipment. After application, incorporate with enough irrigation to move the chemical to the root zone.

For planting systems where multiple rows are planted on beds, apply Durivo according to one of the above methods.

- **Pre-Harvest Interval (PHI):** 30 Days

**Refer to Resistance Management section.**

Crop	Pest	Rate Per Acre Per Application
<p><b>Cucurbit Vegetables</b>                      Chayote                      Chinese waxgourd                      Citron melon                      Cucumber                      Gherkin                      Gourd, edible (hyotan, cucuzza, hechima, Chinese okra)  <i>Momordica</i> species (balsam apple, balsam pear, bittermelon, Chinese cucumber)                      Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i>) includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon)                      Pumpkin                      Squash: summer (crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini) and winter (butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash)                      Watermelon (includes hybrids and/or varieties of <i>Citrullus lanatus</i>)</p>	<p>Aphids                      Cucumber Beetles<sup>1</sup>                      Flea Beetles                      Leafhoppers                      Leafminers<sup>1</sup>                      Melonworm                      Pickleworm                      Thrips                      Whiteflies</p>	<p>10.0 – 13.0 fl. oz./A</p>

**Use Restrictions**

- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per growing season.
- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Method:** Apply specified dosage in sufficient water volume to insure uniform application and incorporation into the soil using one of the following methods:
  1. In-furrow spray at the seeding or transplant depth, or a narrow surface band above the seedline during planting. For surface-banded applications, incorporate to the seeding depth with sufficient sprinkle or drip irrigation within 24 hours.

2. Post seeding, transplant, or hill drench using sufficient water volume to ensure incorporation into the root zone.
3. In drip (trickle) chemigation.
4. Shanked into the root zone after establishment or transplanting using fertilizer knives or other similar equipment. After application, incorporate with enough irrigation to move the chemical to the root zone.

- **Pre-Harvest Interval (PHI):** 30 Days
- **Pest Control:** <sup>1</sup>Suppression.

**Refer to Resistance Management section.**

Crop	Pest	Rate Per Acre Per Application
<b>Fruiting Vegetables</b> Tomato Peppers: bell, chili, cooking, pimento, and sweet Eggplant Groundcherry Pepino Tomatillo	Aphids Beet Armyworm Colorado Potato Beetle Fall Armyworm Flea Beetles Hornworm species Leafhoppers Leafminers <sup>1</sup> Loopers Potato Psyllid Southern Armyworm Thrips Tomato Fruitworm Tomato Pinworm Whiteflies Yellowstriped Armyworm	10.0 – 13.0 fl. oz./A

**Use Restrictions**

- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. ai of chlorantraniliprole containing products per acre per growing season.
- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Method:** Apply specified dosage in sufficient water volume to insure uniform application and incorporation into the soil using one of the following methods:
  1. In-furrow spray at the seeding or transplant depth, or a narrow surface band above the seedline during planting. For surface-banded applications, incorporate to the seeding depth with sufficient sprinkle or drip irrigation within 24 hours.
  2. Post seeding, transplant, or hill drench using sufficient water volume to ensure incorporation into the root zone.
  3. In drip (trickle) chemigation.
  4. Shanked into the root zone after establishment or transplanting using fertilizer knives or other similar equipment. After application, incorporate with enough irrigation to move the chemical to the root zone.
- **Pre-Harvest Interval (PHI):** 30 Days
- **Pest Control:** <sup>1</sup>Suppression

**Refer to Resistance Management section.**

Crop	Pest	Rate Per Acre Per Application
<b>Leafy Vegetables</b> Amaranth Arugula Cardoon Celery Celery, Chinese Celtuce Chervil Chrysanthemum: edible- leaved & garland Corn Salad Cress: garden and upland (yellow rocket, winter cress) Dandelion Dock (sorel) Endive (escarole) Fennel, Florence (finochio) Lettuce: head & leaf Orach Parsley Purslane: garden & winter Radicchio (red chicory) Rhubarb Spinach: New Zealand and Vine (Malabar, Indian) Swiss chard	Aphids Beet Armyworm Cabbage Looper Corn Earworm Diamondback Moth Fall Armyworm Flea Beetles Imported Cabbageworm Leafhoppers Leafminers <sup>1</sup> Whiteflies	10.0 – 13.0 fl. oz./A

**Use Restrictions**

- **Maximum Durivo Allowed per Growing Season:** Do not exceed a total of 13.0 fl. oz./Acre (0.257 lb. a.i./A) of Durivo or 0.172 lb. a.i. of thiamethoxam containing products or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per growing season.
- **Application Rate:** Use lower rates for short residual control and higher rates for long residual control. See rate conversion chart for rate per 1000 linear feet.
- **Application Method:** Apply specified dosage in sufficient water volume to ensure uniform application and incorporation into the soil using one of the following methods:
  1. In-furrow spray at the seeding or transplant depth, or a narrow surface band above the seedline during planting. For surface-banded applications, incorporate to the seeding depth with sufficient sprinkle or drip irrigation within 24 hours.
  2. Post seeding, transplant, or hill drench using sufficient water volume to ensure incorporation into the root zone.
  3. In drip (trickle) chemigation.
  4. Shanked into the root zone after establishment or transplanting using fertilizer knives or other similar equipment. After application, incorporate with enough irrigation to move the chemical to the root zone.

For planting systems where multiple rows are planted on beds, apply Durivo according to one of the above methods.

- **Pre-Harvest Interval (PHI):** 30 Days
- **Pest Control:** <sup>1</sup>Suppression

**Refer to Resistance Management section.**

Durivo® and the Syngenta logo are trademarks of a Syngenta Group Company  
 © 2009 Syngenta  
 Syngenta Crop Protection, Inc.  
 P.O. Box 18300  
 Greensboro, North Carolina 27419-8300

Durivo 1318A-S1 0709-Ig7-17-09