

FIFRA SECTION 24(C) SPECIAL LOCAL NEED LABEL  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

**CURIO® HERBICIDE**

EPA REG. NO. 71368-82

EPA SLN NO. WA-100004

This label for Curio® Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027.

ACTIVE INGREDIENT	(by weight):
Chlorimuron Ethyl	
Ethyl 2-[[[(4-chloro-6-methoxypyrimidin-2-yl)amino]carbonyl]amino]sulfonyl]benzoate .....	25.0%
OTHER INGREDIENTS .....	75.0%
TOTAL .....	100.0%

**For post-emergence control / suppression of creeping buttercup  
in cranberries (bearing and nonbearing)**

**DIRECTIONS FOR USE**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the main EPA-registered label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label.
- Read and understand the Pacific Coast Cranberry Research Foundation Special Conditions and Disclaimer for use before using Curio® Herbicide on cranberries.

**Environmental Hazards**

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Do not apply where or when conditions favor runoff.

**Groundwater Advisory:** Chlorimuron-ethyl is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorimuron-ethyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

**Non-Target Organism Advisory Statement:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of the main EPA-registered label.

**Windblown Soil Particles:** Curio® Herbicide has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying Curio® Herbicide if prevailing local conditions may be expected to result in off-site movement.

## APPLICATION INSTRUCTIONS

### Crop:

Cranberries (bearing and nonbearing)

### Application Rate:

0.5 to 1.0 oz Curio® Herbicide/acre (0.008 to 0.016 lbs chlorimuron ethyl/acre)

### Postemergence:

- Apply Curio® Herbicide as a broadcast spray using 20 to 40 gallons of water per acre with ground spray equipment only.
- Spot treatment can be applied only through a calibrated boom to prevent misapplication.
- Apply Curio® Herbicide any time after March 1 but no later than 60 days before harvest. Optimum control of creeping buttercup can be achieved from an application made in March or April when the weed is most actively growing.
- Curio® Herbicide is a water dispersible granule that does not include a spray adjuvant. Applications must include one of the following adjuvants: a nonionic surfactant at the rate of 2 pints per 100 gallons of spray solution (0.25% v/v), a crop oil concentrate at the rate of 8 pints per 100 gallons of spray solution (1.0% v/v), or a modified vegetable oil concentrate at the rate of 8 pints per 100 gallons of spray solution (1.0% v/v).

### Restrictions:

- **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- For use under this SLN **DO NOT** apply this product through any type of irrigation system. **DO NOT** apply by air.
- **DO NOT** make more than one application per year.
- **DO NOT** apply more than 1.0 oz Curio® Herbicide/acre/year (0.016 lbs chlorimuron ethyl/acre/year).
- **DO NOT** apply within 60 days of harvest.
- **DO NOT** apply directly to water, to areas where surface water is present, or within 5 feet of any water moving off or through the cranberry field.

### Precautions:

- Crop injury may result if crop is under stress conditions caused by cold weather, drought, low fertility, disease or insect damage.
- Avoid application if rainfall is expected within one hour.

### WSDA Container Disposal Guidance:

Completely empty the contents of the pesticide container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington. Information regarding the disposal of unwanted or waste pesticides is available on the WSDA Waste Pesticide Program website at <https://agr.wa.gov/wastepesticide>.

### Pacific Coast Cranberry Research Foundation

#### Special Conditions and Disclaimer for Use of Curio® Herbicide on cranberries:

The Pacific Coast Cranberry Research Foundation (PCCRF) intends that this Section 24(c) label be distributed only by the PCCRF and distributed only to growers who agree in writing to the terms and conditions required by the PCCRF and Nufarm, including a waiver and release from all liability and indemnification by the user and/or grower of the PCCRF, Nufarm and others for failure to perform and crop damage from the use of Curio® Herbicide on cranberries. If such terms and conditions are unacceptable, return Curio® Herbicide at once unopened.

This product when used on cranberries may lead to crop injury, loss or damage. The PCCRF recommends that the user and/or grower test this product in order to determine its suitability for such intended use. The PCCRF and Nufarm make this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use Curio® Herbicide must be made by each individual Curio® Herbicide user and/or grower on the basis of possible crop injury from Curio® Herbicide, the severity of weed infestation, the cost of alternative (pest) controls, and other factors. The PCCRF intends that because of the risk of failure to perform or crop damage that all such use is at the user's and/or grower's risk, to the extent consistent with applicable law.

This Special Conditions and Disclaimer statement is required by the PCCRF and Nufarm, and is not required or enforced by the USEPA or the WSDA.

24(c) Registrant:  
Pacific Coast Cranberry Research Foundation  
2500 Sandridge Road  
Ilwaco, WA 98624

Manufactured by:  
Nufarm Inc.  
4020 Aerial Center Parkway  
Morrisville, NC 27560

RV071822