

# SUPPLEMENTAL LABEL

## FOR USE ON PEACH AND NECTARINE



**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

### DIRECTIONS FOR USE

CROP	PEST	RATE	COMMENTS
Peach Nectarine	Aphids* Sharpshooters Leafhoppers	<b>FOLIAR:</b> 3.5 to 7 fl oz/A (0.090 to 0.180 lb ai/A)	<b>Higher water volumes provide improved insect control.</b>  Begin applications when first pest activity is noticed or when insects reach threshold levels per State and County Extension Service recommendations. Repeat as needed to maintain control, but not more often than every 7 days. For best results, time application before a damaging population becomes established.
	Peach tree borer* Plum curculio* Aphids* Stinkbugs	<b>FOLIAR:</b> 5.25 to 7 fl oz/A (0.135 to 0.180 lb ai/A)	
	Aphids* Sharpshooters Leafhoppers Peach tree borer*	<b>SOIL:</b> 10.5 fl oz/A (0.270 lb ai/A)	Under severe pest pressure, use the higher specified rates.  The rate applied affects the length of control. Use the high rate where infestations occur later in crop development, or where pest pressure is continuous.  SCORPION 35SL INSECTICIDE may be mixed and/or alternated with commonly used insecticides to comply with local IPM and resistance management programs.

**NOTE:** Regardless of application method do not apply more than a total of 14.25 fl oz/A of SCORPION (0.360 lb. ai) per acre per season.

#### Foliar Application

- Apply with air or ground equipment in adequate water for uniform coverage (A minimum of 5 gals/A by air or 50 gals/A by ground).
- Do not apply SCORPION within three (3) days of harvest.
- Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.270 lb ai) per acre per season.
- Interval between applications cannot be less than 7 days.

#### Soil Application

- Do not apply SCORPION within twenty one (21) days of harvest
- Apply with ground equipment in adequate water for uniform coverage (A minimum of 100 gals/A).
- Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.270 lb ai) per acre per year.

Apply specified dosage in sufficient carrier volume to insure uniform application and distribution within and around the root zone of each tree using one of the following methods:

1. As a drench. Applications should be made with sufficient water to insure incorporation into the root zone.
2. Using drip, trickle, micro sprinkler or any customized irrigation system derived from those systems to water trees independently.

\* Provides suppression only

**Gowan**  
The Go To Company

EPA Reg. No. 10163-317  
EPA Est. No.

Distributed By:  
Gowan Company  
PO BOX 5569  
Yuma, AZ 85366