Product Bulletin

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Transform® WG

EPA Reg. No. 62719-625

Special 2(ee) Recommendation[†]

(For Distribution and Use in the States of Idaho, Michigan, Oregon, Pennsylvania, Utah and Washington)

For Control of Mealybugs in Pome (Crop Group 11) and Stone Fruits (Crop Group 12)

Refer to product label for Transform® WG for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

- [†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.
- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Transform WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Transform WG according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Transform WG insecticide.

Directions for Use

Pests	Transform WG (oz/acre)
mealybugs	2.25 – 2.75
	(0.071 – 0.086 lb ai/acre)

Application Timing: Consult your Corteva Agriscience representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Restrictions:

- Preharvest Interval: Do not apply within 7 day of harvest.
- Minimum Treatment Interval: Do not make applications less than 7 days apart.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 8.5 oz of Transform WG (0.266 lb ai of sulfoxaflor) per acre per year.
- Do not apply this product at any time between 3 days prior to bloom and until after petal fall.

R396-282 Issued: 03/22/22 Initial Printing

^{™®} Trademarks of Corteva Agriscience and its affiliated companies