

GROUP 14 HERBICIDE

Lock Down

Herbicide

Directions for use to maintain bare ground non-crop areas

ACTIVE INGREDIENT:

*Flumioxazin 51.0%

OTHER INGREDIENTS: 49.0%

TOTAL: 100.0%

*(2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione)

Lock Down Herbicide is a water dispersible granule containing 51% active ingredient.

**KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

EPA Reg. No. 71368-103

Manufactured for
Nufarm, Inc.
150 Harvester Drive
Burr Ridge, IL 60527



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUTION**

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust and spray mist. Avoid contact with skin, eyes or clothing.

FIRST AID	
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes and socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to non-target plants and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to non-target plants and aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is toxic to plants and should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions this product may have a potential to run off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands or on the downhill side of fields where run-off could occur will minimize water run-off and is recommended.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Use strictly in accordance with precautionary statements and directions, and with applicable State and Federal regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

Do not enter or allow others to enter the treated areas until sprays have dried.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

SPECIMEN

TABLE OF CONTENTS

PRODUCT INFORMATION	
Use Precautions	5
Resistance Management.....	5
Preemergence Application	5
Postemergence Application	5
APPLICATION EQUIPMENT	5
Sprayer Preparation	5
Mixing Instructions	6
Sprayer Cleanup.....	6
Spray Drift Reduction.....	6
WEEDS CONTROLLED	
TABLE 1. WEEDS CONTROLLED BY LOCK DOWN HERBICIDE	7
DIRECTIONS FOR USE TO MAINTAIN BARE GROUND NON-CROP AREAS	
Preemergence Application	8
Postemergence Application	8
Soil Characteristics	8
Carrier Volume and Spray Pressure	8
Preemergence Application	8
Postemergence Application	8
Additives	
Postemergence Application	8
Jar Test to Determine Compatibility of Adjuvants and Lock Down Herbicide	8
Application Equipment	9
Broadcast Application.....	9
Band Application.....	9
Handgun Application.....	9
Aerial Application	9
Tank Mix Application	9
Tank Mix Combinations for Non-Selective Vegetation Control.....	9
Restrictions and Limitations.....	9
STORAGE AND DISPOSAL	10

PRODUCT INFORMATION

This product is a selective herbicide to maintain bare ground non-crop areas when used in accordance with this label. This product is effective as a preemergence and/or postemergence herbicide for control of selected grass and broadleaf weeds.

This product controls weeds by inhibiting protoporphyrinogen oxidase, an essential enzyme required by plants for chlorophyll biosynthesis. Seedling weeds are controlled preemergence when exposed to sunlight following contact with the soil applied herbicide.

USE PRECAUTIONS

- Do not apply when weather conditions favor spray drift from treated areas.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply more than 12 oz of this product per acre per application.
- Do not apply more than 24 oz of this product per acre per year.
- Do not apply to moist or wet desirable plant foliage.
- Do not apply within 300 feet of non-dormant pome or stone fruit crops.
- Treatment of powdery, dry soil or light sandy soil, or light sandy soil when there is little to no likelihood of rainfall soon after may result in off target movement and possible damage to actively growing susceptible crops when soil particles are moved by wind or water. Do not apply when these soil and environmental conditions are present.

RESISTANCE MANAGEMENT

Any weed population may contain or develop plants naturally resistant to herbicides in various mode of action classes. Resistant biotypes may eventually dominate the weed population if the same class of chemistry/mode of action herbicides are used repeatedly in the same field or in successive years. These resistant biotypes may not be adequately controlled by herbicides in a mode of action class for which resistance has developed. A gradual or total loss of weed control may occur over time. Other resistance mechanisms that are not linked to site of action, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To Delay Herbicide Resistance

- Avoid the use of herbicides that have a similar target site mode of action in consecutive years.
- Herbicide use should be based on an Integrated Pest Management (IPM) program that includes scouting, record keeping, and consideration of cultivation practices, water management, weed free crop seed, crop rotation, and other chemical or cultural control practices.
- Monitor treated weed population for resistance development and report suspected resistance.
- Contact your local extension or crop expert (advisor) for any additional pesticide resistance management and/or IPM recommendations for specific crops and weed biotypes.
- For further information contact Nufarm Inc. at the following toll free number 800-345-3330.

PREEMERGENCE APPLICATION

Preemergence application of this product should be made prior to weed emergence. Moisture is necessary to activate this product for residual weed control. Moisture is needed to move this product into the soil for preemergence weed control. Dry weather following application of this product may reduce effectiveness.

POSTEMERGENCE APPLICATION

For best results, this product should be applied to actively growing weeds. Applying this product under conditions that do not promote active weed growth will reduce herbicide effectiveness.

Do not apply this product when the weeds are under stress due to drought, excessive water and extremes in temperatures or disease. This product is most effective when applied under sunny conditions at temperatures above 65° F.

This product is rainfast one hour after application. Applications should not be made if rain is expected within one hour of application or efficacy may be reduced.

APPLICATION EQUIPMENT

Important: Spray equipment, including all tanks, hoses, booms, screens and nozzles, should be thoroughly cleaned. **Spray equipment used to apply this product should not be used to apply other materials to any desirable plant foliage.** Equipment with this product's residue remaining in the system may result in crop injury to subsequently treated crops.

SPRAYER PREPARATION

Before applying this product, start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms should be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to the sulfonylurea and phenoxy herbicides, are active at very small amounts and can cause crop injury when applied to susceptible crops. The spray equipment should be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply this product. If two or more products were tank mixed prior to this product's application, the most restrictive cleanup procedure should be followed.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. To ensure a uniform spray mixture, pre-slurry the required amount of this product with water prior to addition to the spray tank. Use a minimum of 1 gal of water per 10 oz of this product.
3. While agitating, slowly add the pre-slurried this product to the spray tank. Agitation should create a rippling or rolling action on the water surface.
4. If tank mixing this product with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
5. Add any required adjuvants.
6. Fill spray tank to desired level with water. **Agitation should continue until spray solution has been applied.**
7. Mix only the amount of spray solution that can be applied the day of mixing. This product should be applied within 24 hours of mixing.

SPRAYER CLEANUP

Except for dedicated bare ground herbicide application equipment, spray equipment should be cleaned each day following this product application. The following steps should be used to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank, add suitable commercial spray tank cleaning material, following label directions, or add 1 gal of 3% household ammonia for every 100 gals of water, circulate through sprayer for 5 minutes, and then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes.
4. Drain tank completely.
5. Add enough clean water to the spray tank to allow all hoses, booms, screens and nozzles to be flushed for 2 minutes.
6. Remove all nozzles and screens and rinse them with clean water.

SPRAY DRIFT REDUCTION

Do not apply under circumstances where possible drift to unprotected persons or to food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur.

- Use the largest droplet size consistent with acceptable efficacy. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. For ground boom and aerial applications, use medium or coarser spray nozzles according to ASAE 572 definition for standard nozzles or a volume mean diameter (VMD) of 300 microns or greater for spinning atomizer nozzles.
- Make aerial or ground applications when the wind velocity favors on-target product deposition. Drift potential is lowest between wind speeds of 2-10 mph. For all non-aerial applications, wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.
- Do not make aerial or ground applications into areas of temperature inversions. Inversions are characterized by stable air and increasing temperatures with increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. Where permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For ground boom applications, apply with nozzle height at the lowest boom height which provides uniform coverage and reduces exposure to evaporation and wind.

WEEDS CONTROLLED

When this product is applied preemergence or postemergence at recommended rates and weed stages, the following grasses and broadleaf weeds are controlled:

TABLE 1. WEEDS CONTROLLED BY LOCK DOWN HERBICIDE

COMMON NAME	SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME
Alyssum, Hoary	<i>Berteroa incana</i>	Moss	<i>Bryum</i> spp.
Amaranth		Mustard	
Palmer	<i>Amaranthus palmeri</i>	Tumble	<i>Sisymbrium altissimum</i>
Spiny	<i>Amaranthus spinosus</i>	Wild	<i>Brassica kaber</i>
Barnyardgrass*	<i>Echinochloa crus-galli</i>	Nightshade	
Beggarweed, Florida	<i>Desmodium Tortuosum</i>	Black	<i>Solanum nigrum</i>
Bittercress, Hairy	<i>Cardamine hirsute</i>	Eastern Black	<i>Solanum ptycanthum</i>
Bluegrass, Annual*	<i>Poa annua</i>	Hairy	<i>Solanum sarrachoides</i>
Burclover, California	<i>Medicago Polymorpha</i>	Panicum	
Carpetweed	<i>Mollugo verticillata</i>	Fall*	<i>Panicum dichotomiflorum</i>
Chickweed		Texas*	<i>Panicum texanum</i>
Common	<i>Stellaria media</i>	Parsley-Peirt	<i>Alchemilla arvensis</i>
Mouseear	<i>Cerastium vulgatum</i>	Pearlwork, Birdseye*	<i>Sagina procumbens</i>
Crabgrass		Pennycress, Field	<i>Thlaspi arvense</i>
Large*	<i>Digitaria sanguinalis</i>	Phyllanthus, Longstalked	<i>Phyllanthus tenellus</i>
Smooth*	<i>Digitaria ishaemum</i>	Pigweed	
Southern*	<i>Digitaria ciliaris</i>	Prostrate	<i>Amaranthus blitoides</i>
Croton, Tropic	<i>Croton glandulosus</i> var. <i>septentrionalis</i>	Redroot	<i>Amaranthus retroflexus</i>
	<i>Taraxacum officinale</i>	Smooth	<i>Amaranthus hybridus</i>
Dandelion*	<i>Eupatorium capillifolium</i>	Tumble	<i>Amaranthus albus</i>
Donfennel	<i>Eclipta prostrate</i>	Pineapple-weed*	<i>Matricaria matricarioides</i>
Eclipta		Plantain	
Foxtail		Broadleaf*	<i>Plantago major</i>
Bristly*	<i>Setaria verticillata</i>	Buckhorn*	<i>Plantago lanceolata</i>
Giant*	<i>Setaria faberi</i>	Poinsettia, Wild	<i>Euphorbia heterophylla</i>
Green*	<i>Setaria viridis</i>	Pondweed, Sago	<i>Potamogeton pectinatus</i>
Yellow*	<i>Setaria glauca</i>	Puncturevine	<i>Tribulus terrestris</i>
Galinsoga, Hairy	<i>Galinsoga ciliata</i>	Purslane, Common	<i>Portulaca oleracea</i>
Geranium, Carolina	<i>Geranium carolinianum</i>	Pusley, Florida	<i>Richardia scabra</i>
Goosegrass*	<i>Eleusine indica</i>	Ragweed	
Groundsel, Common	<i>Senecio vulgaris</i>	Common	<i>Ambrosia artemisiifolia</i>
Henbit	<i>Lamium amplexicaule</i>	Giant	<i>Ambrosia trifida</i>
Horseweed*	<i>Conyza Canadensis</i>	Redmaids	<i>Calandrinia ciliata</i>
Indigo, Hairy	<i>Indigofera hirsuta</i>	Redweed	<i>Melochia corchorifolia</i>
Ivy, Ground*	<i>Glechoma hederacea</i>	Rocket, Yellow	<i>Barbarea vulgaris</i>
Jimsonweed	<i>Datura stramonium</i>	Senna, Coffee	<i>Cassia occidentalis</i>
Kochia	<i>Kochia scoparia</i>	Sesbania, Hemp	<i>Sesbania exaltata</i>
Kyllinga, Green*	<i>Kyllinga brevifolia</i>	Shepherd's-Purse	<i>Capsella bursa-pastoris</i>
Ladysthumb	<i>Polygonum persicaria</i>	Sida, Prickly (Teaweed)	<i>Sida spinosa</i>
Lambsquarters, Common	<i>Chenopodium album</i>	Signalgrass*	<i>Brachiaria platyphylla</i>
Lovegrass, California*	<i>Eragrostis diffusa</i>	Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>
Mallow		Sowthistle, Annual	<i>Sonchus oleraceus</i>
Common	<i>Malva neglecta</i>	Spurge	
Little	<i>Malva parviflora</i>	Prostrate	<i>Euphorbia humistrata</i> Engelm
Venice	<i>Hibiscus trionum</i>	Spotted	<i>Euphorbia maculata</i>
Mayweed*	<i>Anthemis cotula</i>	Starbur, Bristly*	<i>Acanthospermum hispidum</i>
Morningglory		Thistle	
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integruscula</i>	Canada*	<i>Cirsium arvense</i>
Ivyleaf	<i>Ipomoea hederacea</i>	Russian	<i>Salsola iberica</i>
Red/Scarlet	<i>Ipomoea coccinea</i>	Velvetleaf	<i>Abutilon theophrasti</i>
Smallflower	<i>Jacquemontia tamnifolia</i>	Waterhemp	
Tall	<i>Ipomoea purpurea</i>	Common	<i>Amaranthus rudis</i>
		Tall	<i>Amaranthus tuberculatus</i>
		Woodsorrel, Yellow*	<i>Oxalis stricta</i>

*Preemergence control only

DIRECTIONS FOR USE

TO MAINTAIN BARE GROUND NON-CROP AREAS

This product, when used as directed, can be used for non-selective vegetation control to maintain bare ground non-crop areas that must be kept weed-free. Apply this product only to:

- Bare ground under guard rails, above-ground pipelines, and railroad beds, railroad yards and surrounding areas
- Bare ground in parking and storage areas, plant sites, substations, pumping stations, and tank farms
- Bare ground areas of airports, brick yards, industrial plant sites, lumber yards, military installations, and storage areas
- Bare ground around farm buildings, and along ungrazed fence rows, wind breaks and shelter belts
- Road surfaces, improved roadside areas and gravel shoulders.

Follow all applicable directions as outlined above under General Information. See Table 1 for a list of broadleaf weeds and grasses controlled by this product.

This product offers residual and postemergence control of susceptible broadleaf and grass weeds as well as additional mode of action to assist in the control of ALS (acetolactate synthase) resistant weeds. The length of residual control is dependent on the rate applied as well as on rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase.

PREEMERGENCE APPLICATION

Apply 8 to 12 oz (0.25 to 0.38 lb ai/A) of this product per broadcast acre as a preemergence application. Preemergence (to weed emergence) applications of this product should be made to a weed free soil surface. Preemergence applications of this product must be completed prior to weed emergence. Moisture is necessary to activate this product on soil for residual weed control. Dry weather following application of this product may reduce effectiveness. However, when adequate moisture is received after dry conditions, this product will control susceptible germinating weeds.

POSTEMERGENCE APPLICATION

Apply 8 to 12 oz (0.25 to 0.38 lb ai/A) of this product per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 qt/A crop oil concentrate). The addition of an adjuvant enhances this product activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of this product. Emerged weeds are controlled postemergence with this product, however, translocation of this product within a weed is limited, and control is affected by spray coverage and by the addition of an adjuvant. The most effective postemergence weed control with this product occurs when applied in combination with a surfactant to weeds less than 2 inches in height.

SOIL CHARACTERISTICS

Application of this product to soils with high organic matter and/or high clay content may require higher dosages than with soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

CARRIER VOLUME AND SPRAY PRESSURE

PREEMERGENCE APPLICATION

To ensure uniform coverage, use 10 to 30 gals of spray solution per acre. Nozzle selection should meet manufacturer's gallonage and pressure recommendation for preemergence herbicide application.

POSTEMERGENCE APPLICATION

To ensure thorough coverage, use 15 to 30 gals of spray solution per acre. Use 20 to 30 gals per acre if dense vegetation or heavy residue is present on the soil surface. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence herbicide application.

ADDITIVES

POSTEMERGENCE APPLICATION

When applying this product after weed emergence, mix with an agronomically approved adjuvant. A crop oil concentrate which contains at least 15% emulsifiers and 80% oil or a non-ionic surfactant containing at least 80% active ingredient should be used when applying this product as part of a postemergence weed control program. Mixing compatibility should be verified by a jar test before using.

A spray grade nitrogen source (either ammonium sulfate at 2.0 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with a crop oil concentrate or non-ionic surfactant to enhance weed control. The addition of a nitrogen source does not replace the need for crop oil concentrate or non-ionic surfactant.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND LOCK DOWN HERBICIDE

A jar test should be performed before mixing commercial quantities of this product, when using this product for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pt of water to a quart jar. The water should be from the same source and have the same temperature as the water used in the spray tank mixing operation.
2. Add 3 grams (approximately 1 level tsp) of this product for the 8 oz/A rate or 4 grams (approximately 1 1/2 tsp) for 12 oz/A rate to the jar. Gently mix until product disperses.
3. Add 60 ml (4 Tbsp or 2 fl oz) of additive to the quart jar and gently mix.
4. If nitrogen is being used, add 16 ml (1 Tbsp) of the 28 to 32% nitrogen source to the quart jar. If ammonium sulfate is being used, add 19 grams of AMS to the quart jar in place of the 28 to 32% nitrogen.

5. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
6. An ideal tank mix combination will be uniform and free of suspended particles. If any of the following conditions are observed the choice of adjuvant should be questioned:
 - a) Layer of oil or globules on the solution surface.
 - b) Flocculation: Fine particles in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: Thickening texture (coagulated) like gelatin.

APPLICATION EQUIPMENT

Application equipment should be clean and in good repair. Nozzles should be uniformly spaced on boom and frequently checked for accuracy.

BROADCAST APPLICATION

Apply this product, and this product tank mixes, with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume.

BAND APPLICATION

When banding, use proportionately less water and this product per acre.

HANDGUN APPLICATION

Applications may also be made using a handgun sprayer. Use a spray volume of at least 40 gals per acre to insure uniform coverage.

AERIAL APPLICATION

- **Aerial applications are limited to maintaining weed free railroad beds, railroad yards and surrounding areas and military installations.**

To obtain satisfactory weed control with aerial applications of this product, uniform coverage must be obtained. Do not spray when drift is possible or when wind velocity is more than 10 mph. Avoid spraying this product within 200 feet of dwellings, adjacent sensitive crops or environmentally sensitive areas. To obtain satisfactory application and drift, the following directions must be observed:

Volume Pressure

Use this product in 5 to 10 gals of water per acre with a maximum spray pressure of 40 PSI. Application at less than 5 gals per acre will provide inadequate weed control. Higher gallonage applications provide more consistent weed control.

Nozzle and Nozzle Operation

Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles, such as diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzle must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.

Adjuvants

Refer to the additive section or the tank mix partner's label for adjuvant recommendation.

TANK MIX APPLICATION

In addition to weeds controlled by this product used alone, tank mixtures with other preemergence and postemergence herbicides registered for use in non-crop areas provide a broader spectrum of weed control. This product must be tank mixed with other non-crop herbicides including, but not limited to those products listed below.

TANK MIX COMBINATIONS FOR NON-SELECTIVE VEGETATION CONTROL

2,4-D	hexazinon	picloram
bromacil	imazapic	pramitol
chlorsulfuron	imazapyr	prodiamine
dicamba	metsulfuron-methyl	simazine
diuron	norflurazon	sulfometuron-methyl
chlorpyralid	oryzalin	tebuthiuron
glyphosate	pendimethalin	triclopyr

IMPORTANT: Completely read and follow the label of any potential this product tank mix partner. When using tank mixtures, use conditions must be in accordance with the most restrictive of the label limitations and precautions on either herbicide label.

RESTRICTIONS AND LIMITATIONS

- Do not apply more than 2 applications at 12 oz/A or 3 applications at 8 oz/A per year.
- Do not re-apply this product within 30 days.

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Keep pesticide in original container.

Store in a cool, dry, secure place.

Do not put formulation or dilute spray solution into food or drink containers.

Do not store or transport near feed or food.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night **(800) 892-0099**. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

[Container Disposal language for container sizes less than 5 gals]

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[Container Disposal language for container sizes greater than 5 gals or 50 lbs]

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water.

Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal.

Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV013013)