

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®]-4E

EPA Reg. No. 62719-220

EPA 24(c) Special Local Need Registration SLN DE-930004
For Distribution and Use Only in the State of Delaware

To Control Armyworms and Corn Earworm Larvae Infesting Sweet Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®]-4E insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban-4E according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban-4E .

Directions for Use

Apply Lorsban-4E at the rate of 1 pint per acre as a postemergence broadcast spray in sufficient waer to ensure thorough coverage of plants using either aerial (fixed-wing or helicopter) or power-operated ground equipment. Use no less than 15 gallons of water per acre for ground spray equipment or at least 3 gallons of water per acre for aerial application. Control may be reduced at low spray volumes under high temperature and wind conditions. Treat when field counts indicate that worms are or may become a problem.

Chemigation: Lorsban-4E may be broadcast applied postemergence through sprinkler irrigation systems at recommended application rates. See Chemigation (Sprinkler Irrigation) section on product label for application instructions.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 7 days before harvest.
- Do not allow meat or dairy animals to graze in treated areas.
- Do not feed treated corn silage, forage, or fodder to meat or dairy animals.
- Do not make more than 3 applications of Lorsban-4E per season.
- Do not make a second application of Lorsban-4E within 10 days of the first application.
- Maximum single application rate is 0.5 lb ai chlorpyrifos per acre.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

[®]Trademark of Dow AgroSciences LLC

R062-014

Approved: 04/22/2005

Replaces 122-58-032N.

Revisions: Added Restricted Use Pesticide; revised paragraph under Directions for Use; revised Chemigation paragraph; added "Specific Use" to "Restrictions"; revised PHI point; changed "livestock" to "meat or dairy animals"; added four points.