

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## RESTRICTED USE PESTICIDE

Due to high acute inhalation toxicity and carcinogenicity.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## Telone<sup>®</sup> II

EPA Reg. No. 62719-32

**For Distribution and Use Only in the States of Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, South Carolina and Texas**

### Nematode Suppression in Cotton

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Telone<sup>®</sup> II soil fumigant before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Telone II according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Telone II.

#### Directions for Use

Apply Telone II as a row treatment at the rate of 28 to 56 fl oz per 1000 feet of row (3 to 6 gallons per acre based on a 38-inch row spacing). Use suitable ground application equipment that insures placement of the fumigant at least 14 inches below the final planting surface. Use one chisel per row to treat only the soil where the crop is to be planted. Soil should be sealed immediately after application. This may be accomplished using bedding equipment. After application and sealing, leave the soil undisturbed for at least a week before planting.

#### Precaution:

- Do not plant if the odor of the fumigant is still present at the depth of injection.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R018-016

Approved: 12/21/91

Replaces 122-70-001N.