

CLOPYRALID | GROUP 4 | HERBICIDE

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

For Distribution and Use Only in the State of Oregon

Stinger[®]

EPA Reg. No. 62719-73
EPA SLN No. OR-230007

This label is valid until December 31, 2027, or until otherwise amended, withdrawn, cancelled, or suspended.

Postemergence Broadleaf Weed Control in Radish Grown for Seed, Including Daikon Grown for Seed

Active Ingredient:

cloprialid: 3,6-dichloro-2-pyridinecarboxylic acid, monoethanolamine salt.....	40.9%
Other Ingredients	59.1%
Total	100.0%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid - 31% (3 lb/gal)

Keep Out of Reach of Children

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

Groundwater Advisory:

Clopyralid is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory:

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of clopyralid and its degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Non-Target Organism Advisory:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This SLN label and the main EPA-registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label. Read this SLN label and the label affixed to the container for Stinger™ before applying.

Application Rates:

Target Weeds	Application Rate (pint/acre)	Use Restrictions (All Rates)
buckwheat, wild chamomile clover cocklebur, common dandelion galinsoga lettuce, prickly pineappleweed ragweed smartweed	1/4 - 1/2 (0.09375 - 0.187 lb ae/acre)	<ul style="list-style-type: none">• Make 1 to 2 broadcast application per crop per year, not to exceed a total of 1/2 pint per acre (0.187 lb ae/acre) per year.• Do not apply more than ½ pint of Stinger (0.187 lb ae) per acre per year.• Do not make more than 2 applications per acre per year.• For use under this SLN label, do not tank mix Stinger with other herbicides.

Target Weeds	Application Rate (pint/acre)	Use Restrictions (All Rates) – CONT.
sowthistle, annual ¹ thistle, Canada ¹	1/3 – 1/2 (0.125 – 0.187 lb ae/acre)	<ul style="list-style-type: none"> • Make 1 to 2 broadcast application per crop per year, not to exceed a total of 1/2 pint per acre (0.187 lb ae/acre) per year. • Do not apply more than ½ pint of Stinger (0.187 lb ae) per acre per year. • Do not make more than 2 applications per acre per year. • For use under this SLN label, do not tank mix Stinger with other herbicides. • Do not apply Stinger within 6 to 8 hours of expected rainfall or irrigation. • Chemigation: For use under this SLN label, do not apply this product through any type of irrigation system. • For use under this SLN label, do not aerially apply Stinger. • Avoid spray drift. Follow guidelines in the Avoiding Injury to Non-Target Plants and Spray Drift Management sections of the main EPA-registered label. • Follow the rotational crop restrictions and precautions under the Crop Rotation Intervals section of the main EPA-registered label. • Do not use plant residues, including hay or straw from treated areas, seed screenings, or manure or bedding straw from animals that have grazed or consumed forage from treated areas, for composting or mulching, where susceptible plants may be grown the following season. Do not spread manure from animals that have grazed or consumed seed screenings, forage or hay from treated areas on land used for growing susceptible broadleaf crops. Because crop injury may occur, do not use seed screenings/plant residue (even if composted in growing media). Do not apply seed screenings/plant residue to land that might be planted to a susceptible crop. To promote herbicidal decomposition, plant residues should be evenly incorporated or burned. Breakdown of clopyralid in crop residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation. Follow all applicable precautions and restrictions on this SLN label and the main EPA-registered label regarding composting and the use of treated plant soil, treated plant material, and manure of livestock that have grazed on treated plant material.

¹Suppression only.

Broadcast Application Rates: Apply uniformly with ground equipment in a minimum of 10 to 40 gallons of water per acre. For suppression of Canada thistle, apply after the majority of basal leaves have emerged but prior to bud stage and at least 30 days prior to harvest.

Application Timing: Risk of crop injury may increase when Stinger is applied during the bolting and/or

flowering stages of crop growth. If sprays are needed at this time it is recommended to use a lower rate within the application rate range.

Specific Use Precautions:

- Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

Special Crop Use Restrictions:

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown for Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to, green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

Special Conditions and Risks of Use

USE OF Stinger® herbicide (THE "PRODUCT") ON RADISH GROWN FOR SEED (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this Special Local Need Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Special Local Need Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Special Local Need Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and/or Crop loss;**
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Special Local Need Labeling;**
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop**

and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;

- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to inability to sell the Crop, Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of Oregon.

24(c) Registrant: Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, Indiana 46268
800-992-5994 (option #4)

™® Trademarks of Corteva Agriscience and its affiliated companies

R043-202
Approved: 02/16/23
Replaces: R043-114