



FIFRA

Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
PENNSYLVANIA**

Dual MAGNUM[®]

EPA Reg. 100-816

EPA SLN No. PA-080003

FOR CONTROL OF WEEDS IN EDAMAME

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Dual MAGNUM label.

This labeling must be in the possession of the user at the time of herbicide application.

Dual MAGNUM will not control emerged weeds. For effective weed control, Dual MAGNUM must be applied to clean-tilled soil.

Chemigation: Do not apply this product through any irrigation system.



EDAMAME

Dual MAGNUM may be applied preplant or pre-emergence for the control or suppression of grass and small seeded broadleaf weeds in edamame. For specific rates, see the rate table listed below.

Preplant Surface-Applied: For minimum-tillage or no-tillage systems only, Dual Magnum alone may be applied up to 45 days before planting. Use only split applications for treatments made 30-45 days before planting, with 2/3 the recommended broadcast rate for the crop and soil texture applied initially and the remaining 1/3 applied at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank mixture combination with a burndown herbicide (for example, Gramoxone Inteon™, Touchdown® or Roundup®). Observe directions for use, precautions, and restrictions on the label of the burndown herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated: Apply Dual Magnum to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate Dual Magnum after bed formation, unless specified otherwise.

Preemergence: Apply Dual Magnum during planting (behind the planter) or after planting, but before weeds or crops emerge.

Dual Magnum Broadcast Rates Per Acre

Soil Texture	Percent Organic Matter in Soil	
	< 3%	≥ 3%
Coarse	1 – 1.33 pts.	1.33 pts.
Medium	1.33 – 1.67 pts.	1.33 – 1.67 pts.
Fine	1.33 – 1.67 pts.	1.67 – 2.0 pts.

Restrictions and Precautions:

1. Dual Magnum will not control emerged weeds
2. Do not cut for hay within 120 days following a Dual MAGNUM application
3. Do not use for forage within 60 days following a Dual II MAGNUM application
4. Do not apply more than 2.0 pts./A of Dual MAGNUM during any one crop year.

Dual Magnum®, Touchdown® trademark of Syngenta Group Company,
Roundup® trademark of Monsanto Technology LLC

©2008 Syngenta

24(c) Registrant:
Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27419

PA0816014AA0408