



FIFRA 24(c) SPECIAL LOCAL NEED 24(C) LABELING

Lorox[®] DF

Agricultural Herbicide

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WISCONSIN ON CARROTS GROWN ON COARSE-TEXTURED, LOW ORGANIC MATTER SOILS TO CONTROL BROADLEAF AND GRASS WEEDS

EPA Reg. No. 61842-23 EPA SLN No. WI-150004

This SLN expires and must not be distributed or used in accordance with this SLN registration after 12/31/2019

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label. This labeling must be in the possession of the user at the time of pesticide application. This Section 24(c) SLN label contains new or supplemental instructions for use of this product, in Wisconsin, which do not appear on the EPA-registered label. Follow the instructions carefully.

Because carrot varieties vary in their resistance to herbicides, determine tolerance to LOROX DF prior to adoption as a field practice to prevent possible crop injury.

- Do not exceed 4 lbs. LOROX DF per acre per year.
- Do not apply within 14 days of harvest.
- Do not apply by air.

Do not apply LOROX DF to the following soils if the depth to groundwater is 30 feet or less:

- Sand with less than 3% organic matter
- Loamy sand with less than 2% organic matter
- Sandy loam with less than 1% organic matter

Do not apply LOROX DF in any manner that would contaminate irrigation water used for crops other than corn or water used for domestic purposes.

Do not apply LOROX DF within 150 feet of any well where depth to groundwater is 30 feet or less.

Preemergence on coarse-textured soils (such as sands or loamy sands and soils with less than 1% organic matter):

A single application of 0.25 to 0.5 lb. per acre should be made after planting but prior to carrots emerging. Seed should be planted at least 1/2 inch deep. The lower rate should be used on lighter soils and the higher rate on heavier soils. Additional postemergence applications may be made as long as the total does not exceed the seasonal maximum of LOROX DF.

Postemergence on coarse-textured soils (such as sands or loamy sands and soils with less than 1% organic matter):

Apply up to 1 lb. per acre as a broadcast spray. Applications to carrots that are less than 3 inches tall may result in crop injury; grower and/or applicator assumes all crop injury risk if applications are made to carrots less than 3 inches tall. Repeat application may be made as long as the total LOROX DF use does not exceed the seasonal maximum of LOROX DF.

Precautions for Postemergence Carrot Applications:

- Do not treat susceptible varieties which show an initial burning of foliage following postemergence treatment with LOROX DF.
- Do not exceed 40 psi spray nozzle pressure as crop injury may result.
- Do not apply when the temperature exceeds 85° F as crop injury may result.

The activity of LOROX DF on both carrots and weeds is increased if applied after 3 or more cloudy days. If spraying is done under these conditions, the dosage of LOROX DF must be reduced.

The addition or tankmix of Stoddard solvent, surfactants, nitrogen, or fertilizer solution, or other pesticides may cause crop injury. The grower/applicator assumes all risks in the tankmix situations.

LOROX DF often interacts with other herbicides or insecticides and may damage carrots when the chemicals are tank mixed or applied sequentially at close intervals. Several days, preferably a week should elapse between LOROX DF applications and application of insecticides.

Environmental Hazards

- It is a violation of federal law to harm an endangered species. Use of this product may be in counties which have endangered and threatened species present; therefore, applicators should evaluate the areas to be treated and take necessary precautions to avoid harming endangered species or their habitat/environment. Applicators may refer to http://datcp.wi.gov/Farms/Endangered_Species/ for help in identifying the species and their habitats. Growers who are uncertain of the presence of a protected species should contact the Wisconsin Department of Agriculture, Trade and Consumer Protection at 608- 224-4547.
- Linuron is known to leach through soil into groundwater under certain conditions as a result of labeled use on coarse textured soils with low organic matter. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

For product information, call 1-800-525-2803. You may also contact 1-866-374-1975 for emergency medical treatment information.

To the extent consistent with applicable law, Tessenderlo Kerley, Inc makes this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use LOROX DF must be made by each individual LOROX DF user and/or grower on the basis of possible crop injury from LOROX DF, the severity of weed infestation, the cost of alternative weed controls, and other factors. Read the Warranty and Limitation of Damages on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Warranty and Limitation of Damages on the product label.

Section 24(c)

Rev. 05/01/2015

Tessenderlo Kerley, Inc.
2255 N. 44th Street, Suite 300 Lorox® and NovaSource® are registered trademarks of Tessenderlo Kerley, Inc
Phoenix, AZ 85008 USA ©2013, 2014 Tessenderlo Kerley, Inc. All Rights Reserved
1-800-525-2803
www.novasource.com

