



SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR LOROX DF AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

This labeling must be in the possession of the user at the time of pesticide application.

"Label" as used in this supplemental labeling refers to the label booklet for LOROX® DF and this supplement.

This supplemental labeling expires on May 15, 2019 and must not be used or distributed after this date.

Lorox® DF

Agricultural Herbicide

EPA Reg. No. 61842-23

FOR WEED CONTROL IN ASPARAGUS

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ASPARAGUS

Do not exceed a maximum of 4 lbs./acre/year of LOROX® DF (2 lbs./acre/year of linuron active ingredient (a.i.)).

Do not use surfactant or fertilizer solution in spray mixture.

Do not apply within 1 day of harvest.

Direct Seeded or Newly Planted Crowns:

Preemergence: Make a single application of 2 to 4 lbs. per acre of LOROX® DF (1 to 2 lbs. per acre linuron a.i.) after planting seed 1/2 inches deep in coarse soil and 1 inch deep in fine soils. During planting operation, spray activated charcoal as a 1 inch band on soil surface directly over seed rows at the rate of 300 lbs. of LOROX® DF per acre (150 lbs. per acre linuron a.i.) (equivalent to 15 lbs. of LOROX® DF (7.5 lbs. linuron a.i.)) per acre of crop with a 20 inch row spacing). Preemergence weed control will be reduced in soils with high organic matter (greater than 5% and peat or muck).

Postemergence: Make 1 or 2 applications of 1 to 2 lbs. per acre of LOROX® DF (0.5 to 1 lb. linuron a.i.) per acre when ferns are in the 6 to 18 inch stage and weeds are not over 4 inches tall.

Established Beds:

Preemergence: Make a single application of 2 to 4 lbs. of LOROX® DF (1 to 2 lbs. linuron a.i.) per acre. Preemergence weed control will be reduced in soils with high organic matter (greater than 5% and peat or muck).

Postemergence: Make 1 to 3 applications of 1 to 2 lbs. of LOROX® DF (0.5 to 1 lb. linuron a.i.) per acre before weeds exceed 4 inches in height. Apply before cutting season or immediately after cutting.

Directed Postemergence (Fern Stage): Make a single application of 4 lbs. of LOROX® DF (2 lbs. linuron a.i.) per acre as directed. Spray to base of plants for control of dudain melon.

In California, dudain melon and annual nightshade are controlled.

Read the **WARRANTY AND LIMITATION OF DAMAGES** on the **Section 3 Federal product label** before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Warranty and Limitation of Damages on the Section 3 Federal product label.

Tessengerlo Kerley, Inc.
2255 N. 44th Street, Suite 300
Phoenix, AZ 85008 USA
1-800-525-2803
www.novasource.com

Lorox is a registered trademark of Tessengerlo Kerley, Inc.
05042016EPA