

For additional information  
contact ARVESTA CORPORATION  
Toll Free 877-448-6636

## SUPPLEMENTAL LABELING

# ELEVATE™

## 50 WDG FUNGICIDE



EPA REG NO. 66330-35

For Control of *Botrytis cinerea* Diseases of Bushberries, Caneberries and Pistachios  
Approved by EPA April 22, 2002

-It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
-This labeling must be in the possession of the user at the time of application.  
-Read the label affixed to the container for ELEVATE 50 WDG fungicide before applying. Carefully follow all precautionary statements and applicable use directions.  
-Use of ELEVATE 50WDG according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for ELEVATE 50 WDG except for the AGRICULTURAL USE REQUIREMENTS section. As requested by EPA 4/22/02 this section is revised as indicated below.

### DIRECTIONS FOR USE

**BUSHBERRIES (Blueberries, gooseberries, currants and huckleberries):** For control of *Botrytis cinerea* (Gray mold), apply 1.5 pounds ELEVATE 50 WDG Fungicide per acre (0.75 lb AI/A). Begin application at 10 % bloom and continue through harvest. Applications should be made every 7 days or when conditions favor disease development. The final application may be made up to and including the day of harvest (PHI=0). DO NOT apply more than 6.0 pounds of product per acre per season (3.0 lb AI/A/season)

**CANEBERRIES (Red and black raspberries, blackberries, and loganberries):** For control of *Botrytis cinerea* (Gray mold), apply 1.5 pounds ELEVATE 50 WDG fungicide per acre (0.75 lb AI/A) Begin application at 10% bloom and continue through harvest. Applications should be made every 7 days or when conditions favor disease development. The final application may be made up to and including the day of harvest (PHI=0). DO NOT apply more than 6.0 pounds of product per acre per season (3.0 lb AI/A/ season).

**PISTACHIOS:** For the control of Botrytis blossom and shoot blight (*Botrytis cinerea*) apply 1.0 to 1.5 lbs of ELEVATE 50 WDG Fungicide per acre (0.5 to 0.75 lb AI/A) at 5-10% bloom, full bloom, then every 7-10 days or when conditions favor disease development up to 28 days after petal fall. Apply in sufficient water to obtain complete coverage of flowers, fruit and twigs. DO NOT apply more than 6.0 pounds of product per acre per season (3.0 lb AI/A/ season).

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training decontamination, notification, and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notifications to workers, restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, socks shoes and chemical resistant gloves made of any waterproof material.

ARVESTA CORPORATION, 100 First St., Suite 1700, San Francisco, CA. 94105