

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA Reg. No. 279-3108

For Agricultural Use Only

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

CROP	PEST	RATE OF APPLICATION
Citrus	Fuller Rose Beetle <i>Asynonychus godmani</i>	2.5 – 5.0 lbs product/A (0.25 – 0.5 lb ai/A)

DIRECTIONS FOR USE: A single or split application as a soil or trunk treatment may be made depending on pest emergence.

Soil Treatment: Make a uniform application to the soil beneath trees from the trunk to the drip line in a minimum of 40 gallons finished spray solution per acre. See product label for full soil application details.

Trunk Treatment: Apply to the lower trunk as a directed spray. Uniform coverage of the trunk is critical for pest control. Spray the trunks of 10 trees with water to determine the amount of spray that is needed for thorough coverage of the trunk, without runoff, from the soil line to approximately 18 inches up the trunk. Measure the amount of water needed to refill the spray tank to its initial level. Then calculate the amount of spray needed per acre, e.g. if it took 0.5 gallons to spray 10 trees and there are 180 trees per acre then you will need $0.5 \times (180/10) = 9$ gal/A of spray mix. Based on the number of acres to be sprayed, put that amount of water per acre in the tank along with the appropriate amount of Brigade WSB (5 lbs/A for a single application or 2.5 lbs/A for a split application)

Single Application: A single application of 5.0 lbs product/A (0.5 lb ai/A) may be made before pest emergence.

Split Application: A split application using one application of 2.5 lbs product/a (0.25 lb ai/A) followed later by a second application of 2.5 lbs product/A (0.25 lb ai/A) may be made. Applications should be made prior to peak pest emergence.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Brigade is a FMC Trademark
©2012 FMC Corporation

RESTRICTIONS:

- Do not apply through irrigation systems.
- Do not allow Brigade WSB to contact fruit or foliage. Contact with fruit may result in illegal residues.
- Do not apply more than 5.0 pounds product (0.5 lbs active ingredient) per acre per year.
- Do not apply within 1 day of harvest.
- Do not apply by air.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ARE ON THE EPA REGISTERED BRIGADE® WSB INSECTICIDE/MITICIDE LABEL (EPA REG. NO. 279-3108) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Brigade is a FMC Trademark
©2012 FMC Corporation