

**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms.**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



**2(ee) Recommendation**

**FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA**

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING**

**For Agricultural or Commercial Use Only**

CROP	PEST	RATE OF APPLICATION
Strawberries (0 Day PHI)	<i>Vinegar Flies (adults)</i> ( <i>Drosophila spp.</i> Including Spotted-wing <i>Drosophila</i> )	8 - 32 oz / acre.

**DIRECTIONS FOR USE:** Through coverage is essential to achieve control. Use a minimum spray volume of 5 gallons by air or 50 gallons by ground to obtain full coverage of foliage. Multiple applications may be required.

**RESTRICTIONS:**

Do not apply more than 80 ounces product (0.5 lbs active ingredient) per acre per season. Harvesters and other personnel performing tasks with all-day foliage contact in treated fields within five (5) days of application must wear long-sleeved shirt, long pants, and shoes plus socks.

Following treatment of strawberry fields at rates of Brigade® WSB greater than 0.1 lb ai/acre, harvesters must wear gloves for five (5) days following application.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED BRIGADE® WSB INSECTICIDE/MITICIDE LABEL (EPA REG. NO. 279-3108) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**



FMC Corporation  
Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Mustang and FMC - FMC Trademarks  
©2009 FMC Corporation