

# FIFRA Section 24(c) Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Strongarm®

EPA Reg. No. 62719-288

EPA 24(c) Special Local Need Registration SLN SC-160002

### Postemergence Application for Suppression of Tropical Spiderwort (*Commelina benghalensis*) Control in Peanuts

(For Distribution and Use Only in the State of South Carolina)

This label expires and must not be distributed or used in accordance with the SLN registration after December 31, 2021

#### Active Ingredients:

diclosulam: N-(2,6-dichlorophenyl)-5-ethoxy-7-fluoro[1,2,4]triazolo-[1,5-c]pyrimidine-2-sulfonamide .....	84%
Other Ingredients .....	16%
Total .....	100%

Contains 0.84 lb of active ingredient per pound of product

U.S. Patent No. 5,163,995

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Strongarm® herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as described in this Special Local Need labeling, use of Strongarm is subject to all precautions and limitations imposed by the label affixed to the product container.

#### Directions for Use

Apply Strongarm in a broadcast spray when tropical spiderwort is in the 1 to 2 leaf stage of growth and actively growing. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control. Optimum weed control will be obtained by applying Strongarm under favorable growing conditions (i.e., adequate soil moisture and temperature). Applications may occur from peanut cracking, to the initiation of flowering (bud stage), but no later than 30 days after planting.

Weeds Suppressed	Strongarm (oz/acre)
Tropical Spiderwort ( <i>Commelina benghalensis</i> )	0.45

**Surfactant Use:** Applications of Strongarm must include a nonionic surfactant at 1 quart per 100 gallons (0.25% v/v). When mixing, add nonionic surfactant last.

**Tank Mix Applications:** Strongarm may be applied alone or in tank mix combination with other postemergence herbicides registered for use in peanuts. When applied in tank mix combination with other herbicides, follow applicable use instructions, including application rates, precautions and restrictions for each product used in the tank mixture, including use of adjuvants. The most restrictive limitations on each tank mix label must be followed.

**Rotation to Cotton:** Except for rotation to cotton, follow crop rotation intervals specified in the Crop Rotation Intervals for Common Crops section of the label affixed to the product container. For postemergence use in peanuts, the minimum interval for rotation to cotton is 18 months or a successful field bioassay. See instructions for Field Bioassay in the Crop Rotation Interval for Common Crops section of the label affixed to the product container.

**Maximum Application Rates:** In peanuts, do not apply more than 0.45 oz of Strongarm per acre per year in any combination of preplant incorporated, preplant surface, preemergence through cracking or postemergence applications.

**Crop Tolerance:** Strongarm may cause temporary yellowing or chlorosis of peanut foliage after application. These symptoms will vary in degree of severity, depending on environmental and crop growth factors (e.g., temperature and soil moisture). If these symptoms as described are not acceptable, do not use Strongarm as a postemergence application in peanuts.

®Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

R044-010

Accepted: 05/18/16

Initial printing