

**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms.**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



**2(ee) Recommendation**

**FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA**

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING**

EPA Registration No. 279-3126

CROP	PEST	RATE OF APPLICATION
Fruiting Vegetables (except Cucurbits) Leafy Vegetables (except Brassica) Head and Stem Brassica Vegetables Leafy Brassica Greens Legume Vegetables (except Soybeans)	Symphylans	4.3 fl oz/A (0.05 lb ai/A)

**DIRECTIONS FOR USE:**

Apply Mustang Insecticide before, during, or after planting for protection from symphylans. Placement of Mustang Insecticide is critical to get plant protection and will depend on seed type and placement, or transplant placement. Successful protection from symphylans is obtained by having Mustang Insecticide-treated soil around or in contact with developing plant roots. Whatever application method is used must result in proper placement of Mustang Insecticide to get plant protection. Successful application methods include (but may not be limited to) the following:

**Direct-seeded crops:** Mustang Insecticide should be applied in a narrow band (3-6 inches) and incorporated throughout the seed line. Incorporation should result in treated soil around and below (1-3 inches) the seed. Mechanical and sprinkler incorporation have been successful, but mechanical incorporation has been more consistent. Where more than one seed line per bed are planted, multiple bands or bed top applications will be necessary. Use as narrow a band as practical.



FMC Corporation  
Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Mustang is a FMC Trademark  
©2014 FMC Corporation  
Version 6-20-2014

**Transplanted crops:** Apply Mustang Insecticide to the transplant row prior to transplanting in as narrow a band as practical (but not less than 4-6 inches) and mechanically incorporate to a depth of 4-6 inches. Mustang Insecticide may also be applied as a soil drench at the time of transplanting. Mustang insecticide has also given plant protection from symphylans when applied to the transplant row with two shanks four inches apart and six inches deep. Mustang Insecticide may also be applied with a spray blade so a barrier of treated soil immediately below the transplant is obtained.

**RESTRICTIONS:**

- Do not apply more than 4.3 fl oz/A (0.05 lb ai/A) per application.
- Do not apply more than 25.8 fl oz/A (0.3 lb ai/A) per season.
- Do not make applications of Mustang Insecticide less than 7 days apart on the listed crops except for Legume Vegetables where applications can be made 5 days apart.
- Do not apply within 1 day of harvest (1-Day PHI) for Fruiting Vegetables, Leafy Vegetables (except Brassica), Head and Stem Brassica Vegetables, Leafy Brassica Greens, and Legume Vegetables (succulent shelled or edible-podded peas or beans).
- Do not apply with 21 days of harvest (21-Day PHI) for Legume Vegetables (dried shelled peas and beans).

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG INSECTICIDE LABEL (EPA REG. NO. 279-3126) MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**



FMC Corporation  
Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Mustang is a FMC Trademark  
©2014 FMC Corporation  
Version 6-20-2014