

MEFENOXAM	GROUP	4	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

WARDEN® RTA®

Fungicide

A seed treatment fungicide for protection against damage from listed seed-borne, soil-borne, and seedling diseases of soybeans and other specified legume crops.

Active Ingredients:

Mefenoxam*	2.21%
Fludioxonil**	0.72%
<hr/>	
Other Ingredients:	97.07%
Total:	100.00%

*CAS Nos. 70630-17-0 and 69516-34-3

**CAS No. 131341-86-1

Warden RTA is a flowable concentrate for seed treatment containing 0.197 lb mefenoxam and 0.064 lb fludioxonil per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

[See additional precautionary statements and directions for use in booklet.]

EPA Reg. No. 100-1146-1381

EPA Est. 100-NE-001

SCPSR-WIN-1146C-L1F 0719 SU 59750
 2.5 gallons
 Net Contents

SCPSR-WIN-1146C-L4D 0719 SU 59749
 15 gallons
 Net Contents

Distributed by:
 Winfield Solutions, LLC
 PO Box 64589
 St. Paul, MN 55164-0589

1/0719/9

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24-Hour Medical Emergency Assistance (Human or Animal) Call 1-877-424-7452 or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-424-9300	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants

- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils or Viton® ≥14 mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not contaminate water when disposing of equipment wash-water or rinsate.

Groundwater Advisory

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER AND USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED

CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USER OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUES, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the seller, and purchase price will be refunded.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use is permitted on-farm and in commercial seed treatment facilities. Do not use for at-plant applications (e.g. hopper box, planter box, etc.). This product is to be applied as a water-based slurry through standard slurry- or mist-type commercial seed treatment equipment.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Winfield Solutions, LLC no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed when treated with Warden RTA.

PRODUCT INFORMATION

Warden RTA is a seed treatment fungicide that protects against damping-off and seed rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* spp., and early season *Phytophthora* root rot. Warden RTA also suppresses seed-borne *Sclerotinia* and *Phomopsis* spp. Warden RTA contains the active ingredient mefenoxam found in Apron XL® that provides protection against *Pythium* and early season *Phytophthora* root rot. Warden RTA also contains the active ingredient fludioxonil found in Maxim® 4FS that provides protection against the remaining diseases mentioned in this label.

RESISTANCE MANAGEMENT

MEFENOXAM	GROUP	4	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

For resistance management, please note that Warden RTA contains both a Group 4/mefenoxam and Group 12/fludioxonil fungicide. Any fungal population may contain individuals naturally resistant to Warden RTA and other Group 4 or Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Warden RTA or other Group 4 or Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide application. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Winfield Solutions, LLC at 1-855-494-6343. You can also contact your pesticide distributor or university extension specialist to report resistance.

Winfield Solutions, LLC encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

MIXING PROCEDURES

Important: Always re-circulate Warden RTA thoroughly before using.

Follow the manufacturer's application instructions for the seed treatment equipment being used.

Apply Warden RTA as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the

specified amount of Warden RTA into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

Certain crops require addition of inoculants when the seed is treated or planted. Warden RTA is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Winfield Solutions, LLC representative for directions before applying Warden RTA with inoculants.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants. The minimum slurry volume to achieve adequate coverage is 4.0 fl oz/100 lb seed. More diluent may be required to obtain complete coverage.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution. Clean out any unused product from the treater after treating or maintain constant agitation if the leftover slurry will be maintained overnight. Warden RTA already contains an EPA-approved dye or colorant that imparts an unnatural color to the seed as stated in 40 CFR 153.155(c). Allow seed to dry before bagging.

SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that containers of treated seeds shall be labeled with the following statements:

- This seed has been treated with mefenoxam and fludioxonil fungicides.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on containers of seeds treated with Warden RTA:

- Ground Water Advisory: Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Excess treated seed may be used for ethanol production only if:
 - (1) By-products are not used for livestock feed, and
 - (2) No measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Do not allow children, pets, or livestock to have access to treated seed.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.

- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed by burying away from bodies of water.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Do not use at a rate that will result in more than 0.004 lb fludioxonil per acre (2.0 grams ai/A) per year and 0.013 lb mefenoxam per acre (5.7 grams ai/A) per year as a seed treatment application.

CROP USE DIRECTIONS

When applied according to the **WARDEN RTA RATE TABLE**, Warden RTA provides protection against damping-off and seed borne rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* species and early season *Phytophthora* root rot. Warden RTA also suppresses seed-borne *Sclerotinia* and *Phomopsis* species. Additional Apron XL, EPA Reg. No. 100-799, may be necessary for high levels of *Phytophthora* or *Pythium*. Refer to the table below to determine the appropriate use rate of Apron XL, if needed. Read and follow all label directions for Apron XL use.

Legume Vegetables (Succulent or Dried) Crop Group 6 & Foliage of Legume Vegetables Crop Group 7

Bean (*Lupinus* spp.)

(grain lupin, sweet lupin, white lupin, white sweet lupin)

Bean (*Phaseolus* spp.)

(field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

Bean (*Vigna* spp.)

(adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean)

Broad bean (fava bean)

Chickpea (garbanzo bean)

Guar

Jackbean

Lablab bean (hyacinth bean)

Lentil

Pea (*Pisum* spp.)

(dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

Pigeon pea

Sword bean

Soybean

Soybean (immature seed)

WARDEN RTA RATE TABLE

Pathogens	Use Rate			
	fl oz/100 lb seed		g ai/100 kg	
	Warden RTA	Apron XL	Warden RTA	Apron XL
<i>Pythium</i> , <i>Fusarium</i> , <i>Rhizoctonia</i> , seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> seed decay	5.0	-	Fludioxonil 2.5 Mefenoxam 7.5	-
<i>Pythium</i> (high pressure), <i>Phytophthora</i> , <i>Fusarium</i> , <i>Rhizoctonia</i> , seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> seed decay	5.0	0.32	Fludioxonil 2.5 Mefenoxam 7.5	Mefenoxam 7.5

WARDEN RTA RATE TABLE (continued)

Pathogens	Use Rate					
	fl oz/140,000 seeds*		Total mg ai/seed*		g ai/100 kg	
	Warden RTA	Apron XL	Warden RTA	Apron XL	Warden RTA	Apron XL
<i>Pythium</i> , <i>Fusarium</i> , <i>Rhizoctonia</i> , seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> seed decay	2.33	–	0.015	–	Fludioxonil 2.5 Mefenoxam 7.5	–
<i>Pythium</i> (high pressure), <i>Phytophthora</i> , <i>Fusarium</i> , <i>Rhizoctonia</i> , seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> seed decay	2.33	0.15	0.015	0.011	Fludioxonil 2.5 Mefenoxam 7.5	Mefenoxam 7.5

*Rates expressed as fl oz/140,000 seeds and mg ai/seed are based on an average seed weight of 3,000 seeds/lb.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place. Store in an area designated for pesticides and away from food or feed. Store in original containers only. Keep container closed when not in use.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons – mini-bulk]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons - bulk]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, or accident, call CHEMTREC 1-800-424-9300

Apron XL®, Maxim® and RTA® are trademarks of a Syngenta Group Company

Warden® is a registered trademark of Winfield Solutions, LLC.

©2019 WINFIELD

Distributed by:
Winfield Solutions, LLC
PO Box 64589
St. Paul, MN 55164-0589

SCPSR-WIN-1146C-L1F 0719
2.5 gallons
Net Contents

SU 59750

SCPSR-WIN-1146C-L4D 0719
15 gallons
Net Contents

SU 59749

Base Label – 2.5 gallons and 15 gallons

MEFENOXAM	GROUP	4	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Warden® RTA®

Fungicide

A seed treatment fungicide for protection against damage from listed seed-borne, soil-borne, and seedling diseases of soybeans and other specified legume crops.

Active Ingredients:

Mefenoxam*	2.21%
Fludioxonil**	0.72%
<hr/>	
Other Ingredients:	97.07%
Total:	100.00%

*CAS Nos. 70630-17-0 and 69516-34-3

**CAS No. 131341-86-1

Warden RTA is a flowable concentrate for seed treatment containing 0.197 lb mefenoxam and 0.064 lb fludioxonil per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

[See additional precautionary statements and directions for use in booklet.]

EPA Reg. No. 100-1146-1381

EPA Est. 100-NE-001

SCPSR-WIN-1146C-L1F 0719
2.5 gallons
Net Contents

SU 59750

SCPSR-WIN-1146C-L4D 0719
15 gallons
Net Contents

SU 59749

Distributed by:
Winfield Solutions, LLC

PO Box 64589
St. Paul, MN 55164-0589

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See directions for use in attached booklet.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **HOT LINE NUMBER:** For 24 Hour Medical Emergency Assistance (Human or Animal), Call **1-877-424-7452** or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), **Call 1-800-424-9300.**

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not contaminate water when disposing of equipment wash-water or rinsate.

Groundwater Advisory

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place. Store in an area designated for pesticides and away from food or feed. Store in original containers only. Keep container closed when not in use. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, or accident, call CHEMTREC 1-800-424-9300

RTA® is a trademark of a Syngenta Group Company

Warden® is a registered trademark of Winfield Solutions, LLC