

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid[®] 2F EPA Reg. No. 62719-442

Control of Lepidoptera Larvae Infesting Pome Fruit (Crop Group 11-10)¹

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid[®] 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

¹Pome fruit (crop group 11-10) including apple, Asian pear, azarole, crabapple, loquat, mayhaw, medlar, pear, quince, tejocote, cultivars, varieties, and/or hybrids of these

Refer to product label for General Information, Mixing and Application directions.

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
codling moth (suppression only) For use against low to moderate infestations in conjunction with alternate control measures such as in established mating disruption blocks.	16 (0.25 lb ai/acre)	For each generation, apply at the initiation of egg lay (usually occurs at 100 to 200 DD, base 50°F, following biofix). Reapply 10 to 18 days later.	<ul style="list-style-type: none"> • Do not apply more than a total of 64 fl oz of Intrepid 2F (1 lb ai) per acre per season • Preharvest Interval: Do not apply within 14 days of harvest. • Aerial application is allowed only for the last two applications prior to harvest.
lesser appleworm oriental fruit moth	12 – 16 (0.19 - 0.25 lb ai/acre)		
obliquebanded leafroller pandemis leafroller	8 – 16 (0.12 - 0.25 lb ai/acre)	<p>Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending upon infestation level.</p> <p>Summer generation: Make the first application during the period of peak egg lay to early egg hatch (usually 200 to 400 DD following biofix). Reapply 10 to 18 days later (usually 500 to 700 DD).</p>	

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
eyespotted bud moth fruittree leafroller light brown apple moth redbanded leafroller variegated leafroller		For control of surface or foliar feeding leafroller larvae, apply when larvae are feeding.	
tufted apple bud moth	6 – 10 (0.09 - 0.16 lb ai/acre)	For each generation, apply at 10 to 30% egg hatch.	
spotted tentiform leafminer western tentiform leafminer	8 – 12 (0.12 - 0.19 lb ai/acre)	First generation: Apply at pink to petal fall. Second, third generation: Apply at early egg hatch for each generation.	
lacanobia fruitworm	12 (0.19 lb ai/acre)	Apply at egg hatch or at the first sign of larval infestation. Reapply within 10 to 14 days.	

For best protection, begin applications before egg hatch of each generation. For pests that penetrate fruit apply Intrepid 2F before the larval hatch and penetrate the fruit. Intrepid 2F provides 10 to 18 days of protection depending upon application rate and how rapidly fruit and/or leaves are expanding. Most effective crop protection results from an application of Intrepid 2F made at the initiation of egg hatch. For heavy infestations, continuous moth flight and egg laying, or extended egg hatch, use the maximum specified rates. Maintain coverage on the fruit surface with 10- to 18-day re-treatment intervals.

Intrepid 2F may also be used in a program approach alternated or interspersed with other insecticides. Make sure the re-treatment interval does not exceed the period of effectiveness of the alternate products and Intrepid 2F.

Consult local spray timing advisories or follow biofix dates based upon pheromone trap catches to time sprays appropriately.

Ground Application: Apply Intrepid 2F by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trellised trees or trees 10 feet tall or less. For trees greater than 10 feet tall use a minimum of 100 gallons per acre.

Aerial Application: Aerial application is allowed only for the last two applications prior to harvest. Apply Intrepid 2F in a minimum of 20 gallons per acre. Intrepid 2F can be applied by aerial applications when conditions warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Spray Adjuvants: The addition of agricultural adjuvants to sprays of Intrepid 2F may improve initial spray deposits, redistribution and weatherability. Select adjuvants that are recommended and registered for your specific use pattern and follow their use directions. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Always add adjuvants last in the mixing process.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

® Trademark of Dow AgroSciences LLC

R846-079

EPA accepted 10/29/13

Initial printing.