

## Supplemental Label



### Reduced preharvest interval (PHI) in Bermudagrass and other perennial warm-season grasses grown for forage or hay production, or in pastures or rangeland

This supplemental label expires April 30, 2019, and must not be used or distributed after this date.

**Active Ingredient\*:**

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine ..... 38.7%

**Other Ingredients:** ..... 61.3%

**Total:** ..... 100.0%

\*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

**EPA Reg. No. 241-418**

**CAUTION**

---

#### Directions For Use

---

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Prowl<sup>®</sup> H<sub>2</sub>O herbicide** container label, EPA Reg. No. 241-418, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Prowl H<sub>2</sub>O** before applying.
- Use of **Prowl H<sub>2</sub>O** according to this labeling is subject to the use precautions and restrictions imposed by the label affixed to the container for **Prowl H<sub>2</sub>O**.

---

#### Product Information

---

**Prowl H<sub>2</sub>O** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to the **Prowl H<sub>2</sub>O** container label for a complete list of weeds controlled.

---

#### Application Instructions

---

**Prowl H<sub>2</sub>O** may be applied by ground or chemigation, by air, or on dry bulk fertilizer.

#### Use Method, Rate, and Timing

##### Established Bermudagrass and other Perennial Warm-season Grasses

Apply **Prowl H<sub>2</sub>O** in fields of Bermudagrass (and other perennial warm-season grasses including Bahiagrass, buffalograss, switchgrass, and others) grown for forage or hay production, and/or grown in pastures or rangeland for livestock grazing.

Apply **Prowl H<sub>2</sub>O** only to established (defined as planted in fall or spring which has gone through a first cutting/mowing) Bermudagrass and other perennial warm-season grasses.

Apply **Prowl H<sub>2</sub>O** before target weed germination in the dormant season [i.e., when grasses are not actively growing in fall (postharvest), during winter dormancy period, or in early spring before greenup] or in-season between cuttings. **DO NOT** apply to Bermudagrass and other perennial warm-season grasses after greenup in spring before the first cutting.

Uniformly apply at a broadcast rate of 1.1 quarts (1.0 lb ai/A) to 4.2 quarts (4.0 lbs ai/A) of **Prowl® H<sub>2</sub>O herbicide** per acre in a single application.

**Prowl H<sub>2</sub>O** may also be applied as a sequential or split program when the initial application(s) is made in the dormant season and the subsequent application(s) is made in-season between cuttings. **DO NOT** apply more than a maximum cumulative total of 4.2 quarts of **Prowl H<sub>2</sub>O** per acre per year.

Apply the higher rate of **Prowl H<sub>2</sub>O** when higher weed pressure is anticipated or when a longer duration of residual weed control is desired.

**Prowl H<sub>2</sub>O** may be applied in a sequential use program or as a tank mix with other registered herbicides that control emerged weeds.

**Prowl H<sub>2</sub>O** may cause temporary injury to Bermudagrass and other perennial warm-season grass stands. Disease, extremely cold weather, drought, extensive frost heaving, salinity, low pH, or high pH may weaken stands and make the crop more susceptible to herbicide injury.

### **Chemigation Application**

**Prowl H<sub>2</sub>O** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the **Spraying Instructions** section of the **Prowl H<sub>2</sub>O** container label.

### **Tank Mixes**

**Prowl H<sub>2</sub>O** may be tank mixed with other herbicides labeled for use in Bermudagrass and other perennial warm-season grass fields. BASF recommends testing **Prowl H<sub>2</sub>O** tank mixes on a small portion of the target crop to determine if damage is likely to occur.

Application of postemergence herbicides may cause crop injury, such as stunting or chlorosis of Bermudagrass and other perennial warm-season grasses. Consult your local BASF dealer for local tank mix options. Always perform a mixing test to check the compatibility of **Prowl H<sub>2</sub>O** with all potential tank mix partners and fertilizers. Follow all precautions and restrictions on the labels of all products applied in combination with **Prowl H<sub>2</sub>O**. Always follow the most restrictive label.

**Physical incompatibility, reduced weed control, or crop injury may result from mixing Prowl H<sub>2</sub>O with other pesticides, additives, or fertilizers.**

### **Crop-specific Restrictions**

- **DO NOT** apply **Prowl H<sub>2</sub>O** if surface water is present in the field.
- **DO NOT** apply more than a maximum cumulative total of 4.2 quarts of **Prowl H<sub>2</sub>O** per acre per year.
- There is no preharvest interval for grass forage or hay, or for livestock grazing after application of **Prowl H<sub>2</sub>O**.

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Prowl is a registered trademark of BASF.*

© 2016 BASF Corporation  
All rights reserved.

000241-00418.20160118b.**NVA 2016-04-195-0015**  
Based on: NVA 2016-04-195-0013  
Supersedes: NVA 2014-04-195-0026

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



We create chemistry