

TECHNICAL INFORMATION LABELING

ACEQUINOCYL GROUP 20B INSECTICIDE

KANEMITE® 15 SC MITICIDE

EPA Reg. No. 66330-38

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA.

For Use on Citrus Fruit Groups for Control of Yellow Mite and Flat Mites.

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of **KANEMITE® 15 SC MITICIDE (EPA 66330-38)**.

Citrus Fruit Group (Crop Group 10-10): Australian Desert Lime; Australian Finger-Lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus Hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Orange, Sour; Orange, Sweet; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate Orange; Uniq Fruit; Cultivars, Varieties, and/or Hybrids of these.

PEST	USE RATE
Yellow mite (<i>Lorryia formosa</i>)	21 fl oz/A – 31 fl oz/A
Flat mites (<i>Brevipalpus</i> spp.)	(0.2 lb ai/A – 0.3 lb ai/A)
For optimal control, use 150 gallons of water per acre or higher.	
RESTRICTIONS: <ul style="list-style-type: none">Do not use less than 100 gallons of water volume.Allow a minimum of 21 days between treatments.Do not make more than two applications of KANEMITE 15 SC MITICIDE per year.Do not apply more than 62 fl oz of KANEMITE 15 SC MITICIDE per acre (0.6 lb ai) per year.Preharvest interval is 7 days.Restricted-entry interval (REI) is 12 hours.	

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **KANEMITE 15 SC MITICIDE** BEFORE APPLYING. USE OF **KANEMITE 15 SC MITICIDE** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **KANEMITE 15 SC MITICIDE (EPA Reg. No. 66330-38)**.

120817V003CA2ee

Produced For:
ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

KANEMITE is a registered trademark of Agro-Kanesho Company.

The KANEMITE logo is a registered trademark of an Arysta LifeScience Group Company.

©2019 Arysta LifeScience Group Company.