

# RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

GROUP

3A

INSECTICIDE



I N S E C T I C I D E

For mixing directly with liquid fertilizer to control listed soil insect pests

EPA Reg. No. 279-3302

EPA Est. 279-NY-1

Active Ingredient:

By Wt.

Bifenthrin\* ..... 17.15%

Other Ingredients: ..... 82.85%

100.0%

\*Cis isomers 97% minimum, trans isomers 3% maximum.

This product contains 1.5 pounds active ingredient per gallon.



**KEEP OUT OF REACH OF CHILDREN  
WARNING  
AVISO**

This label must be in the possession of the user at the time of application.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for additional precautionary information.

**Net Contents: 2.5 Gallons**

Sold By



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

## FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency medical treatment information.

## NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

**For Emergency Assistance Call (800) 331-3148**

## PRECAUTIONARY STATEMENTS

### Hazards to Humans (and Domestic Animals)

#### Warning

May be fatal if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### Personal Protective Equipment:

Applicators and other handlers (other than mixers and loaders) must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

#### Mixers and Loaders must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

##### Users should:

- Remove PPE after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

### DIRECTIONS FOR USE

#### RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, and Shoes plus socks.

### Resistance Management

Some pests are known to develop resistance to insecticides that have been used repeatedly. While the development of insect resistance is well understood, it is not easily predicted. Therefore insecticides should be used in conjunction with the resistance management strategies in the area. Consult the local or State agricultural advisors for details. If insect resistance should develop in the area, this product used alone may not continue to provide sufficient levels of pest control. If the reduced levels of control can not be attributed to improper application techniques, improper use rates, improper application timing, unfavorable weather conditions or abnormally high pest pressure, a resistant strain may have developed.

To reduce the potential for pesticide resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the recommended rates and in accordance with the use directions. Do not use less than recommended label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. For optimum performance, scout fields carefully and begin applications when pests are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

### Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, drip irrigation, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For LEPA irrigation a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluents, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distributions is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment. Capture LFR Insecticide should be applied continuously for the duration of the water application. Capture LFR Insecticide should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inches per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

### Application and Mixing Instructions

Capture LFR Insecticide is an insecticide/miticide that contains 1.5 pounds of bifenthrin per gallon. Capture LFR Insecticide can be mixed directly with liquid fertilizer or with water. The rate of application is variable according to pest pressure, timing of treatments and field scouting. Use lower rates under light to moderate pest infestations, and higher rates under heavier pest pressure. In arid climates, applications rates are generally higher. Fill the tank one-half full with liquid fertilizer or water and begin spray tank agitation. Add the proper amount of Capture LFR Insecticide, and then add the rest of the fertilizer or water. Maintain agitation until the mixture has been applied.

Shake well before using.

Agitate the Capture LFR Insecticide spray solutions in nurse tanks prior to moving the solution to spray system.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Capture LFR Insecticide can be applied in-furrow with the seed, as a T-band (band over the open furrow), as a broadcast application, as a band over the row, as a pre-emergent (PRE) application, as a post plant incorporated (PPI) application, as a foliar application (includes chemigation), or as a transplant-water drench during setting.

Use rates in the individual crop sections for at-plant soil applications are listed as Fluid oz/1000 Linear ft based on 30 inch row spacings. For conversion to application rates applicable to other row spacings see the table below.

The maximum application rate per acre per season (pounds active ingredient /acre/season) as listed under the use directions for each crop must not be exceeded. This seasonal maximum includes at-plant plus foliar applications of Capture LFR Insecticide and other products containing bifenthrin.

For at-plant applications, Capture LFR Insecticide can be mixed with commonly used liquid starter or pop-up fertilizers. Follow liquid fertilizer recommendations regarding seed safety and use guidelines. Conduct a preliminary jar test using the appropriate ratio of fertilizer and Capture LFR Insecticide. Do not allow a tank mixture to set overnight, but if this occurs agitate tank mixture prior to application.

Row Spacing	36"	30"	20"	15"	Twin Row 30" centers
Linear row ft/acre	14,520 ft	17,424 ft	26,136 ft	34,848 ft	34,848 ft
<b>Conversion</b>					
0.2 Fluid oz /1000 Linear ft =	2.9 fl oz /acre	3.5 fl oz /acre	5.2 fl oz/acre	7.0 fl oz /acre	7.0 fl oz /acre
0.24 Fluid oz /1000 Linear ft =	3.5 fl oz /acre	4.2 fl oz /acre	6.3 fl oz/acre	8.4 fl oz /acre	8.4 fl oz /acre
0.3 Fluid oz /1000 Linear ft =	4.4 fl oz /acre	5.2 fl oz /acre	7.8 fl oz/acre	10.5 fl oz /acre	10.5 fl oz /acre
0.39 Fluid oz /1000 Linear ft =	5.7 fl oz /acre	6.8 fl oz /acre	10.2 fl oz b, /acre	13.6 fl oz /acre	13.6 fl oz /acre
0.49 Fluid oz /1000 Linear ft =	7.1 fl oz /acre	8.5 fl oz /acre	12.8 fl oz /acre		
0.73 Fluid oz /1000 Linear ft =	10.6 fl oz /acre	12.7 fl oz /acre			
0.78 Fluid oz /1000 Linear ft =	11.3 fl oz /acre	13.6 fl oz /acre			
0.98 Fluid oz /1000 Linear ft =	14.2 fl oz /acre	17.1 fl oz /acre			
1.47 Fluid oz /1000 Linear ft =	21.3 fl oz /acre	25.6 fl oz /acre			

### Crop Rotation Restrictions

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following the final application of bifenthrin.

### Tank-Mixtures

Capture LFR Insecticide may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products. To insure successful applications, product compatibility tests should be conducted.

### Maximum Allowable Capture LFR Insecticide Use Per Acre Per Season

Refer to the individual crop sections for maximum allowable Capture LFR Insecticide usage per acre per season. The maximum allowable use must include all registered use patterns including at-plant, soil applied and/or foliar applications for the 12 months period. The 12 month period is to begin upon the initial application to the acre.

### Buffer Zones

#### Vegetative Buffer Zones

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing bifenthrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21pp.

[http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs143\\_023819.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs143_023819.pdf).

**Buffer Zone for Ground Application (groundboom, overhead chemigation, or airstblast)** – Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

**Buffer Zone for ULV Aerial Application** - Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

**Buffer Zone for Non-ULV Aerial Application** – Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, natural ponds, estuaries, and commercial fish ponds).

### Spray Drift Requirements

#### Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

#### Temperature Inversion

Do not make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

#### Droplet Size

Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

#### Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airstblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

#### Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

### ARTICHOKE

#### At-Plant

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Cribrate Weevil (Grubs)	8.5	0.49	0.1	Apply as a 5 to 7 inch band (T-band) over an open furrow, or in-furrow with the seed.

#### PPI & PRE

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Cribrate Weevil Grubs	PRE 8.5	PRE 0.1	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides.
	PPI 8.5	PPI 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Cribrate Weevil Artichoke Plume Moth	8.5	0.1	Apply when pest population reaches damaging threshold and repeat as necessary to maintain control, but not more often than 15 day intervals.  <b>Application by ground:</b> Apply a full cover spray in a minimum of 75 gallons of finished spray per acre. <b>Application by air:</b> Apply specified dosage in a minimum of 10 gallons of finished spray per acre.
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply within 5 days of harvest.</li> </ul>			

**Artichoke Restrictions:**

- Do not apply more than 0.5 lb ai/A per season including at-plant, PPI, PRE and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

**BRASSICAS, Head and Stem (Crop Subgroup 5A)**

Including: Broccoli, Chinese Broccoli (gailon, white flowering broccoli), Brussels Sprouts, Cauliflower, Cavalo broccoli, Kohlrabi, Cabbage, Chinese Cabbage (napa), and Chinese Mustard Cabbage (gai choy)

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Army cutworm Armyworm species Cabbage maggot Cutworm species Grubs Root aphids Root maggot Seedcorn maggot Wireworm	3.4 – 6.8	0.2 – 0.39	0.04-0.08	Apply as a 5-7 inch band (T-band) over the open seed or transplant furrow, or in-furrow with the seed or transplant.  Cutworm and armyworm treatments may be applied as broadcast treatments to the soil surface. May be applied through transplant water at time of transplanting
<b>At Plant Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.1lb ai/A per season as an at-plant application.</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Root Aphids Root Maggots Seed corn maggot Wireworms Garden Symphylans	PRE 3.4 – 6.8	PRE 0.04 – 0.08	Capture LFR Insecticide can be tank mixed and applied with PRE labeled herbicides and fungicides for pre-transplant application.
	PPI 3.4 – 6.8	PPI 0.04 - 0.08	Incorporation of Capture LFR Insecticide should not be incorporated any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting or transplant depth.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Armyworms Cutworms Corn Earworm Crickets Cucumber Beetles Diamondback Moth Flea Beetles Ground Beetles Imported Cabbageworm Leafhoppers Loopers Saltmarsh Caterpillar Stink Bugs Thrips Tobacco Budworm Whitefly Wireworm (adults) Black burrowing bug	2.8 – 8.5	0.033 – 0.1	Thorough coverage is necessary to attain acceptable control. Make application at the onset of infestation reaching locally determined economic thresholds.  Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1 - 2 quarts of emulsified oil may be substituted for 1 - 2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.
Banks Grass Mite Carmine Mite Lygus Species Pacific Spider Mite Twospotted Spider Mite	6.8 – 8.5	0.08 – 0.1	
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make more than 5 applications after bloom.</li> <li>Do not make applications less than 7 days apart.</li> <li>Do not apply within 7 days of harvest.</li> </ul>			

**Brassica, Head and Stem (Crop Subgroup 5A) Restrictions:**

- Do not apply more than 0.5 lb ai/A per season including at-plant, PPI, PRE and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

**CILANTRO, CORIANDER**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Armyworm species Cutworm species Flea beetle larvae Wireworm	3.4 – 6.8	0.2 – 0.39	0.04 – 0.08	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band) in-furrow with the seed, or broadcast to the soil surface.
<b>At Plant Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application.</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root aphids Seed corn beetle Seed corn maggot Wireworms (PPI Only)	PRE 3.4 – 6.8	PRE 0.04 – 0.08	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides
	PPI 3.4 – 6.8	PPI 0.04 – 0.08	Capture LFR Insecticide can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.

## Foliar

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Beet Armyworm Cabbage Looper Cutworm Flea beetle Grasshoppers Leafminer Saltmarsh caterpillar Spotted Cucumber beetle Thrips Whitefly	2.8 – 8.5	0.033 - 0.1	Apply using sufficient water to obtain uniform coverage.  Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.
Two Spotted Spider Mite	6.8 – 8.5	0.08 - 0.1	
<b>Foliar Restrictions:</b>			
<ul style="list-style-type: none"> <li>Do not make applications less than 7 days apart.</li> <li>Do not apply within 3 days of harvest.</li> </ul>			

### Cilantro and Coriander Restrictions:

- Do not apply more than 0.5 lb ai/A per season including at-plant, PPI, PRE and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

## CITRUS (Crop Group 10-10)

Including: Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; unqi fruit; cultivars, varieties, and/or hybrids of these.

### General Use Directions:

When applied as directed, Capture LFR Insecticide will provide control of the following pests listed in the table below. Apply Capture LFR Insecticide by ground equipment to bare soil beneath citrus trees. Capture LFR Insecticide must be uniformly applied from the trunk to the drip line of tree. Apply in a minimum of 40 gallons of dilute spray per acre.

Greater spray volume should insure greater uniformity of coverage. A pre- and post-application irrigation may aid in the uniformity of coverage as well.

Capture LFR Insecticide protects citrus tree roots from Diaprepes and other citrus root weevil feeding by forming a barrier which provides contact activity on newly hatched larvae (neonates). As citrus root weevil eggs hatch in new foliage, neonates fall to the soil surface beneath the tree and come in contact with Capture LFR Insecticide as they attempt to burrow into the root zone. Disturbance of the soil beneath trees should be minimized.

Timing of Capture LFR Insecticide applications is critical. Current information suggests that peak emergence of adult Diaprepes Weevil varies by citrus growing region and these emergence peaks can be dramatically affected by environmental factors, such as soil moisture. Typically, two peaks are observed for Diaprepes, first in spring then late summer or early fall. Southern Blue- Green and Blue-Green Citrus Weevils and Fuller Rose Beetle typically exhibit a single emergence peak in the spring. Brown and Little Leaf Notchers typically exhibit three emergence peaks, spring, summer and fall. Since emergence varies seasonally and by location, timing of Capture LFR Insecticide application can be accurately forecast by observing adults. Adults are most active early morning and late afternoon; numbers can be estimated by trapping throughout spring and summer (emergence periods). Egg laying will occur for 8 to 10 weeks following adult emergence from the soil; larval invasion of the soil will begin 2 to 3 weeks following adult emergence. It is critical to have the Capture LFR Insecticide soil barrier in place prior to drop of the neonates.

Capture LFR Insecticide is one of several effective tools in an integrated pest management program for Citrus Root Weevils. Application of Capture LFR Insecticide should be used in conjunction with good cultural practices, biological control of larvae and foliar control of adults. Consult local university extension personnel for current information to protect citrus trees from Citrus Root Weevils and other pests.

Apply to individual citrus resets, when not in solid planted rows, using hand-gun or shielded sprayer.

Peak emergence of Diaprepes root weevil generally occurs in the spring. Depending on weather conditions, a minor emergence of Diaprepes root weevil may also occur in the fall.

If the citrus grove to be treated is in an area where weather conditions are conducive to primary emergence occurring in the spring, 42.5 fluid ounces formulated product should be used to obtain the longest residual management of Diaprepes root weevil. If the citrus grove to be treated is in an area where weather conditions will promote more than one peak of pest emergence, 21.25 fluid ounces formulated product can be

applied early season and 21.25 fluid ounces formulated product can be applied later in the season.

## BARE SOIL SURFACE UNDER DRIP LINE

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Asian cockroach Fire ants ( <i>Solenopsis species</i> )	8.5 - 21.25	0.1 - 0.25	Apply the specified dosage in a minimum of 40 gallons of finished spray per acre.
Blue Green Citrus Root Weevil Brown Leaf Notcher Diaprepes Root Weevil Little Leaf Notcher Southern Blue Green Citrus Root Weevil	21.25- 42.5	0.25 - 0.5	
<b>Restrictions:</b>			
<ul style="list-style-type: none"> <li>Do not allow any application of Capture LFR Insecticide to contact fruit or foliage.</li> <li>Do not apply more than a total of 42.5 fluid ounces of formulated product (0.5 pound active ingredient) per acre per year.</li> <li>Do not apply by air.</li> </ul>			

## CORN

**Field Corn (Grain and Silage), Popcorn, Field Corn Grown for Seed, Sweet Corn, Sweet Corn Grown for Seed**

### Use Directions:

Heavy Corn Rootworm Pressure Management Program: In areas where large corn rootworm populations are present, a multi-approach system may be needed for optimal pest management. However, if the population level is not known and if a corn rootworm adult scouting program along with threshold adult control measures were not completed during the previous growing season, then utilize a maximum dosage seed treatment program or genetically modified corn rootworm resistant hybrid in addition to Capture LFR Insecticide.

### At-Plant

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Corn Rootworm larvae (Northern, Southern and Western)	6.8 - 17.0	0.39 - 0.98	0.08 - 0.2	Apply as a 5 to 7 inch band (T-band) over an open furrow, or in-furrow with the seed. For Army cutworm, Stalkborer, Cutworm species, True armyworm or Armyworm species, apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), in-furrow with the seed, or broadcast to the soil surface.
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root aphids Seed corn beetle Seed corn maggot Stalkborer Sugarcane beetle True armyworm Wireworm	3.4 - 13.6	0.2 - 0.78	0.04 - 0.16	
<b>At Plant Restrictions:</b>				
<ul style="list-style-type: none"> <li>Do not apply more than 0.2 pound active per acre per season as an at-plant application.</li> <li>For field corn - Do not apply more than 0.3 lb ai/A per season including ppi, at-plant, pre-emergence, and foliar applications of other bifenthrin products (such as Brigade 2EC).</li> <li>For sweet corn - Do not apply more than 0.2 lb ai/A total per season including ppi, at-plant, pre-emergence, and foliar applications of other bifenthrin products (such as Brigade 2EC).</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Armyworm species Black Cutworm Grape colaspis Seed corn beetle Seedcorn Maggot White Grub Wireworm	<b>PPI</b> 4 to 5.3	<b>PPI</b> 0.047 to 0.062	Capture LFR Insecticide can be tank mixed and applied with PPI herbicides. Incorporation of Capture should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.
Armyworm species Black Cutworm Seed corn beetle Stalkborer	<b>PRE</b> 3.4	<b>PRE</b> 0.04	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Army Cutworm Beet Armyworm Cereal Leaf Beetle Chinch Bug Common Stalk Borer Corn Earworm <sup>1</sup> Corn Rootworm Adults Cucumber Beetle Adult Cutworm species European Corn Borer <sup>2</sup> Fall Armyworm Flea Beetle Grasshoppers Greenbug Japanese Beetle Adult Sap Beetle Southern Armyworm Southern Corn Leaf Beetle Southwestern Corn Borer <sup>2</sup> Stinkbugs Tarnished Plant Bug True Armyworm or Armyworm species Webworms Western Bean Cutworm Yellowstriped Armyworm	2.8 – 8.5	0.033 - 0.1	Apply in a minimum of 2-5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft, use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal.  When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray.  Thorough coverage is essential to achieve control.  <b>1To Control Ear Attacking Pests:</b> Apply Capture LFR Insecticide just before silking and repeat as necessary to maintain control.  <b>2Southwestern Corn Borer, European Corn Borer:</b> Make application for corn borer control with initial application at or shortly before egg hatch.  <b>For Control of Other Insect Pests:</b> Apply when pests first appear and repeat as necessary.  <b>3For Control of Mites:</b> Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.  <b>For Twospotted Spider Mite and Carmine Mite control,</b> apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.  Higher rates will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb. active per acre in tank mixture has demonstrated good control under these conditions.  <b>For Mite Control in Texas, New Mexico, Oklahoma, and Arizona:</b> Apply in a minimum of 5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment
Banks Grass Mite <sup>3</sup> Carmine Mite <sup>3</sup> Twospotted Spider Mite <sup>3</sup>	6.8 – 8.5	0.08 - 0.1	

**For field corn**

- Do not apply within 30 days of harvest for field corn (grain and silage), popcorn, field corn grown for seed.
- Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application for field corn (grain and silage), popcorn, field corn grown for seed.

**For sweet corn**

- Do not apply within 1 day of harvest for sweet corn or sweet corn grown for seed.
- Do not graze livestock in treated areas or cut treated crops for feed within 1 day of the last application for sweet corn or sweet corn grown for seed.

Use of ultra low volume (ULV) application on corn is prohibited.

Do not make aerial or ground applications to corn if heavy rainfall is imminent.

**Corn Restrictions:**

• **For field corn**

- Do not apply more than 0.3 lb ai/A total per season including ppi, at-plant, pre-emergence, and foliar applications of Capture LFR Insecticide and other bifenthrin products.

• **For sweet corn**

- Do not apply more than 0.2 lb ai/A total per season including ppi, at-plant, pre-emergence, and foliar applications of Capture LFR Insecticide and other bifenthrin products.

**CUCURBITS (Crop Group 9)**

Including: **Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd, edible *Lagenaria* species (includes hyotan, cucuzza), *Luffa* species (includes hechima, Chinese okra), *Momordica* species (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin (*Cucurbita* species), Squash, summer (includes crook-neck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Squash, winter (includes butternut squash, calabaza, hubbard squash (*C. mixta*; *C. pepo*) includes acorn squash, spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus* spp.).**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Cucumber beetle larvae	6.8 - 8.5	0.39 - 0.49	0.08 - 0.1	Apply as a 5 to 7 inch band (T-band) over an open furrow, or in-furrow with the seed. May be applied through transplant water at time of transplanting.
Army cutworm Armyworm species Cutworm species Flea beetle larvae Grubs True Armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	To control cucumber beetle larvae, apply as a 5 to 7 inch band over an open furrow (T-band), or in-furrow with the seed.  To control wireworm, grubs, and flea beetle larvae, apply as a 5 to 7 inch band over an open furrow (T-band), or in-furrow with the seed or transplant  To control army cutworm, cutworm species, true armyworm and armyworm species, apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), in-furrow with the seed, broadcast to the soil surface or banded over the row.

**At Plant Restrictions:**

- Do not apply more than 0.1 lb ai/A per season as an at-plant application

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Seed Corn Maggot Wireworms Army cutworm Armyworm species Cutworm species Flea beetle larvae Grubs True Armyworm True armyworm	<b>PRE</b> 6.8 – 8.5	<b>PRE</b> 0.08 – 0.1	Capture LFR Insecticide can be tank mixed and applied with PRE pesticides.  Apply through drip of drip tape. Apply when soil is moist towards the end of the irrigation run.
	<b>PPI</b> 6.8 – 8.5	<b>PPI</b> 0.08 - 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI labeled pesticides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth. Incorporation depth should be close to the intended depth  Apply through drip or Drip tape. Apply when soil is moist towards the end of the irrigation run

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Armyworms Cabbage Looper Corn Earworm Cucumber beetles Cutworms Grasshopper Leafhoppers Melonworm Pickleworm Plant Bug Rindworm Squash Bugs Squash Vine Borer Stink Bugs Tobacco Budworm	3.4 – 8.5	0.04 – 0.1	Thorough coverage is necessary to attain acceptable control. Make application at the onset of infestation reaching locally determined economic thresholds  Apply in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment When applying by air 1 2 quarts of emulsified oil may be substituted for 1 – 2 quarts of water in the finished spray Thorough coverage is essential to achieve control.
Carmine Mite Lygus species Mite Twospotted Spider Mite Whitefly	6.8 – 8.5	0.08 – 0.1	
<b>Foliar Restrictions:</b>			
<ul style="list-style-type: none"> <li>Do not make more than two applications after bloom.</li> <li>Do not make applications less than 7 days apart.</li> <li>Do not apply within 3 days of harvest</li> </ul>			

**Cucurbits (Crop Group 9) Restrictions:**

- Do not apply more than 0.3 lb ai/A per season including **At-plant, PRE & PPI, and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.**

**DRIED BEANS AND PEAS (Crop Subgroup 6C)**  
Including: Dried cultivars of: Bean (Lupinus); Bean (Phaseolus), Field bean, Kidney bean, Lima bean (dry), Navy bean, Pinto bean, Tepary bean; Bean (Vigna), Adzuki bean, Blackeyed pea, Catjang, Cowpea, Crowder pea, Moth bean, Mung bean, Rice bean, Southern pea, Urd bean; Broad bean (dry), Chickpea, Guar, Lablab bean, Lentil; Pea (Pisum), Field pea, Pigeon pea., purple hulled peas

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Corn rootworm larvae	6.8 – 8.5	0.39 – 0.49	0.08 – 0.1	Apply as a 5 to 7 inch band over the row on the soil surface, 5 to 7 inch band (T-band) over an open furrow, or in-furrow with the seed.
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root maggot True armyworm Wireworm	3.4 - 6.8	0.2 – 0.39	0.04 – 0.08	Apply broadcast to the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species.
<b>At Plant Restrictions:</b>				
<ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application.</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root maggot True armyworm Wireworm (PPI only)	<b>PRE</b> 6.8 – 8.5	<b>PRE</b> 0.08 – 0.1	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides. Apply in a minimum of 10 gallons per acre.
	<b>PPI</b> 6.8 – 8.5	<b>PPI</b> 0.08 – 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth. Apply in a minimum of 10 gallons per acre.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aster Leafhopper Flea Beetle Grasshoppers Leafhoppers	2.1 – 8.5	0.025 – 0.1	Apply in a minimum of 2 gallons finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. Thorough coverage is essential to achieve control.
Alfalfa Caterpillar Aphids Bean Leaf Beetle Beet Armyworm Cloverworm Corn Earworm Corn Rootworm (Adult) Cucumber Beetles Cutworms European Corn Borer Fall Armyworm Grasshoppers Imported cabbage-worm Japanese beetle (Adult) Leafminer Loopers Mexican Bean Beetle Pea Leaf Weevil Pea Weevil Plant Bug Saltmarsh caterpillar Sap Beetle Southern Armyworm Stink Bugs Tarnished Plant Bug Thrips Twospotted Spider Mite Tobacco budworm Webworms Western Bean Cutworm Whitefly Yellowstriped Armyworm	2.8 – 8.5	0.033 – 0.1	When applying by air 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control
Banks Grass Mite Carmine Mite Lygus Species	6.8 - 8.5	0.08 – 0.1	
<b>Foliar Restrictions:</b>			
<ul style="list-style-type: none"> <li>Do not apply within 14 days of harvest</li> <li>Do not make applications less than 7 days apart</li> </ul>			

**Dried Beans and Peas (Crop Subgroup 6C) Restrictions:**

- Do not apply more than 0.2 lb ai/A to peas, or 0.3lb ai/A to beans per season including **At-Plant, PRE & PPI, and Foliar applications of Capture LFR Insecticide and other bifenthrin containing products.**

**EGGPLANT**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Grubs Root maggot True Armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), or in-furrow with the transplant or seed. Apply broadcast to the soil surface for control of Army Cutworm, Cutworm species, True Armyworm or Armyworm species
<b>At Plant Restrictions:</b>				
<ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application.</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Grubs Root maggot True armyworm Wireworm	<b>PRE</b> 8.5	<b>PRE</b> 0.1	Capture LFR Insecticide can be tank mixed and applied with PRE pesticides.  <b>Post Plant Soil Applied:</b> Apply through drip or Drip tape. Apply when soil is moist towards the end of the irrigation run.
	<b>PPI</b> 3.4 – 8.5	<b>PPI</b> 0.04 – 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI labeled pesticides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth. Incorporation depth should be close to the intended depth.  <b>Post Plant Soil Applied:</b> Apply through drip or Drip tape. Apply when soil is moist towards the end of the irrigation run.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Armyworms (Including Beet) Armyworm Fall Armyworm Southern Cabbage Looper Colorado Potato Beetle Corn Earworm Cucumber Beetle Cutworms European Corn Borer Flea Beetle Leafminers Loopers Pepper weevil Plant Bug Stink Bug Thrips Tomato Hornworm Tomato Pinworm Vegetable Leafminer Whitefly Yellowstriped Armyworm	2.8 – 8.5	0.033 – 0.1	Thorough coverage is necessary to attain acceptable control. Make application at the onset of infestation reaching locally determined economic thresholds.  Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.  When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray.  Thorough coverage is essential to achieve control.
Banks Grass Mite Broad Mite Carmine Mite Lygus species Pacific Spider Mite Twospotted Spider Mite	6.8 – 8.5	0.08 – 0.1	

**Foliar Restrictions:**

- Do not make applications less than 7 days apart.
- Do not apply within 7 days of harvest.

**Eqplant Restrictions:**

- Do not apply more than 0.2 lb ai/A per season including **At-Plant, PRE & PPI, and Foliar applications of Capture LFR Insecticide and other bifenthrin containing products.**

**HEAD LETTUCE**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Rootworm larvae	6.8 - 8.5	0.39 - 0.49	0.08 - 0.1	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), or in-furrow with the seed. Apply broadcast to the soil surface for control of Army cutworm, Cutworm species, True armyworm, armyworm species, or bulb mites.
Army cutworm Armyworm species Bulb mites Cutworm species Grubs Lettuce root aphid Root maggot True armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	

**At Plant Restrictions:**

- Do not apply more than 0.1 lb ai/A per season as an at-plant application.

**PPI**

PEST	USE RATES	
	Fluid oz/acre	Pound ai/acre
Lettuce Root Aphid Garden Symphians	6.8-8.5	0.08 – 0.1

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Armyworms Corn earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Loopers Salt Marsh Caterpillar Stink Bug species Tobacco Budworm Whitefly	2.8 – 8.5	0.033 – 0.1	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.
Carmine Mite Lygus Species Twospotted Spider Mite	6.8 – 8.5	0.08 – 0.1	

**Foliar Restrictions:**

- Do not make applications less than 7 days apart.
- Do not apply within 7 days of harvest.

**Head Lettuce Restrictions:**

- Do not apply more than 0.5 lb ai/A per season **At-Plant, PRE & PPI, and Foliar applications of Capture LFR Insecticide and other bifenthrin containing products.**

**LEAFY BRASSICAS (Crop Subgroup 5B), TURNIP GREENS**

Including: Broccoli Raab, Bok Choy, Collards, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens, Turnip Greens

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Rootworm larvae	6.8 – 8.5	0.39 – 0.49	0.08 – 0.1	Apply as a 5-7 inch band over the row on the soil surface, a 5-7 inch band over the open furrow (T-band), or in-furrow with the seed or transplant. May be applied through transplant water at time of transplanting.
Army cutworm Armyworm species Cutworm species Grubs Lettuce root aphid Root maggot True armyworm Wireworm	3.4 – 6.8	0.2 – 0.39	0.04 – 0.08	Apply broadcast over the soil surface for control of Army cutworm, Cutworm species, True armyworm or armyworm species.

**At Plant Restrictions:**

- Do not apply more than 0.1 lb ai/A per season as an at-plant application.

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Flea beetle larvae Grubs Lettuce root aphid Root Maggots True armyworm Wireworms	<b>PRE</b> 3.4 – 6.8	<b>PRE</b> 0.04 – 0.08	Capture LFR Insecticide can be tank mixed and applied with PRE pesticides.  <b>Post Plant Soil Applied:</b> Apply thorough drip or drip tape. Apply when soil is moist towards the end of the irrigation run.
	<b>PPI</b> 3.4 – 6.8	<b>PPI</b> 0.04 – 0.08	Capture LFR Insecticide can be tank mixed and applied with PPI labeled pesticides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth. Incorporation depth should be close to the intended depth.  <b>Post Plant Soil Applied:</b> Apply thorough drip or drip tape. Apply when soil is moist towards the end of the irrigation run.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Armyworms Corn Earworm Cutworms Crickets Cucumber Beetles Diamondback Moth Flea Beetles Grasshoppers Ground Beetles Imported Cabbageworm Japanese Beetle (adult) Leafhoppers Loopers Saltmarsh Caterpillar Stink Bugs Tobacco Budworm Thrips Whitefly Wireworm (adults)	2.8 – 8.5	0.033 – 0.1	Thorough coverage is necessary to attain acceptable control. Make application at the onset of infestation reaching locally determined economic thresholds.  Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.
Banks Grass Mite Carmine Mite Lygus species Pacific Spider Mite Twospotted Spider Mite	6.8 – 8.5	0.08 – 0.1	
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make applications less than 7 days apart.</li> <li>Do not apply within 7 days of harvest.</li> </ul>			

**Leafy Brassica (Crop Subgroup 5B) and Turnip Greens Restrictions:**

- Do not apply more than 0.4 lb ai/A per season including at-plant, PPI, PRE, and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

**OKRA**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Armyworm Cutworm species Flea beetle larvae Grape colaspis Root maggot Wireworm White Grub	3.4 – 6.8	0.2 – 0.39	0.04 – 0.08	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), in-furrow with the seed, or broadcast to the soil surface.
<b>At Plant Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application.</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Cutworm species	<b>PRE</b> 3.4 – 8.5	<b>PRE</b> 0.04 – 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI or PRE herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.
Cutworm species Flea beetle larvae Grape colaspis Root maggot Wireworm White Grub	<b>PPI</b> 3.4-8.5	<b>PPI</b> 0.04 to 0.1	

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Armyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Japanese Beetle (Adult) Leafminers Loopers Stink bugs Thrips Whitefly	2.8 – 8.5	0.033 - 0.1	Apply as needed using sufficient water to obtain uniform coverage.  Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.
Broad Mite Carmine Mite Lygus species Twospotted Spider Mite	6.8 – 8.5	0.08 - 0.1	
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make applications less than 7 days apart.</li> <li>Do not apply within 7 days of harvest.</li> </ul>			

**Okra Restrictions:**

- Do not apply more than 0.2 lb ai/A per season including At-Plant, PRE & PPI, and Foliar applications of Capture LFR Insecticide and other bifenthrin products.

**PEPPERS (BELL and NON-BELL) & PEPINO**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Flea beetle larvae Grubs Pepper maggot Root aphid Root maggot Stalk borer True armyworm Wireworm	3.4 –6.8	0.2 – 0.39	0.04-0.08	Apply as a 5-7 inch band (T-band) over the open seed or transplant furrow, or in-furrow with the seed or transplant.  Cutworm and armyworm treatments may be applied as broadcast treatments to the soil surface. May be applied through transplant water at time of transplanting
<b>At Plant Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.1lb ai/A per season as an at-plant application.</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Flea beetle larvae Grubs True Armyworm Wireworm	<b>PRE</b> 8.5	<b>PRE</b> 0.1	Capture LFR Insecticide can be tank mixed and applied with PRE pesticides  <b>Post Plant Soil Applied:</b> Apply through drip or drip tape. Apply when soil is moist towards the end of the irrigation run.
	<b>PPI</b> 3.4 – 8.5	<b>PPI</b> 0.04 – 0.1	

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Armyworms (Including Beet) Armyworm Fall Armyworm Southern Cabbage Looper Colorado Potato Beetle Corn Earworm Cucumber Beetle Cutworms European Corn Borer Flea Beetle Leafminers Loopers Pepper weevil Plant Bug Stink Bug Thrips Tomato Hornworm Tomato Pinworm Vegetable Leafminer Whitefly Yellowstriped Armyworm	2.8 – 8.5	0.033 - 0.1	Thorough coverage is necessary to attain acceptable control. Make application at the onset of infestation reaching locally determined economic thresholds.  Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.  When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.
Broad Mite Carmine Mite Lygus species Pacific Spider Mite Twospotted Spider Mite	6.8 – 8.5	0.08 - 0.1	

**Foliar Restrictions:**

- Do not make applications less than 7 days apart.
- Do not apply within 7 days of harvest.

**Pepper (Bell and Non-Bell) Restrictions:**

- Do not apply more than 0.2 lb ai/A per season including at-plant, PPI, PRE and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

**SOD FARMS**

**At-Plant**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Cutworms <sup>1</sup> White Grub Wireworm Crickets Earwigs Ants Chinch Bugs <sup>5</sup> Imported Fire Ants <sup>8</sup>	8.5	0.1	Apply as a 5 to 7 inch band (T-band) over an open furrow, or in-furrow with the seed.

**PRE & PPI**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Ants Chinch Bugs <sup>5</sup> Crickets Cutworms <sup>1</sup> Earwigs Imported Fire Ants <sup>8</sup> White Grub Wireworm	<b>PRE</b> 8.5	<b>PRE</b> 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI and PRE herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.
	<b>PPI</b> 8.5	<b>PPI</b> 0.1	

**Foliar**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/ 1000 sq. ft.	Pound ai /acre	
Armyworms <sup>1</sup> Cutworms <sup>1</sup> Sod Webworm <sup>1</sup>	2.8 – 4.35	0.066 - 0.1	0.033 - 0.05	Apply as a broadcast treatment. Use higher volumes up to 10 gallons of carrier per 1000 square feet to get uniform coverage when treating dense grass foliage.  Irrigation to treated area within a few hours following application can improve efficacy to sub-surface pests including mole crickets.  The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, Capture LFR Insecticide may be applied at up to 0.4 fluid oz. per 1000 square feet to control each of the pests listed in this table. The higher application rates should be used when maximum residual control is desired or heavy pest populations occur.
Annual Bluegrass Weevil (Hyperodes) (Adult) <sup>2</sup> Banks Grass Mite <sup>6</sup> Billbugs (Adult) <sup>3</sup> Black Turfgrass Ataenius (Adult) <sup>4</sup> Crickets Earwigs Fleas (Adult) Grasshoppers Mealybugs Mites <sup>5</sup>	4.35 - 8.7	0.1 - 0.2	0.05 - 0.1	
Ants Chinch Bugs <sup>5</sup> Fleas (Larvae) <sup>7</sup> Imported Fire Ants <sup>8</sup> Japanese Beetle (Adult) Mole Cricket (Adult) <sup>9</sup> Mole Cricket (Nymph) <sup>10</sup> Ticks <sup>11</sup>	8.7 - 17.42	0.2 - 0.4	0.1 - 0.2	

<sup>1</sup>**Armyworms, Cutworms and Sod Webworms:** To ensure optimum control, delay watering (irrigation) or mowing for 24 hours after application. If the grass area is being maintained at a mowing height of greater than 1 inch, then higher application rates (up to 0.4 fluid oz. per 1000 square feet) may be required during periods of high pest pressure.

<sup>2</sup>**Annual Bluegrass Weevil (Hyperodes) adults:** Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when *Forsythia* is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

<sup>3</sup>**Billbug adults:** Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

<sup>4</sup>**Black Turfgrass Ataenius adults:** Applications should be made during May and July to control the first and second generation of black turfgrass ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of *Vanhoutte spiraea* (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of *Rose of Sharon* (*Hibiscus syriacus*).

<sup>5</sup>**Chinch Bugs:** Chinch Bugs infest the base of grass plants and are often found in the thatch layer. Irrigation of the grass area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located. Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained. Chinch Bugs can be one of the most difficult pests to control in grasses and the higher application rates (up to 0.4 fluid oz. per 1000 square feet) may be required to control populations that contain both nymphs and adults during the middle of the summer.

<sup>6</sup>**Mites:** To ensure optimal control of eriophyid mites, apply in combination with the labeled application rate of a surfactant. A second application, five to seven days after the first, may be necessary to achieve acceptable control.

<sup>7</sup>**Flea larvae:** Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Use a higher volume application when treating these areas to ensure penetration of the insecticide into the soil. Note: if the lawn area is being treated with this product at 0.1 fluid oz. per 1000 square feet for adult flea control, then the larval application rate may be achieved by increasing the application volume two- to four-fold.

<sup>8</sup>**Imported Fire Ants:** Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound drenches that will control existing colonies. If the soil is not moist, then it is important to irrigate before application or use a high volume application. Broadcast treatments should apply 0.4 fluid oz. per 1,000 square feet. Mounds should be treated by diluting 0.05 fluid oz of Capture LFR Insecticide per gallon of water and applying 1 to 2 gallons of finished spray per mound. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. A four foot diameter circle around the mound should also be treated. For best results, apply in cool weather (65 - 80°F) or in early morning or late evening hours.

<sup>9</sup>**Mole Cricket adults:** Achieving acceptable control of adult mole crick-

ets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Grass areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

**<sup>10</sup>Mole Cricket nymphs:** Grass areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

**<sup>11</sup>Ticks (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted fever):** Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher spray volumes when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed activity. Repeat application must be limited to no more than once per seven days.

**Deer ticks (*Ixodes sp.*)** have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter.

**American dog ticks** may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

**Sod Farm Restrictions:**

- In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
- In New York State, make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

**SOYBEANS**

**At-Plant**

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Rootworm larvae	6.8 - 8.5	0.39 - 0.49	0.08 - 0.1	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), or in-furrow with the seed. Apply broadcast over the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species.
Army cutworm Armyworm species Bean Leaf Beetle Larvae Cutworm species Grape colaspis Grubs Root maggot Seed corn beetle Seedcorn maggot True armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	
<b>At Plant Restrictions:</b>				
<ul style="list-style-type: none"> <li>• Do not apply more than 0.1 lb ai/A per season as an at-plant application</li> </ul>				

**PPI & PRE**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Armyworm species Black Cutworm	<b>PRE</b> 3.4	<b>PRE</b> 0.04	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides.
Armyworm species Bean leaf Beetle Larvae Black Cutworm Seed corn beetle Seed corn Maggot Stalkborer White Grub Wireworm	<b>PPI</b> 4 to 5.3	<b>PPI</b> 0.047 to 0.062	Capture LFR Insecticide can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Alfalfa Caterpillar Aphids Armyworms Bean Leaf Beetle Blister Beetle species Corn Earworm Corn Rootworm Adult Cowpea Curculio Cucumber Beetle Adult Cutworms Dectes Stem Borer European Corn Borer False Cinch Bug Flea Beetle Grasshoppers Green cloverworm Hornworms Imported Cabbageworm Japanese Beetle Adult Leaf Skeletonizer species Leafhoppers Leafminers Adults Lesser Cornstalk Borer Loopers Kudzu Bug Mexican Bean Beetle Painted Lady (Thistle) Caterpillar Pea Leaf Weevil Saltmarsh Caterpillar Seedcorn Maggot Adult Silverspotted Skipper Spittlebug Stink Bug Three Cornered Alfalfa Hopper Thrips Tobacco Budworm Velvetbean Caterpillar Webworm Woollybear Caterpillar	2.8 - 8.5	0.033 - 0.1	Apply in a minimum of 10 gallons per acre with ground equipment or 2 gallon per acre by aircraft.  Pyrethroid resistance is common for Beet Armyworm and Tobacco Budworm. Please consult your local or state agricultural authority to determine if resistant pest populations are in your area. If so, refer to the resistance management statement in the DIRECTIONS FOR USE section of this label.
Lygus species Whitely Twospotted spider mite	6.8 - 8.5	0.08 - 0.1	
<b>Foliar Restrictions:</b>			
<ul style="list-style-type: none"> <li>• Do not make applications less than 30 days apart.</li> <li>• Do not apply within 18 days of harvest.</li> </ul>			

**Soybean Restrictions:**

- Do not apply more than 0.3 lb ai/A per season including **At-Plant, PRE & PPI, and Foliar applications of this and other bifenthrin products.**

# SPINACH

## At-Plant

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Rootworm larvae Garden Symphylans	6.8 - 8.5	0.39 - 0.49	0.08 - 0.1	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), or in-furrow with the seed. Apply broadcast to the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species.
Army cutworm Armyworm species Cutworm species Grubs Root maggot Seedcorn maggot True armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	
<b>At Plant Restrictions:</b>				
<ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application</li> </ul>				

## PPI

PEST	USE RATES	
	Fluid oz/acre	Pound ai/acre
Seed Corn Maggot Wireworms Garden Symphylans	<b>PPI</b> 3.4 - 6.8	<b>PPI</b> 0.04 - 0.08

## Foliar

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Armyworms Colorado Potato Beetle Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Leafminers Loopers Pepper Weevil Tomato Hornworm Tomato Pinworm Thrips Whitefly	2.8 - 8.5	0.033 to 0.1	For control of whiteflies, apply foliar treatments of Capture LFR Insecticide by ground or air at rates of up to 0.1 lb active per acre at minimum 7 day intervals up to a maximum of 4 applications. Do not apply within 40 days of harvest.
Banks Grass Mite Broad Mite Carmine Mite Fire Ants Lygus species Pacific Spider Mite Twospotted spider mite	6.8 - 8.5	0.08 to 0.1	For control of fire ants apply Capture LFR Insecticide to the soil (at planting) or as a foliar treatment by ground or air at rates of up to 0.1 lb active per acre at minimum 7 day intervals up to a maximum of 4 applications.  Apply the specified dosage in 5-50 gallons of finished spray per acre by air or 10-50 gallons finished spray per acre by ground.
<b>Foliar Restrictions:</b>			
<ul style="list-style-type: none"> <li>Do not make applications less than 7 days apart.</li> <li>Do not apply within 40 days of harvest.</li> </ul>			

## Spinach Restrictions:

- Do not apply more than 0.4 lb ai/A per season including At-Plant, PRE & PPI, and Foliar applications of this and other bifenthrin products.

# SUCCULENT PEAS AND BEANS (Crop Subgroups 6A and 6B)

Pea (*Pisum* spp.): Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea, Pigeon pea; Bean (*Phaseolus* spp.): Broadbean (succulent), Lima bean (green), Runner bean, Snap bean, Wax bean; Bean, *Vigna* spp.): Asparagus bean, Blackeyed pea, Chinese longbean, Cowpea, Moth bean, Southern pea, Yardlong bean, Jackbean, Soybean (immature seed), Sword bean

## At-Plant

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Rootworm larvae	6.8 - 8.5	0.39 - 0.49	0.08 - 0.1	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), or in-furrow with the seed. Apply broadcast over the soil surface for control of Army cutworm, Cutworm species, True armyworm, or Armyworm species.
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root maggot Seedcorn maggot True armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	
<b>At Plant Restrictions:</b>				
<ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application</li> </ul>				

## PPI & PRE

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Grape colaspis Grubs Root maggot True armyworm Wireworm (PPI only)	<b>PRE</b> 6.8 - 8.5	<b>PRE</b> 0.08 - 0.1	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides.  <b>Post Plant Soil Applied:</b> Apply through drip or Drip Tape. Apply when soil is moist towards the end of the irrigation run.
	<b>PPI</b> 6.8 - 8.5	<b>PPI</b> 0.08 - 0.1	Capture LFR Insecticide can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth. Apply in a minimum of 10 gallons per acre.  <b>Post Plant Soil Applied:</b> Apply through drip or Drip tape. Apply when soil is moist towards the end of the irrigation run.

## Foliar

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aster Flea Beetle Leafhopper	2.1 – 8.5	0.025 - 0.1	Apply in a minimum of 2 gallons finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.
Adult Sap Beetle Alfalfa Caterpillar Aphids Armyworm, Beet Armyworm, Fall Armyworm, Southern Armyworm, Yellowstriped Bean Leaf Beetle Cloverworm Corn earworm Corn Rootworm Adult Cucumber Beetle Cutworms European Corn Borer Grasshoppers Japanese Beetle Loopers Pea Leaf Weevil Pea Weevil Plant Bug Stink Bugs Tarnished Plant Bug Thrips Webworms Western Bean Cutworm Whitefly	2.8 – 8.5	0.033 - 0.1	When applying by air, 1 – 2 quarts of emulsified oil may be substituted for 1 – 2 quarts of water in the finished spray.  Thorough coverage is essential to achieve control.  Make application at the onset of infestation reaching locally determined economic threshold.
Banks Grass Mite Carmine Mite Lygus species Twospotted spider mite	6.8 – 8.5	0.08 to 0.1	
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply within 3 days of harvest.</li> <li>Do not make applications less than 3 days apart.</li> </ul>			

### Succulent Peas and Beans (Crop Subgroups 6A and 6B) Restrictions:

- Do not apply more than 0.2 lb ai/A per season including At-Plant, PRE & PPI, and Foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

## TOBACCO

### Pre-Transplant & At-Transplant

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Armyworm species Cutworm species Flea beetle larvae Mole cricket Stalkborer White grubs Wireworm	3.4 – 8.5	0.2 - 0.49	0.04 - 0.1	<b>Pre-transplant soil applications:</b> Use of suitable equipment to incorporate into top 4" of the soil is required to control below ground pests.  <b>At-transplant water treatment application:</b> Apply 0.0625 to 0.1 pound ai/A in a water treatment application volume of 10 to 200 gal/A.  May be tank mixed with Command, Spartan, and other herbicides approved for tobacco use.
<b>Pre-Transplant &amp; At-Transplant Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply later than layby.</li> </ul>				

## Foliar

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphid species* Armyworm species Chinch Bugs Cutworm species Flea Beetle (Adults) Grasshoppers Green Bugs Japanese Beetles Stink Bugs Tarnished plant bugs Thrips Whiteflies	3.4 – 8.5	0.04 - 0.1	Apply 0.04 to 0.10 lb ai/A per foliar application up to, and including, layby in a minimum of 10 gal/A.  May be tank mixed with Command, Spartan and other herbicides approved for tobacco use.  *See resistance statement under "Directions for Use" section.
Hornworm Tobacco Budworm	6.8 - 8.5	0.08- 0.1	
Spider mites Lygus species	8.5	0.1	
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make more than 2 foliar applications per season.</li> <li>Do not apply later than layby.</li> </ul>			

### Tobacco Restrictions:

- Do not apply more than 0.2 lb ai/A per season including at-plant, PRE, PPI and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

## TOMATOES, TOMATILLOS, & GROUND-CHERRIES

### At-Plant

PEST	USE RATES			DIRECTIONS
	Fluid oz/acre	Fluid oz/1000 Linear ft.	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Flea beetle larvae Grubs Root maggot Stalkborer True armyworm Wireworm	3.4 - 6.8	0.2 - 0.39	0.04 - 0.08	Apply as a 5 to 7 inch band over the row on the soil surface, a 5 to 7 inch band over the open furrow (T-band), or in-furrow with the transport or seed. May be applied through transplant water at time of transplanting. Apply broadcast to the soil surface for control of Army cutworm, Cutworm species, True armyworm, Armyworm species, or Stalkborer.
<b>At Plant Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.1 lb ai/A per season as an at-plant application.</li> </ul>				

### PPI & PRE

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Army cutworm Armyworm species Cutworm species Flea beetle larvae Garden symphylans Grubs True Armyworm Wireworm	<b>PRE</b> 6.8	<b>PRE</b> 0.08	Capture LFR Insecticide can be tank mixed and applied with PRE herbicides.  <b>Post Plant Soil Applied:</b> Apply through drip or Drip Tape. Apply when soil is moist towards the end of the irrigation run.
	<b>PPI</b> 3.4 – 6.8	<b>PPI</b> 0.04 – 0.08	Capture LFR Insecticide can be tank mixed and applied with PPI labeled herbicides. Incorporation of Capture LFR Insecticide should not be any deeper than the intended planting depth. Incorporation depth should be close to the intended depth  <b>Post Plant Soil Applied:</b> Apply through drip or Drip Tape. Apply when soil is moist towards the end of the irrigation run.

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Aphids Armyworms (including Beet) Armyworm, Fall Armyworm, Southern Bean Leaf Beetle Cabbageworm Carmine Mite Cloverworm Corn earworm Corn Rootworm Cucumber Beetles Cutworms Diamondback Moth European Corn Borer Flea Beetles Flea Hopper Grasshopper Japanese Beetle (Adult) Leafhoppers Loopers Lygus species Melonworm Pea Weevil Pea Leaf Weevil Pickleworm Plant Bug Rindworm Salt Marsh Caterpillar Sap Beetle Seedpod Weevil Squash Bugs Stink bug species Tobacco Budworm Tarnished Plant Bug Thrips Whitefly Yellowstriped Armyworm	2.8 - 6.8	0.033 to 0.08	Thorough coverage is necessary to attain acceptable control. Make application at the onset of infestation reaching locally determined economic levels.  Apply in water. Apply the specified dosage in 5 to 50 gallons of finished spray per acre by air or 10 to 50 gallons of finished spray per acre by ground. Thorough coverage is essential to achieve control.
Twospotted spider mite	6.8 – 8.5	0.08 - 0.1	
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make applications less than 10 days apart.</li> <li>Do not apply within 1 day of harvest.</li> </ul>			

**Tomato, Tomatillo, and Groundcherry Restrictions:**

- Do not apply more than 0.40 lb ai/A per season including at-plant, PRE, PPI and foliar applications of Capture LFR insecticide and other bifenthrin containing products.

**TUBEROUS AND CORM VEGETABLES (Crop Subgroup 1C)**

Including: Potato, Sweet potato, Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Edible canna, Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (taro), Ginger, Leren, Tanier, Turmeric, Yam bean, True yam

**At-Plant**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Grape colaspis Rootworms Sweet potato flea beetle White grub Wireworms	12.75 – 25.5	0.15 – 0.3	Capture LFR insecticide may be applied as a soil incorporated broadcast, directed bed spray or a T-band spray into the planting furrow for the control of wireworms, sweet potato flea beetle, and white grubs. Apply Capture LFR Insecticide at the rate of 0.15 to 0.3 pounds active ingredient (12.75 to 25.5 fluid ounces formulated) per acre in a minimum of 10 gallons per acre of spray.
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.3 lb ai/A per season as an at-plant application.</li> </ul>			

**Lay-By**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Grape colaspis Rootworms Wireworms White grub	12.75 – 25.5	0.15 – 0.3	Capture LFR insecticide may be applied as one or more soil directed and incorporated treatments at cultivation or layby for the control of wireworms and white grubs. Apply Capture LFR insecticide to the drill area and incorporate by cultivation equipment set to throw soil towards the drill area. Apply in a minimum of 10 gallons per acre of spray.

**PPI**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Grape colaspis Rootworms Wireworms White grub	12.75 – 25.5	0.15 – 0.3	Apply Capture LFR Insecticide to the transplant area and incorporate to planting depth. Apply Capture LFR Insecticide in a minimum of 10 gallons per acre of spray. May be applied as a broadcast application or an incorporated band application

**Foliar**

PEST	USE RATES		DIRECTIONS
	Fluid oz/acre	Pound ai /acre	
Banded Cucumber beetle Black flea beetle Corn wireworm Cucumber beetle Japanese beetle grubs June beetle Rootworms Southern potato wireworm Sugarcane beetle Sweetpotato flea beetle Sweetpotato weevil Tobacco wireworm Whitefringed beetle White grub	2.8 – 8.5	0.033 - 0.1	Apply in a minimum of 3 gallons finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.  Capture LFR Insecticide may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles and May/June beetles (white grubs).
<b>Foliar Restrictions:</b> <ul style="list-style-type: none"> <li>Do not make more than 2 foliar applications per season</li> <li>Do not make applications less than 21 days apart.</li> <li>Do not apply within 21 days of harvest.</li> </ul>			

**Tuberous and Corm Vegetables (Crop Subgroup 1C) Restrictions:**

- Do not apply more than 0.5 lb ai/A per season including soil, lay-by, at-plant, PPI and foliar applications of Capture LFR Insecticide and other bifenthrin containing products.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

### Pesticide Storage

If storing this product below freezing, user should shake or roll the container to ensure proper product consistency.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call **CHEMTREC (Transportation and Spills): (800) 424-9300**.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### Container Disposal

U-Turn® Container: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

**Metal or Plastic Container: Non-refillable container (in sizes 5 gallons or less)** - Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds, pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers.

**Non-refillable container (in sizes greater than 5 gallons)** - Do not reuse or refill this container. Triple rinse or pressure rinse. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Returnable/Refillable Containers (if other than U-Turn Container):** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## Conditions of Sale and Limitation of Warranty and Liability:

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.**

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

## LABEL TRACKING INFORMATION

Label Code: 03-17-17

Brigade, Capture, LFR and FMC are Trademarks of FMC Corporation or an affiliate.

© 2016 FMC Corporation. All rights reserved.