



# SHARK<sup>®</sup>

## H2O

### HERBICIDE

## INTENDED FOR AGRICULTURAL OR COMMERCIAL USE

EPA Reg. No. 279-3194

EPA Est. 279-

Active Ingredient:	By Wt.
Carfentrazone-ethyl .....	40.0%
<b>Other Ingredients:</b> .....	<b>60.0%</b>
<b>Total</b>	<b>100.0%</b>

This product contains 40% of ingredient per pound of product.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

**Note to Physician:** Carfentrazone-ethyl is expected to have low oral and dermal toxicity, and moderate inhalation toxicity. It is expected to be slightly irritating to the skin and minimally irritating to the eyes. This product contains a granular material (sand) that may cause mechanical irritation to the eyes. Treatment is otherwise controlled removal of exposure followed by symptomatic and supportive care.

See other panels for additional precautionary information.

**ACTIVE INGREDIENT MADE IN CHINA, FORMULATED AND PACKAGED IN USA.**

## PRECAUTIONARY STATEMENTS

### Hazards to Humans (and Domestic Animals)

#### Caution

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

#### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. Do not contaminate water when disposing of equipment washwaters

#### For ground water:

"Residues of this chemical have properties and characteristics associated with chemicals detected in ground water. Residues of this chemical may leach into ground water if the chemical is used in areas where soils are permeable, particularly where the water table is shallow."

#### For surface water:

"This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of carfentrazone-ethyl residues from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours."

### Physical/Chemical Hazards

Do not use or store near heat or open flame.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

## TABLE OF CONTENTS

Section	Page
Active ingredient	1
First Aid	1
Precautionary Statements	1
Environmental Hazards	1
Directions for Use	2
Agricultural Use Requirements	2
Storage and Disposal	2
Product Information	2
Tank Mixtures	2
Adjuvant Use Requirements	2
On-Farm Testing	2
Mixing Information	2
Spray Equipment Clean-Out	3
Application Methods	3
Allowable Shark H2O Herbicide Use Information	3
Preharvest Intervals	3
Crop Rotation Restrictions	3
Rice	4
Wild Rice	4
TERMS OF SALE OR USE AND LIMITATION OF WARRANTY AND LIABILITY	5
Disclaimer	5
Label Tracking Information	5

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Long sleeve shirt and pants, Coveralls, waterproof gloves, and shoes plus socks.

### STORAGE AND DISPOSAL

**Do not contaminate water, food, or feed by storage and disposal.**

#### Pesticide Storage

Not for use or storage in or around the home. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put formulated or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills. **Call CHEMTREC (Transportation and spills): (800) 424-9300.**

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

#### Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

#### Container Handling

**Nonrefillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: **(For containers greater than 50 pounds)** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application

equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **(For containers 50 pounds or less)** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Triple rinse (or equivalent). Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Containers** - Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. If unable to return or refill, offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## PRODUCT INFORMATION

SHARK H2O Herbicide is a water dispersible granule formulation. SHARK H2O Herbicide is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops.

Weed control is optimized when the product is applied to actively growing weeds. SHARK H2O Herbicide is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation.

Extremes in environmental conditions e.g. temperature, moisture, soil conditions, and cultural practices may affect the activity of SHARK H2O Herbicide. Herbicide symptoms may be accelerated under moist conditions. Weed Control may be reduced when weeds are hardened off by drought and become less susceptible to SHARK H2O Herbicide.

SHARK H2O Herbicide is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

### TANK MIXTURES (See specific crop application instructions for additional tank mixture restrictions and information)

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SHARK H2O herbicide may be tank-mixed with other registered herbicides for controlling broader spectrum weeds. Refer to this and other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank mix partner. When preparing a new tank mix conduct an appropriate compatibility test by mixing proportional amounts of all spray ingredients in a test vessel (jar) prior to tank mixing with other products. Shake the mixture vigorously and allow it to stand for five to ten minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied. Provided the jar test indicates the mixture to be compatible, prepare the tank mixture as follows: Fill the tank one fourth full with water. With the agitator operating, add the recommended amounts of ingredients using the following order: SHARK H2O Herbicide and dry granules first, and liquid suspensions (flowables) second. As the agitation continues and the tank is filled with water add emulsifiable concentrate products third followed by the addition of water soluble products.

### ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop sections of this label for specific adjuvant type and use rates.

### ON-FARM TESTING

Not all varieties or cultivars of labeled crops have been fully evaluated under all environmental and soil conditions. Consult with your local seed company for additional information. It may also be beneficial to conduct small on-farm trials under actual conditions with specific varieties or cultivars before treating large acreage.

### MIXING INFORMATION

#### Mixing and Loading Instructions

Fill the spray tank 3/4 full with clean water. Make sure the agitation system is operating while adding products. Prepare a slurry of SHARK H2O Herbicide in a clean container using clean water. Slowly add the SHARK H2O Herbicide/water slurry. Carefully rinse the slurry container adding the rinsate to the spray tank. Complete filling the spray tank to the desired level. The spray tank agitation should be sufficient to ensure uniform spray mixture during application and must continue until the spray tank has been emptied. Follow your local extension guidelines for mixing order of products. In spray tank add dry products first, agitate, water emulsions or water soluble liquids next, emulsifiable concentrates, and then adjuvants last. Ensure the compatibility of other products and/or liquid fertilizers with SHARK H2O Herbicide before mixing them together in the spray tank.

### Mixing Precautions

Avoid the overnight storage of SHARK H2O Herbicide spray mixtures. If spray solution is stored overnight or longer, thoroughly agitate spray mixture before applying the solution. Premixing SHARK H2O Herbicide spray solutions in nurse tanks is not recommended. Maintain continuous and adequate spray solution agitation until all the spray solution has been used. Do not use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solution to alter the pH range as appropriate.

### SPRAY EQUIPMENT CLEAN-OUT

Many new pesticides are very active at low rates, especially to sensitive crops. Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying SHARK H2O Herbicide and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with SHARK H2O Herbicide as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high-pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.
2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 3% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water.
5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

Do not apply sprayer cleaning solutions or rinsate to sensitive crops.

Do not store the sprayer overnight or for any extended period of time with SHARK H2O Herbicide spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

If the sprayer has been stored or idle, purge the spray boom and nozzles with clean water before beginning any application.

Should small quantities of SHARK H2O Herbicide remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. FMC accepts no liability for any effects due to inadequately cleaned equipment.

## APPLICATION METHODS

### GROUND APPLICATION

Use ground sprayers designed, calibrated and operated to deliver uniform spray droplets to the targeted plant or plant parts. Adjust sprayer nozzles to achieve uniform plant coverage. Overlaps and slower ground speeds (caused by continuing to spray while starting, stopping or turning) may result in higher application rates and possible crop response.

#### Spray Buffer for Ground Application

Spray buffer zones for ground applications, listed in chart below, are required near desirable perennial vegetation or crops before blossom and after total leaf drop, and/or near other desirable or annual crops.

Buffers For Ground Application		
SHARK H2O USE RATE (lbs. ai per acre)	Low Spray Boom Buffer (ft.)	High Spray Boom Buffer (ft.)
0.024	20	33
0.031	26	46

### Directed Sprayers

For directed sprayers apply SHARK H2O Herbicide with drop nozzles or other post directed spray equipment.

### Post-Directed Applications

Post-directed applications may be utilized when labeled crops have reached minimum growth stages where sprays may be directed to the target weeds, but is not deposited on the green stem, foliage, blooms or fruit of the crop. Do not apply when conditions favor drift or when wind is above 10 miles per hour. Use drop nozzles or other spray equipment capable of directing the spray to target weeds and away from sensitive plant parts. Apply when labeled crops have reached minimum growth stages described in specific crop sections of this label and when spray will not be deposited on green stems, foliage, blossoms or fruit.

### SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

SHARK H2O Herbicide is a contact PPO (protoporphyrinogen oxidase) herbicide. Avoid any drift conditions that would allow the product to contact desirable vegetation. SHARK H2O herbicide is not volatile, however; mist from spray drift may cause injury to sensitive plants.

The interaction of equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications of dry materials.

Where states have more stringent regulations, they must be observed.

### ALLOWABLE SHARK H2O HERBICIDE USE INFORMATION

Refer to the crop section of this label for specific product use directions.

Table 1:

Maximum Allowable SHARK H2O Herbicide Use Per Acre Per Season for Crops or Crop Group.		
Crop	SHARK H2O Herbicide (dry oz/acre) Per Season	Maximum Rate (lb ai/acre) Per Season
Rice	12	0.3
Wild Rice	12	0.3

### PREHARVEST INTERVALS

Refer to the crop section of this label for specific product use directions.

Table 2:

Preharvest Intervals (PHI) or Maximum Growth Stage for SHARK H2O Herbicide Applications	
Crop	PHI (Days Before Harvest) or Growth Stage
Rice (preplant and in-season)	60
Wild Rice	60

### CROP ROTATIONAL RESTRICTIONS

Following an application of SHARK H2O Herbicide, a treated field may be rotated to a registered crop at any time, subject to specific crop restrictions that may be found in the individual crop sections. All other crops may be planted after 12 months.



**RICE:**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Early Post Seeding Applications to Submerged Weeds</b>	Do not apply within 60 days of harvest.	See weed list in table 3 below.	7.6 dry oz (0.19 pounds active ingredient) per acre	Do not apply by air. Do not apply within 1/2 mile of sensitive crops. Do not apply when conditions favoring drift exist.
<b>Foliar Applications to Emerged Weeds Above the Water Surface</b>	Do not apply within 60 days of harvest.	See weed list in table 4 below	Up to 4.0 dry oz (0.10 pounds active ingredient) per acre	Pre-flood treatment, once field is flooded, water must be held for at least 23 days following treatment before release. Do not apply more than 7.6 dry oz (0.19 pounds active ingredient) per single application. Do not apply more than 12.0 dry oz (0.3 pound active ingredient) per acre per season including fallow, preplant, burndown, and labeled crop applications. Do not release water for at least 23 days following a Post Flood treatment in the water.

**DIRECTIONS FOR USE**

Apply SHARK® H2O Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Applications shall be made by ground equipment only using a minimum finished spray volume of 10 gallons of spray per acre.

To control weeds not listed on this label, SHARK® H2O Herbicide may be tank mixed with other herbicides registered for use on rice. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and restrictions.

**Early Post Seeding Applications to Submerged Weeds**

Apply SHARK® H2O Herbicide at 7.6 dry oz (0.19 pounds active ingredient) per acre. Evenly distribute the spray solution over the flooded rice. The floodwater must be 3 to 6 inches deep. Apply at 1.5 leaf stage of rice. Earlier applications may cause unacceptable crop response. Rice must be well rooted and actively growing at the time of application. Hold the floodwater at a static depth for at least five days after application of SHARK® H2O Herbicide. Once field is flooded, water must be held for at least 23 days following treatment before release.

**When used as directed SHARK® H2O Herbicide will provide control of listed weeds at the 2 leaf stage or less.**

**Table 3:**

Arrowhead, California	Bulrush, ricefield
Ammannia, purple (suppression only)	Umbrellaplant, smallflower (suppression only)
Ammannia, redstem (suppression only)	

**Tank Mixtures** (See "Tank Mixtures" for additional tank mixture requirements) It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SHARK® H2O Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturers' label directions for the companion herbicide except for specific directions on this label. Apply SHARK® H2O Herbicide before, after, or with an application of Londax®, Ordram® and Bolero® herbicides. Observe all applicable directions, restrictions (including water holding requirements) and precautions on the Londax, Ordram and Bolero labels.

Do not apply SHARK® H2O Herbicide as a tank mixture with Regiment®.

**Foliar Applications to Emerged Weeds Above the Water Surface**

Apply SHARK® H2O Herbicide to weeds up to 4.0 dry oz (0.10 pound active ingredient) per acre to the foliage of exposed weeds. At least 80% of the weed foliage must be exposed before spraying SHARK® H2O Herbicide. For optimum results, apply to actively growing weeds 20 to 45 days postseeding or the earliest practical opportunity to spray. Weed control is enhanced with greater weed exposure. If the field was drained at application, reflood twenty-four hours after application to the normal flood depth.

**When used as directed SHARK® H2O Herbicide will provide control or suppression of the following weeds.**

**Table 4:**

Bulrush, ricefield	Ammannia, redstem (suppression only)
Arrowhead, California	Umbrellaplant, smallflower (suppression only)
Ammannia, purple (suppression only)	

**Crop Response**

Some temporary leaf speckling may occur shortly after application.

**Tank Mix** (See "Tank Mixtures" for additional tank mixture requirements) It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SHARK® H2O Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. SHARK® H2O Herbicide may be tank mixed with propanil-containing herbicides, Londax®, Bolero®, or Whip® herbicides. Not all combinations of SHARK® H2O Herbicide and other formulated herbicides have been tested. In general, the EC formulations, nonionic and silicone based surfactants and crop oil concentrates, when mixed with SHARK® H2O Herbicide will increase leaf speckling on the rice leaves. These tank mixtures should be tested on a small portion of the field to ensure crop safety prior to general use.

**Crop Rotation Restriction:**

After an application of this product to rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**RICE, Wild: Wild rice grown in cultivated fields where the water discharge / release can be controlled.**

Methods and Timing	PHI	Target Weeds	Rates	Restrictions
<b>Postemergence Weed Control</b>	Do not apply within 60 days of harvest.	See weed list in table 5.	4.0 to 7.6 dry oz (0.1 to 0.19 pounds active ingredient) per acre	Do not apply when conditions favoring drift exist. Do not apply when winds exceed 10 mph. Do not apply more than 7.6 dry oz (0.19 pounds active ingredient) per single application. Do not apply more than 12.0 dry oz (0.3 pound active ingredient) per acre per season, including fallow/preplant, burndown, and labeled crop applications. Do not apply during the floating leaf stage when exposed wild rice leaves are most susceptible to injury. Do not apply to wild rice when there is heavy dew on the leaves or under high humidity conditions. Do not release flood water off wild rice field(s) for a minimum of 23 days after application of this product. (Specific to California) Do not apply within 0.5 mile of sensitive crops (for California wild rice). Do not apply to wild rice by air in California Do not release flood water off wild rice field(s) for a minimum of 23 days after application of this product in California.

**DIRECTIONS FOR USE**

Apply SHARK® H2O Herbicide alone or as a tank mixture with other rice herbicides to emerged and actively growing weeds. Applications shall be made by ground equipment using a minimum finished spray volume of 10 gallons per acre.

Apply SHARK® H2O Herbicide to weeds at the rate of 4.0 to 7.6 dry oz (0.1 to 0.19 pound active ingredient) per acre to the foliage of exposed weeds above the water surface. Make applications after the floating leaf stage through tillering. The water in paddies may be lowered if practical. Smaller weeds with more leaf area exposed will give better control. If water is lowered for application, it may be re-flooded to normal depths 24 hours after the application

**When used as directed Shark H2O Herbicide will provide control or suppression of the following weeds.**

**Table 5:**

Ammannia, purple (Suppression only)	Burrweed, giant (Suppression only)
Ammannia, redstem (Suppression only)	Umbrellaplant, smallflower (Suppression only)
Arrowhead, California	Waterplantain, common (Suppression only)
Bulrush, ricefield	

**Crop Response**

Some temporary leaf speckling may occur following application. Wild rice should be well rooted and vigorously growing at the time of application. Earlier applications may cause unacceptable crop response.

**Tank Mix** (See "Tank Mixtures" for additional tank mixture requirements)

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SHARK® H2O Herbicide may be tank mixed with other herbicides to control weeds not listed on this label. Not all combinations of SHARK® H2O Herbicide and other formulated herbicides and adjuvants have been tested. In general, EC formulations, nonionic and silicone based surfactants, and crop oil concentrates, will increase leaf speckling on the wild rice leaves. These tank mixes should be tested on a small portion of the field to ensure crop safety prior to general use.

**Precautions**

Wet leaf surfaces at the time of application can cause unacceptable injury.

**Crop Rotation Restriction**

After an application of this product to wild rice, you may only rotate the field to a carfentrazone-ethyl registered crop.

**TERMS OF SALE OR USE AND LIMITATION OF WARRANTY AND LIABILITY**

**Terms of Sale and/or Use**

On purchase of this product buyer and user agree to the terms and conditions as follow.

**Packaging**

Distributors/Dealers/Retailers shall sell in original packages only.

**Warranty**

FMC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use section when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of seller or FMC), and buyer assumes the risk of any such use.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risk inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC of Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

**Use of Product**

FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**DISCLAIMER**

FMC Corporation intends that SHARK H2O Herbicide be distributed only to end users and/or growers (and/or applicators acting on the behalf of growers), who agree to the terms and conditions as stated herewith and further agree to a waiver and release from any and all liability by the user and/or grower of FMC for failure to perform and/or crop damage resulting from the use of SHARK H2O Herbicide as recommended on the labeled crops under the those specific sections of this label. If such terms and conditions are unacceptable, FMC requests the return at once of all products in unopened original containers. FMC considers the user to have accepted such terms and conditions upon the use of SHARK H2O Herbicide.

SHARK H2O Herbicide, when used as directed, may result in crop injury, crop loss or crop damage. FMC recommends that the user and/or grower test SHARK H2O Herbicide in order to determine its suitability for the intended use. FMC makes SHARK H2O Herbicide available to the user and/or grower solely to the extent that the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of SHARK H2O Herbicide. The decision to use, or not to use, SHARK H2O Herbicide must be made by each individual user and/or grower on the basis of possible crop injury from SHARK H2O Herbicide, the severity of weed infestations, the cost of alternative weed control measures and other factors. Because of the risk of crop damage, all such use is at the user and/or grower's risk.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC OR SELLER SHALL NOT BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OF OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These TERMS OF SALE OR USE AND LIMITATION OF WARRANTY AND LIABILITY may not be amended by any oral or written agreement.

FMC and SHARK are trademarks of FMC Corporation

Londax is a trademark of E.I. DuPont de Nemours and Company

Whip is a trademark of Bayer

Ordram is a trademark of ZENECA Group Company

Bolero is a trademark of Valent USA Corporation

Regiment is a trademark of Kumiai Chemical Industry Company, Ltd.

© 2016 FMC Corporation. All rights reserved.