

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

Tundra® EC

Agricultural Insecticide

For Agricultural or Commercial Use Only

FIFRA 24(c) Special Local Need Registration

EPA SLN No. WY-21-002

EPA Reg. No. 1381-196

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF WYOMING

This label expires and shall not be distributed or used in accordance with this SLN registration after December 31, 2026.

ACTIVE INGREDIENT:	By Wt.
Bifenthrin* (2 methyl[1,1'-biphenyl]-3-yl) methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-Dimethylcyclopropanecarboxylate	25.1%
INERT INGREDIENTS	74.9%
TOTAL	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.

**Contains xylene range aromatic solvents

This product contains 2 pounds active ingredient per gallon.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered container label.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

CROP	PEST	RATE OF APPLICATION
Alfalfa Grown for Seed	Lygus Bugs Weevils Aphids Spider Mites	3.9 to 6.4 fl. ounces per acre

To control Lygus Bugs, Weevils, Aphids and Spider Mites, apply Tundra EC at the rate of 3.9 to 6.4 fl. ounces (0.06 to 0.1 lb. active) per acre. Apply in a minimum of 5 gallons of water per acre by air equipment or in a minimum of 20 gallons of water by ground equipment.

Maintain a minimum of 21 days between Tundra EC treatments.

DIRECTIONS FOR USE CONTINUED

Make aerial or ground applications when the wind velocity favors on target product disposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. Boom length must not exceed 75% of wingspan or 80% rotor diameter.

All seed from alfalfa treated with Tundra EC is to be tagged at the conditioning plant, "Not for human or animal consumption." It shall be the grower's responsibility to notify the conditioning plant of any alfalfa seed crop treated with Tundra EC. Screenings and seed from seed conditioning are prohibited from feed channels. Seed from alfalfa treated with Tundra EC must not be used to grow sprouts for human or animal consumption.

Refer to the container label for additional **Spray Drift Requirements** and for information on Chemigation.

RESTRICTIONS

Do not apply Tundra EC or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply by ground equipment within 25 feet, or by air (non-ULV) within 150 feet of lakes, reservoirs, rivers, permanent streams, marches or natural ponds, estuaries, and commercial fish farm ponds.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetative filter strip.

Do not apply more than 6.4 ounces (0.1 lb. active) per acre per treatment. Do not make more than 3 applications in a crop year.

Do not cut current year's treated alfalfa seed crop for hay or forage.

Do not graze current year's treated alfalfa seed crop.

All Tundra EC treated alfalfa seed screenings must be removed from the feed market.

ROTATIONAL CROPS

Crops for which bifenthrin residue tolerances exist, may be rotated at any time. For all other crops, wait at least 30 days following the final application of bifenthrin before rotating.

24(c) Registrant:

Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0089