



Surround® WP

Crop Protectant

EPA Reg. No. 61842-18

FOR USE IN AIDING THE ESTABLISHMENT OF TRANSPLANTS OF ANNUAL FRUITS AND VEGETABLES, PERMANENT CROPS AND ALL OTHER CROPS LISTED ON THE SURROUND WP LABEL IN THE USA

This recommendation is made as permitted under FIFRA Section 2 (ee) and has not been submitted to or approved by the EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply **Surround WP** to new transplants in the field to reduce transplant shock and heat stress. Re-apply **Surround** every 5-7 days depending on rainfall/irrigation, plant growth and type of crop. Emerging green tissue should be protected from sunburn and heat stress. Once plants reach full size or start to bear fruit, follow the **Surround** Section 3 label. Certain crops whose fruit cannot be washed post-harvest, such as strawberries, must not be sprayed with **Surround** once they start bearing fruit.

Rate & Concentration: Use concentrations of 25 to 50 lbs **Surround WP** per 100 gallons of water in sufficient volumes per acre to spray just to drip. This will be 6.25 to 37.5 lbs (depending on transplant size and crop) of **Surround WP** per acre. Small seedlings will need less water per acre using a band spray technique. Adjust spray method and water volume for larger or more mature plants to uniformly coat plant surfaces. Adjust the volume of water and rate used throughout the season based on plant size.

Consult the **Surround** label and do not exceed the rates indicated for the type of transplant being treated.

Do not use anti-transpirant type products with **Surround**. These materials may block stomata, thereby restricting efficiency of photosynthesis.

APPLICATION TIPS:

- Avoid foliage run-off. Apply the spray volume that will achieve near-drip coverage.
- Surfactants may improve spreading.
- Consult the label for instructions on aerial application.
- See **Surround** label for more information on application techniques.
- **Surround** can be applied with most other insecticides, miticides and fungicides. Conduct a compatibility test to ensure tank-mix compatibility. Add **Surround** to the spray tank first.
- Do not apply **Surround** through any type of overhead irrigation system or with overhead cooling.
- Do not dip transplants into **Surround** solution.

IMPORTANT NOTICE

BEFORE BUYING OR USING TESSENDERLO KERLEY PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS.

It is a violation of federal law to use this or any pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced Tessenderlo Kerley product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of Surround product labels are available at <http://www.novasource.com/products>.

Tessenderlo Kerley, Inc.
2255 N. 44th Street, Suite 300
Phoenix, AZ 85008 USA
1-800-525-2803
www.novasource.com

Surround is a registered trademark of Tessenderlo Kerley, Inc.
©2009 Tessenderlo Kerley, Inc. All Rights Reserved.

