

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



## 50 SG INSECTICIDE

EPA Reg. No. 71512-10-279

<b>2(ee) Recommendation</b>
<b>FOR DISTRIBUTION AND USE ONLY IN  NEW YORK</b>
THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA
<b>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING</b>

CROP	PEST	RATE OF APPLICATION
Apples	Codling Moth	2.8 oz / acre.
	San Jose Scale	2.8 oz / acre.

**DIRECTIONS FOR USE:**

**Codling Moth**

For preventative use to suppress first generation codling moth. Once biofix has been determined begin applications no earlier than 200 DD's through peak egg deposition and prior to egg hatch. Beleaf will aid in control of second generation codling moth when tankmixed with another insecticide labeled for second generation codling moth control and applied according to the tankmix partners label. Repeat applications may be required under prolonged flight conditions. Thorough coverage is essential to achieve control. Apply by ground in a minimum of 50 gallons of finished spray and 10 gallons by air to obtain full coverage of fruit and foliage.

**San Jose Scale**

For preventative use to control San Jose Scale. Applications can occur as early as pink, petal fall or anytime prior to crawler emergence. Apply by ground in a minimum of 50 gallons of finished spray and 10 gallons by air to obtain full coverage of fruit and foliage.

**RESTRICTIONS:**

- Do not apply more than 2.8 oz/A Beleaf 50 SG (0.089 lbs ai/acre) per application.
- Do not apply more than 8.4 oz/A Beleaf 50 SG (0.267 lbs ai/acre) per season.
- Do not apply more than 3 consecutive applications at the 2.8 oz/A Beleaf 50 SG rate per season.
- Allow a minimum of 7 days between applications.

**PHI: 21 Days**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED BELEAF 50 SG INSECTICIDE LABEL (EPA REG. NO. 71512-10-279) MUST BE FOLLOWED.  
THE USER SHOULD POSSESS THIS RECOMMENDATION AT THE TIME OF PESTICIDE APPLICATION.**