

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA

Group 9C Insecticide



Beleaf
INSECTICIDE

EPA Reg. No. 71512-10-279

For Agricultural or Commercial Use Only

2(ee) Recommendation
NOT FOR USE IN
IN, KY, NH, NM, NY, SD & TX
THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

Crop	Pest	Rate of Application
Potatoes	Potato Psyllid: <i>(First and Second Instar Only)</i>	2.8 oz/acre

DIRECTIONS FOR USE:

Begin applications before populations begin to build and before damage is evident, according to local pest management guidelines. For control of adults, apply in combination with other effective products labeled for use in potatoes. Rapidly growing plants may need to be retreated. Scout fields often and retreat as necessary to maintain populations below damaging levels.

Thorough Coverage is essential for optimum control. Apply sufficient volume to ensure good coverage; use a minimum of 10 gallons per acre by ground and 5 gallons air. Finished volumes should be increased under extreme pest populations or dense plant foliage. Apply a maximum of 0.25 inches if applying through chemigation.

RESTRICTIONS:

Do not apply more than 2.8 oz/acre Beleaf 50SG per application; do not apply more than 8.4oz/acre of Beleaf 50SG per season. Do not apply more than 3 applications at the 2.8 oz/acre Beleaf 50SG rate per season. Allow a minimum of 7 days between applications.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED BELEAF 50 SG INSECTICIDE LABEL (EPA REG. NO.71512-10-279) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation
1735 Market Street
Philadelphia, PA 19103