

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms. For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



FIFRA SECTION 24 (C)
Special Local Need Label
Legume Vegetables (Dried)

EPA Reg. No. 279-3302
 EPA SLN. No. MT-15-0003

FIFRA Section 24 (c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA

This label for Capture LFR Insecticide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2018.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Crop	Pest	Rate of Application
DRIED SHELLED PEAS AND BEANS Pre-Plant Incorporated (PPI)	Wireworm Seed Corn maggot Grape colapsis White Grubs Root maggot Black cutworm Armyworm spp.	5.3 – 6.8 fluid oz/acre (0.062 - 0.1 lb ai/ acre)

DIRECTIONS FOR USE: Dried shelled pea and bean (except soybean) subgroup.

FOR PRE-PLANT INCORPORATED APPLICATIONS (PPI) TO DRIED SHELLED PEA AND BEANS.

Dried cultivars of bean (*Lupinus*) ; bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (*Pisum*) (includes field pea); pigeon pea.

When applied as directed, Capture LFR will provide control of wireworm. For PPI treatments, the 5.3- 6.8 oz/A rate

must be used. Capture LFR can be tank mixed and applied with PPI herbicides. Incorporation of Capture LFR should no deeper than 3 inches. Incorporation should be close to the intended seed planting depth.

Restrictions:

Do not apply more than 0.2 pound active ingredient per acre per season to peas including pre, ppi, at-plant and foliar applications of other bifenthrin containing products.

Do not apply more than 0.3 pounds active ingredient per acre per season to beans including pre, ppi, at-plant and foliar applications of other bifenthrin containing products.

Do not apply more than 0.1 pound active ingredient per acre per season as a pre-plant, PPI, or at-plant application.

This pesticide is extremely toxic to fish and aquatic invertebrates. Capture LFR Insecticide should not be used under this SLN label where impact on listed threatened or endangered species is likely. **Refer to U.S. Fish & Wildlife Services website, <http://ecos.fws.gov/ipac/> for location specific information on threatened and endangered species.** Consult the Section 3 label for additional restrictions and precautions to protect aquatic organisms.

FMC and Capture LFR are trademarks of FMC Corporation. Capture LFR is a restricted use pesticide. All rights reserved.



FMC Corporation
Agricultural Solutions
1735 Market Street
Philadelphia, PA 19103

©All rights reserved Rev:10-6-2015