

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



The Chemical Company

SPECIMEN

RESPECT[®]

EC INSECTICIDE

Active Ingredient:

* S-Cyano (3-phenoxyphenyl)methyl (+) cis/trans 3-(2,2-dichloroethenyl)-
2,2 dimethylcyclopropane carboxylate** 9.6%

Inert Ingredients:*** 90.4%

Total: 100.0%

* Contains 0.8 pounds active ingredient per gallon

** Cis/trans ratio: Max. 75% (±) cis and min. 25% (±) trans

*** Contains petroleum distillates

U.S. Patent No. Pending

EPA Reg. No. 279-3327-7969

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Limitation of Warranty and Liability**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call FMC day or night 1-800-331-3148.

Net Contents:

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact FMC at 1-800-331-3148 for emergency medical treatment information.	
<p>Note to Physician: Contains petroleum distillates – vomiting may cause aspiration pneumonia. Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages by aspiration of the solvent.</p> <p>For Emergency Assistance Call FMC, 1-800-331-3148.</p> <p>See other panels for additional precautionary information.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS. Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for **Category G** on an EPA chemical resistance category selection chart.

Handlers who may be exposed to the dilute through application or other tasks must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as nitrile, butyl, neoprene, and/or barrier laminate
- Shoes plus socks.

Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as nitrile, butyl, neoprene, and/or barrier laminate
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates, oysters and shrimp. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

DO NOT contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds.

DO NOT apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

Physical/Chemical Hazards

DO NOT use or store near heat or open flame.

Directions For Use

RESTRICTED USE PESTICIDE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as Barrier Laminate or Viton
- Shoes plus socks

STORAGE AND DISPOSAL

Pesticide Storage

Store in a cool, dry, well-ventilated place. **DO NOT** store below -6.6° C (20° F). If solids are observed, warm to above 4.4° C (40° F) and roll or shake containers to redissolve. **DO NOT** use near heat, open flame or hot surfaces. Store in original containers only. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put concentrate or dilute material into food or drink containers. **DO NOT** contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Keep out of reach of children and animals.

In Case of Spill

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC 1-800-331-3148

To Confine Spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Metal containers. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **DO NOT** cut or weld metal containers.

Plastic containers. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Returnable/refillable sealed containers. DO NOT rinse container. **DO NOT** empty remaining formulated product. **DO NOT** break seals. Return intact to point of purchase.

Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system. **DO NOT** connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact state Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Respect® EC insecticide should be applied continuously for the duration of the water application. **Respect EC** should be diluted in sufficient volume to ensure accurate application over the area to be treated. Use the appropriate amount of water to carry the product to the target pest. Agitation is not required when a suitable diluent is used.

BUFFER ZONES

Vegetative Buffer Zones

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing zeta-cypermethrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services*. USDA, NRCS. 2000. Fort Worth, Texas. 21pp. <http://www.in.nrcs.usda.gov/technical/agronomy/new-conbuf.pdf>.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)

DO NOT apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application

DO NOT apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application

DO NOT apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Spray Drift Requirements

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. **DO NOT** apply when the wind velocity exceeds 15 mph.

Temperature Inversion

DO NOT make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

- Wind speed must be adjacent to the application site on the upwind side, immediately prior to application.
- For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.
- For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

- The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.
- Flight speed and nozzle orientation must be considered in determining droplet size.
- Spray must be released at the lowest height consistent with pest control and flight safety. **DO NOT** release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
- When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

General Instructions

Use low rate under light-to-moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. **DO NOT** exceed maximum allowable rate.

Preventive Use

For cutworm, armyworm, or stalk borer control, **Respect® EC insecticide** may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control.

Rotational Crops

With the exception of the crops listed below, rotational crops should not be planted within 30 days of last application.

Tank Mixture

Respect EC may be applied in tank mixtures with other products approved for use on Alfalfa and Nongrass Animal Feeds, Brassica Vegetables, Bulb Vegetables, Corn, Cotton, Fruiting Vegetables, Legume Vegetables, Peanut, Root and Tuber Vegetables, Sorghum, Soybeans, Sugarcane, Tree Nuts and Wheat. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

Maximum Usage When Applying Both Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season

DO NOT apply more than the maximum seasonal total for either product when used alone, and **DO NOT** apply more than the combined maximum seasonal total for both products as outlined in the table below.

Crop	Maximum Seasonal Total (lb ai/acre)				Maximum Seasonal Total (lb ai/acre) When Applying Cypermethrin and Zeta-Cypermethrin Products to the Same Crop	Maximum Seasonal Total (lb ai/acre) When Applying Zeta-Cypermethrin Products to the Same Crop
	Zeta-Cypermethrin*			Cypermethrin**	Zeta-Cypermethrin* plus Cypermethrin**	Zeta-Cypermethrin*
	Mustang®	Respect® EC insecticide	HERO™			
Cotton	0.3	0.15	0.1125	0.6	0.6	0.3
Field Corn	0.2	0.10	0.10	NA	NA	0.2
Sweet Corn	0.3	0.15	0.0675	NA	NA	0.3
Eggplant	0.3	0.15	0.0675	NA	NA	0.3
Pepper (Bell and Non-Bell)	0.3	0.15	0.0675	NA	NA	0.3
Tomato	0.3	0.15	0.105	NA	NA	0.3
Head Lettuce	0.3	0.15	0.1125	0.6	0.6	0.3
Head and Stem Brassica	0.3	0.15	0.1125	0.6	0.6	0.3
Succulent Peas and Beans	0.3	0.15	0.0675	NA	NA	0.3
Pecans	0.3	0.15	0.1125	0.6	0.6	0.3

***Mustang** or **Fury**® (1.5 EW or 1.5 EC); **Respect EC** (0.8 EC); **Z-Cype** (0.8 EW); **HERO**; or any **Zeta-Cypermethrin** product approved for crop use

Any Cypermethrin product approved for crop use, including **Ammo®

NA = Not Applicable

Maximum Seasonal Usage and PHI (Preharvest Interval) for Respect® EC insecticide-Labeled Crops

Crop	Maximum Seasonal Total/Acre for Respect EC		PHI (days)
	lb ai	fl ozs	
Alfalfa and Nongrass Animal Feeds (Forage, Fodder, Straw and Hay) Group	0.025/cutting	4.0	3 (cutting or grazing)
	0.075/season	12.0	7 (harvesting seed)
Brassica Vegetables	0.15	24.0	1
Bulb Vegetables	0.125	20.0	7
Corn, Sweet	0.15	24.0	3
Corn Field Seed Pop	0.10	16.0	30 (grain and stover) 60 (forage)
Cotton	0.15	24.0	14
Fruiting Vegetables	0.15	24.0	1
Head Lettuce	0.15	24.0	1
Legume Vegetables	0.15	24.0	1 (succulent shelled or edible-podded) 21 (dried shelled)
Peanut	0.15	24.0	7
Root and Tuber Vegetables (except Sugar Beet)	0.15	24.0	1
Sorghum	0.125	20.0	14 (grain and fodder stover) 45 (forage (silage))
Soybeans	0.15	24.0	21
Sugarcane	0.10	16.0	21
Tree Nuts	0.125	20.0	7
Wheat	0.125	20.0	14

The REI (restricted-entry interval) is **12 hours** for all labeled crops.

Refer to the crop-specific use directions for detailed information on application timing and any use restrictions.

Nongrass Animal Feeds (Forage, Fodder, Straw and Hay) Group including: Alfalfa; Alfalfa grown for seed (Includes lucerne, sainfoin, holy clover, esparcet, birdsfoot trefoil and varieties and/or hybrids of these); **Velvet Bean; Clover; Kudzu; Lespedeza; Lupin; Sainfoin; Trefoil; Vetch; Crown Vetch; and Milk Vetch**

Insects Controlled	Rate of Application	Method of Application
Alfalfa Caterpillar Alfalfa Looper Alfalfa Weevil Cutworms Egyptian Alfalfa Weevil (larvae and adult) Flea Beetles Green Cloverworm Hornworms Meadow Spittlebug Potato Leafhopper Velvetbean Caterpillar Webworms Blue Alfalfa Aphid ¹ Green Peach Aphid ¹ Pea Aphid ¹ Spotted Alfalfa Aphid ¹ Three-cornered Alfalfa Hopper	2.24 to 4.0 ounces (0.014 to 0.025 pound active) per acre	Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment. ULV oil spray application is prohibited. Higher volumes of finished spray may improve insect control under high temperatures, when foliage is dense and/or when insect pressure is high. DO NOT exceed maximum allowable rate.
Armyworms Grasshoppers Plant Bugs (including Lygus spp. and Stink Bugs)	2.8 to 4.0 ounces (0.0175 to 0.025 pound active) per acre	
<p>DO NOT make applications less than 7 days apart.</p> <p>A maximum of 4.0 fl ozs product (0.025 pound active ingredient) per acre may be applied per cutting and a maximum of 12.0 fl ozs product (0.075 pound active ingredient) per acre per season.</p> <p>Applications may be made up to 3 days of cutting or grazing or up to 7 days of harvesting seed.</p> <p>¹Aphid control may be variable depending on species present and host-plant relationships.</p>		

Head and Stem Brassica Vegetables (1-day PHI) including: Broccoli; Chinese Broccoli (gai lon, white flowering broccoli); **Brussels Sprouts; Cauliflower; Cavalo Broccolo; Kohlrabi; Cabbage; Chinese Cabbage** (napa); **Chinese Mustard Cabbage** (gai choy); **Leafy Brassica Greens including: Broccoli Raab** (rapini); **Chinese Cabbage** (bok choy); **Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens; and Turnip Greens**

Insects Controlled	Rate of Application	Method of Application
Corn Earworm Cucumber Beetles Cutworm Diamondback Moth ¹ Flea Beetles Imported Cabbageworm Leafhoppers Saltmarsh Caterpillar Southern Cabbageworm Tobacco Budworm ¹	2.24 to 4.0 ounces (0.014 to 0.025 pound active) per acre	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Respect® EC insecticide should be used under light-to-moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations.
Alfalfa Looper Armyworms Cabbage Looper Cabbage Webworm Crickets Grasshoppers Ground Beetles Leafminers (adults) Lygus Bugs Onion Thrips Stinkbugs Wireworm (adults) Aphids ² Whiteflies ³	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. DO NOT exceed maximum allowable rate.

DO NOT make applications less than 7 days apart.

A maximum of 24.0 fl ozs product (0.15 pound active ingredient) may be applied per acre per season.

¹See the resistance statement under the **Directions For Use** section.

²Aphid control may be variable depending on species present and host-plant relationships.

³Aids in control

Head Lettuce (1 day PHI)

Insects Controlled	Rate of Application	Method of Application
Aphid spp. ^{2,3} Corn Earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Saltmarsh Caterpillar Tobacco Budworm ² Whitefly spp. ^{1,2}	2.24 to 4.0 ounces (0.014 to 0.025 pound active) per acre	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Respect® EC insecticide should be used under light-to-moderate insect pressure. Higher rates should be used to control heavy-to-extremely heavy insect populations. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required.
Armyworms Crickets Ground Beetles Loopers Lygus Bugs Onion Thrips Stink Bugs Wireworm (adults)	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	DO NOT exceed maximum allowable rate.
<p>DO NOT make applications less than 7 days apart.</p> <p>A maximum of 24.0 fl ozs product (0.15 pound active ingredient) may be applied per acre per season.</p> <p>¹Aids in control ²See resistance statement under the Directions For Use section. ³Aphid control may be variable depending on species present and host-plant relationships.</p>		

Bulb Vegetables (*Allium* spp.) (7-day PHI) including: Garlic; Garlic, Great-Headed (elephant); Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welch; Shallots, Dry Bulb and Green; Spring Onion or Scallions

Insects Controlled	Rate of Application	Method of Application
Armyworms Cutworms Leafminers (adults) Onion Maggot Adults Stink Bugs Aphids ¹	2.24 to 4.0 ounces (0.014 to 0.025 pound active) per acre	Apply in a minimum of 20 gallons per acre with ground equipment or in a minimum of 3 gallons per acre by aircraft. Begin applications when pests appear and repeat as necessary to maintain control.
Onion Thrips	2.88 to 4.0 ounces (0.018 to 0.025 pound active) per acre	To control Onion Thrips: Use higher rates as population increases and avoid rescue situations. Use of a crop oil concentrate at 16 fluid ounces per acre is recommended. DO NOT exceed maximum allowable rate.
<p>DO NOT make applications less than 7 days apart.</p> <p>DO NOT apply more than 20.0 fl ozs product (0.125 pound active ingredient) per acre per season.</p> <p>DO NOT graze livestock in treated areas or cut treated crops for feed.</p> <p>¹Aphid control may be variable depending on species present and host-plant relationships.</p>		

Corn, Sweet (3-day PHI)

Insects Controlled	Rate of Application	Method of Application
Chinch Bug Corn Rootworm (adult) Corn Silkfly Cutworms Flea Beetle Leafhoppers Japanese Beetle (adult) Sap Beetle (adults) Tarnished Plant Bug	2.24 to 4.0 ounces (0.014 to 0.025 pound active) per acre	Apply with ground or air equipment using sufficient water and application methods to ensure thorough coverage of foliage. Apply in water using a minimum of 20 gallons of finished spray per acre with ground equipment and a minimum of 2 gallons per acre by air.
Armyworms Corn Borers Corn Earworm Grasshoppers Aphids ¹	2.8 to 4.0 ounces (0.0175 to 0.025 pound active) per acre	

Apply at minimum 3- to 5-day intervals or as needed for control.

A maximum of 24.0 fl ozs product (0.15 pound active ingredient) may be applied per acre per season.

DO NOT apply within 3 days of harvest of ears or forage or livestock grazing.

¹Aphid control may be variable depending on species present and host-plant relationships.

Corn (Field), Field Corn Grown for Seed, Popcorn (At Plant Use)

Insects Controlled	Rate of Application	Method of Application		
Cutworms	0.16 fluid ounce per 1,000 linear feet of row (0.001 pound active) per 1,000 linear feet of row	Apply as an in-furrow, band or T-band treatment using a minimum 4-inch band. Use table below to determine the Respect® EC insecticide needs for each acre.		
Row Spacings (inches)		40	30	20
Respect EC (pounds ai per acre)		0.012	0.018	0.024
Respect EC (formulated ounces per acre)		1.92	2.88	3.84

DO NOT apply more than 16.0 fl ozs product (0.10 pound active ingredient) per acre per season, including at-planting plus foliar applications of **Respect EC**.

DO NOT apply within 30 days of harvest for grain and stover and 60 days for forage.

Corn (Field), Field Corn Grown for Seed, Popcorn

Insects Controlled	Rate of Application	Method of Application
Cutworms	1.28 to 2.8 ounces (0.008 to 0.0175 pound active) per acre	Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results.
Corn Earworm ¹ Green Cloverworm Meadow Spittlebug Western Bean Cutworm ¹	1.76 to 4.0 ounces (0.011 to 0.025 pound active) per acre	
Bean Leaf Beetle Cereal Leaf Beetle Corn Borer, European Corn Borer, Southwestern Corn Rootworm Beetle Flea Beetle Grasshoppers Hop Vine Borer Hornworms Japanese Beetle (adult) Sap Beetle (adult) Southern Corn Leaf Beetle Stalk Borer Stink Bug spp. Tobacco Budworm ² Webworms Aphids ³	2.72 to 4.0 ounces (0.017 to 0.025 pound active) per acre	<p>Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground).</p> <p>For chinch bug control, scout corn fields and make applications when bugs migrate from small grains or wild grasses to small corn. Direct spray to the base of plant. Repeat applications at 3- to 5-day intervals if needed. Respect® EC insecticide may only suppress heavy infestations and/or subsequent migrations.</p> <p>DO NOT exceed maximum allowable rate.</p>
Armyworms (including Fall Armyworms) Chinch Bug	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	
<p>DO NOT apply more than 16.0 fl ozs product (0.10 pound active ingredient) per acre per season, including at-planting plus foliar applications of Respect EC.</p> <p>DO NOT apply within 30 days of harvest for grain and stover and 60 days for forage.</p> <p>¹For control before the larva bores into the plant stalk or ear</p> <p>²See the resistance statement under the Directions For Use section.</p> <p>³Control may be variable depending on species present and host-plant relationships.</p>		

Cotton (14-day PHI)

Insects Controlled	Rate of Application	Method of Application
Preemergent Use: Cutworms	1.28 to 1.92 ounces (0.008 to 0.012 pound active) per acre	Use Respect® EC insecticide in the time period from 14 days prior to planting up to emergence of the crop. Apply as a broadcast spray by ground or air, banded (including T-band) or in-furrow spray using sufficient spray volume to achieve adequate coverage. Reduced volumes of water may be used with specialized equipment. Use the higher rates of Respect EC when incorporating into the soil. DO NOT exceed maximum allowable rate.
Cutworms Tobacco Thrips Soybean (banded) Thrips	1.28 to 1.92 ounces (0.008 to 0.012 pound active) per acre	Respect EC may be applied in water or refined vegetable oil. When water is used, apply a minimum of one gallon of finished spray per acre by air or five gallons of finished spray with ground equipment. When applying in water by air, one quart of emulsified oil may be substituted for one quart of water in the finished spray. When using oil, use a minimum of one quart per acre in the finished spray. Control of lepidopteran eggs may be achieved with proper timing of applications. For boll weevil control, apply Respect EC at a 3- to 4-day interval until pest numbers are reduced to acceptable levels. For control of grasshoppers, applications should be made based on careful field scouting. Treatment decisions should be made based on evidence of feeding damage and presence of grasshoppers in cotton. Loss of cotyledon leaves in seedling cotton should be considered more important than leaf loss in older cotton. Applications should be made on a broadcast basis since grasshoppers are highly mobile.
Armyworm, Fall Armyworm, Yellow-striped Boll Weevil Cabbage Looper Corn Borer, European Cotton Bollworm Cotton Fleahopper Cotton Leaf Perforator Pink Bollworm Saltmarsh Caterpillar Stink Bugs Tarnished Plant Bug Other Plant Bugs Tobacco Budworm ¹	2.64 to 3.6 ounces (0.0165 to 0.0225 pound active) per acre	Adjust rates based on populations of grasshoppers found in fields. Applications should be made on a 3- to 5-day schedule until grasshopper populations are under control or until foliage loss subsides. Increase application rates as grasshopper size and population density increases. DO NOT exceed maximum allowable rate.
Armyworm, Beet ² Cotton Aphid ³ Lygus Bugs Whiteflies ⁴	2.8 to 4.0 ounces (0.0175 to 0.025 pound active) per acre	
Grasshoppers	3.0 to 4.0 ounces (0.01875 to 0.025 pound active) per acre	

A maximum of 24.0 fl ozs product (0.15 pound active ingredient) may be applied per acre per season.

DO NOT graze or feed cotton for forage.

¹See the resistance statement under the **Directions For Use** section.

²For control of beet armyworms only in the high plains of Texas, Arizona, and California

³Aphid control may be variable depending on species present and host-plant relationships.

⁴Aids in control

Fruiting Vegetables (except Cucurbits) **(1-day PHI) including: Eggplant; Groundcherry (*Physalis* spp.); Okra; Pepino (Melon Pear); Pepper** (includes Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper); **Tomatillo; Tomato**

Insects Controlled	Rate of Application	Method of Application
Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier Colorado Potato Beetle Corn Borer, European Corn Borer, Southwestern Corn Earworm Cucumber Beetle Cutworm spp. Flea Beetle Garden Webworm Green Stink Bug Hornworms Leafminers (adults) Leafhopper spp. Meadow Spittlebug Pepper Maggot (adults) Pepper Weevil Plant Bug spp. Tobacco Budworm ² Tomato Fruitworm Tomato Pinworm	2.24 to 4.0 ounces (0.014 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). DO NOT exceed maximum allowable rate.
Aphid spp. ^{2,3} Armyworm, Beet ² Armyworm, Fall Cabbage Looper Grasshoppers Lygus Bugs Brown Stink Bug Tomato Psyllid Thrips spp. ^{1,2} Whitefly spp. ^{1,2}	3.2 to 4.0 ounces (0.020 to 0.025 pound active) per acre	
<p>DO NOT make applications less than 7 days apart.</p> <p>DO NOT apply more than 24.0 fl ozs product (0.15 pound active ingredient) per acre per season.</p> <p>¹Aids in control</p> <p>²See the resistance statement under the Directions For Use section.</p> <p>³Aphid control may be variable depending on species present and host-plant relationships.</p>		

Legume Vegetables - Succulent and Dried (except Soybeans)
1-day PHI for succulent shelled or edible-podded peas or beans
21-day PHI for dried shelled peas or beans

Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas (*Pisum* spp.) including:
Dwarf Pea; Edible-Pod Pea; Snow Pea; Sugar Snap Pea; Pigeon Pea; English Pea; Garden Pea; Green Pea; Lentil

Succulent Edible-Podded Beans, Succulent Shelled Beans, and Dried Shelled Beans including:
Runner Bean; Snap Bean; Wax Bean; Asparagus Bean; Chinese Longbean; Moth Bean; Yardlong Bean; Jackbean; Soybean (immature seed); **Swordbean; Lima Bean; Broad Bean** (*Fava* Bean); **Blackeyed Pea; Southern Pea; Grain Lupin; Sweet Lupin; White Lupin; White Sweet Lupin; Field Bean; Kidney Bean; Navy Bean; Pinto Bean; Tepary Bean; Adzuki Bean; Catjang; Cowpea; Crowder Pea; Moth Bean; Mung Bean; Rice Bean; Urd Bean; Chickpea** (*Garbanzo* Bean); **Guar; Lablab Bean**

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Saltmarsh Caterpillar Silverspotted Skipper Thistle Caterpillar (Painted Lady)	1.28 to 4.0 ounces (0.008 to 0.025 pound active) per acre	Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
Alfalfa Caterpillar Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Bean Leaf Beetle Blister Beetle spp. Colorado Potato Beetle Corn Borer, European Corn Borer, Southwestern Corn Earworm Corn Rootworm Beetle (adult) Cowpea Curculio Cucumber Beetle Flea Beetle Green Cloverworm Ground Beetles Imported Cabbageworm Japanese Beetle Leaf Skeletonizer spp. Leafhopper spp. Leafminers (adults) Mexican Bean Beetle Pea Weevil Pea Leaf Weevil Plant Bug spp. Potato Leafhopper Seedcorn Beetle Seedcorn Maggot (adult) Spittlebug Three-cornered Alfalfa Hopper Tobacco Budworm ² Velvetbean Caterpillar Webworm spp. Woolly Bear Caterpillar	2.72 to 4.0 ounces (0.017 to 0.025 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). DO NOT exceed maximum allowable rate.
Aphid spp. ^{2,3} Armyworm, Beet ² Armyworm, Fall Grasshoppers Lesser Cornstalk Borer ¹ Looper spp. ² Stink Bug spp. Thrips spp. ^{1,2} Whitefly spp. ^{1,2}	3.2 to 4.0 ounces (0.020 to 0.025 pound active) per acre	

Legume Vegetables - Succulent and Dried (except Soybeans)

Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas (*Pisum* spp.)

Succulent Edible-Podded Beans, Succulent Shelled Beans, and Dried Shelled Beans (continued)

DO NOT make applications less than 5 days apart.

DO NOT apply more than 24.0 fl ozs product (0.15 pound active ingredient) per acre per season.

¹Aids in control

²See the resistance statement under the **Directions For Use** section.

³Aphid control may be variable depending on species present and host-plant relationships.

Peanut (7-day PHI)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Green Cloverworm Velvetbean Caterpillar Red-necked Peanut Worm	1.28 to 4.0 ounces (0.008 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels.
Bean Leaf Beetle Leafhopper spp. Southern Corn Rootworm (adult) Vegetable Weevil Whitefringed Beetle (adult)	1.76 to 4.0 ounces (0.011 to 0.025 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). DO NOT make applications less than 14 days apart. DO NOT exceed maximum allowable rate.
Aphid spp. ^{1,2} Armyworm, Beet ^{1,2} Armyworm, Fall ^{1,2} Corn Earworm Grasshopper spp. Lesser Cornstalk Borer ^{1,2} Soybean Looper ^{1,2} Stink Bug spp. ^{1,2} Tobacco Thrips ²	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	
<p>DO NOT apply more than 24.0 fl ozs product (0.15 pound active ingredient) per acre per season.</p> <p>DO NOT graze livestock in treated areas. DO NOT use treated vines or hay for animal feed.</p> <p>¹ Aids in control.</p> <p>² See the resistance statement under the Directions For Use section.</p>		

Root and Tuber Vegetables Group 1 (except Sugar Beet) (1 day PHI) including: **Arracacha; Arrowroot; Artichoke** (Chinese and Jerusalem); **Garden Beet; Edible Burdock; Edible Canna; Carrot; Cassava** (Bitter and Sweet); **Celeriac** (Celery Root); **Chayote** (Root); **Turnip-Rooted Chervil; Chicory; Chufa; Dasheen** (Taro); **Ginger; Ginseng; Horseradish; Leren; Turnip-Rooted Parsley; Parsnip; Potato; Oriental Radish** (Daikon); **Radish; Rutabaga; Salsify** (Oyster Plant); **Black Salsify; Spanish Salsify; Skirret; Sweet Potato; Tanier** (Cocoyam); **Turmeric; Turnip; Yam Bean; and Yam** (True)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp.	1.28 to 4.0 ounces (0.008 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels.
Cabbage Looper Cucumber Beetle European Corn Borer Flea beetle spp. Leafhopper spp. Southern Corn Rootworm (adult) Vegetable Weevil Whitefringed Beetle (adult)	1.76 to 4.0 ounces (0.011 to 0.025 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). DO NOT make applications less than 4 days apart. DO NOT exceed maximum allowable rate.
Aphid spp. ^{1,2} Armyworm, Beet ^{1,2} Armyworm, Yellowstriped Cabbage Maggot Colorado Potato Beetle ² Grasshopper spp. Imported Cabbageworm Potato Leafhopper Tarnished Plant Bug	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	
<p>DO NOT apply more than 24.0 fl ozs product (0.15 pound active ingredient) per acre per season.</p> <p>Leaves of root and tuber vegetables cannot be used for food or feed.</p> <p>¹ Aids in control</p> <p>² See the resistance statement under the Directions For Use section.</p>		

Sorghum (Grain) and Millet (14-day PHI for grain and stover; 45-day PHI for forage)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Sorghum Midge	1.28 to 4.0 ounces (0.008 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
Armyworm, Fall Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Corn Borer, European ¹ Corn Borer, Southwestern ¹ Corn Earworm Flea Beetle spp. Hornworms Stink Bug spp. Webworm spp.	1.76 to 4.0 ounces (0.011 to 0.025 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). The addition of 1 to 2 quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control. For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 10-day intervals if needed. For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of plants with sufficient spray volume to penetrate the soil/stem interface, leaf collars, and sheaths.
Aphid spp. ^{2,3} Armyworm, Beet ³ Chinch Bug False Chinch Bug Grasshopper spp. Lesser Cornstalk Borer ¹ Thrips spp. ^{3,4} Whitefly spp. ^{3,4}	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	DO NOT exceed maximum allowable rate.
<p>DO NOT make applications less than 10 days apart.</p> <p>DO NOT apply more than 20.0 fl ozs product (0.125 pound active ingredient) per acre per season.</p> <p>¹For control before the larva bores into the plant stalk</p> <p>²Aphid control may be variable depending on species present and host-plant relationships.</p> <p>³See the resistance statement under the Directions For Use section.</p> <p>⁴Aids in control</p>		

Soybeans (21-day PHI)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Silverspotted Skipper	1.28 to 4.0 ounces (0.008 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
Alfalfa Caterpillar Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Bean Leaf Beetle ¹ Blister Beetle spp. Colorado Potato Beetle Corn Borer, European Corn Earworm Corn Rootworm Beetle (adult) Cowpea Curculio Cucumber Beetle European Corn Borer Flea Beetle Green Cloverworm Hornworms Imported Cabbageworm Japanese Beetle Leaf Skeletonizer spp. Leafhopper spp. Leafminers (adults) Mexican Bean Beetle Pea Leaf Weevil Plant Bug spp. Potato Leafhopper Seedcorn Maggot (adult) Soybean Aphid Spittlebug Three-cornered Alfalfa Hopper Tobacco Budworm ² Velvetbean Caterpillar Webworm spp. Woollybear Caterpillar	2.8 to 4.0 ounces (0.0175 to 0.025 pound active) per acre	Apply with either aerial or ground equipment using sufficient spray volume to obtain full coverage of the plant and foliage. Use a minimum of 2 gallons of finished spray per acre by air or 10 gallons of finished spray per acre by ground. The addition of one to two quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control. DO NOT exceed maximum allowable rate.
Armyworm, Beet Armyworm, Fall Grasshopper spp. Lesser Cornstalk Borer ³ Looper spp. ² Stink Bug spp. Thrips spp. ^{2,3} Whitefly spp. ^{2,3}	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	

DO NOT make applications less than 7 days apart.

DO NOT graze or harvest treated soybean forage, straw, or hay for livestock feed.

DO NOT apply more than 24.0 fl ozs product (0.15 pound active ingredient) per acre per season.

¹Use higher recommended dosage for increased pest pressure, increased residual pest control, or later-season applications.

²See the resistance statement under the **Directions For Use** section.

³Aids in control

Sugarcane (21-day PHI)

Insects Controlled	Rate of Application	Method of Application
Sugarcane Borer Mexican Rice Borer	3.0 to 4.0 ounces (0.01875 to 0.025 pound active) per acre	Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. Apply by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground). DO NOT exceed maximum allowable rate.
DO NOT make applications less than 21 days apart.		
DO NOT apply more than 16.0 fl ozs product (0.10 pound active ingredient) per acre per season.		

Tree Nuts (7 Day PHI) including: almond; beech nut; Brazil nut; butternut; cashew; chestnut; chinquapin; filbert (hazelnut); hickory nut; macadamia nut; pecan; and walnut (black and English)

Insects Controlled	Rate of Application	Method of Application
Black Pecan Aphid Codling Moth Filbert Worm Hickory Shuckworm Leaffooted Bugs Navel Orangeworm Oblique-banded Leafroller Peach Twig Borer Pecan Leaf Casebearer Pecan Nut Casebearer Pecan Phylloxera Pecan Weevil Plant Bugs Stink Bugs Walnut Aphid Walnut Husk Fly Yellow Pecan Aphid	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). DO NOT exceed maximum allowable rate.
DO NOT apply more than 20.0 fl ozs product (0.125 pound active ingredient) per acre per season.		
DO NOT make applications less than 7 days apart.		

Wheat and Triticale (14-day PHI for grain, forage, and hay)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp., including Army Cutworm Painted Lady (Thistle) Caterpillar	1.28 to 4.0 ounces (0.008 to 0.025 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Cereal Leaf Beetle Flea Beetle spp. Pale Western Cutworm Plant Bug spp. Spittlebug Webworm spp.	1.76 to 4.0 ounces (0.011 to 0.025 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). For chinch bug control, begin applications when bugs migrate from small grains or grass weeds. Apply sufficient spray volume to penetrate the soil/stem interface, leaf collars, and sheaths. DO NOT exceed maximum allowable rate.
Aphid spp. ^{1,2} Armyworm, Beet ² Armyworm, Fall Chinch Bug Grass Sawfly Grasshopper spp. Greenbug ^{2,3} Stink Bug spp. Thrips spp. ^{2,3} Wheat Stem Sawfly (adult) ³ Whitefly spp. ^{2,3}	3.2 to 4.0 ounces (0.02 to 0.025 pound active) per acre	
<p>DO NOT make applications less than 14 days apart.</p> <p>DO NOT apply more than 20.0 fl ozs product (0.125 pound active ingredient) per acre per season.</p> <p>¹Aphid control may be variable depending on species present and host-plant relationships.</p> <p>²See the resistance statement under the Directions For Use section.</p> <p>³Aids in control</p>		

Dealers Must Sell in Original Packages Only.

Conditions of Sale and Limitation of Warranty and Liability

NOTICE: Read the entire **Directions for Use** and **Conditions of Sale and Limitation of Warranty and Liability** before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The **Directions for Use** of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of BASF or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold BASF and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the **Directions for Use** when used in accordance with the directions under normal conditions of use. BASF MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or BASF, and buyer assumes the risk of any such use.

In no event shall BASF or seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF BASF AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF BASF OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This **Conditions of Sale and Limitation of Warranty and Liability** may not be amended by any oral or written agreement.

Respect is a registered trademark of BASF.

Ammo, Capture, Fury, Hero, and Mustang are trademarks of FMC Corporation.

Asana is a trademark of E.I. duPont de Nemours & Company.

Ambush and Karate are trademarks of Zeneca, LTD.

Baythroid is a trademark of Bayer Aktiengesellschaft.

Danitol is a trademark of Sumitomo Chemical Company, LTD.

Scout is a trademark of Hoechst Schering Agrevo S.A.

© 2009 BASF Corporation
All rights reserved.

000279-03327-07969.20090401.NVA 2008-04-297-0320
Supersedes: NVA 2008-04-297-0230

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company