

# SUPPLEMENTAL LABELING

**DuPont Crop  
Protection**

**DUPONT™ CORAGEN® INSECT  
CONTROL WITH ACTIVE  
INGREDIENT RYNAXYPYR®  
FOR USE ON COTTON**

## **DUPONT™ CORAGEN® INSECTICIDE**

EPA Reg. No. 352-729

### **FOR FOLIAR INSECT CONTROL ON COTTON**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **GENERAL INFORMATION**

CORAGEN™ is a suspension concentration that can be applied as a foliar spray on cotton to control or suppress the important cotton insect pests listed in the table below. CORAGEN™ is mixed with water for application.

#### **Resistance Management**

For resistance management, CORAGEN® is a Group 28 Insecticide. Repeated and exclusive use of CORAGEN® or other Group 28 Insecticides may lead to the buildup of resistant strains of insects in some crops.

Unless directed otherwise in the specific crop/pest sections of this label, the best practices are to follow these instructions to delay the development of insecticide resistance: Avoid using the same mode of action (same insecticide group) on consecutive generations of insect pests. Make no more than 2 applications of CORAGEN® per generation to the same insect species on a crop. Application(s) to the next generation of target pest(s) must be with an effective product with a different mode of action. Make no more than 2 successive applications within a 30 day period to the same insect species on a crop. The following application to the target pest(s) must be with an effective product with a different mode of action.

See label accompanying product for full resistance management instructions.

#### **APPLICATION**

Apply follow-up treatments of DuPont™ CORAGEN®, as specified in the table on page 2 of this label, to keep pest populations within threshold limits. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area. Refer to the Resistance Management section of the label accompanying product for further guidance on follow-up treatments.

Use sufficient water to obtain thorough, uniform coverage.

#### **IMPORTANT**

Before using CORAGEN®, read and follow all applicable directions; restrictions; and precautions on the EPA-registered label. This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

(See table on Page 2)

R-957 032709-2 03-17-09

© 2009 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

Crop	Application Method	Target Pest	DUPONT™ CORAGEN™ RATE		Last Application (Days to Harvest)	REI (Hours)
			Lb. ai per acre	fluid ounces product per acre		
Cotton	FOLIAR	Beet armyworm Cotton bollworm** Fall armyworm Saltmarsh caterpillar Southern armyworm Tobacco budworm**	0.045 – 0.09	3.5– 7.0	21	4 hr.
		Cabbage looper Soybean looper*	0.065 – 0.098	5.0 – 7.5		
<p>Do not apply more than 15.4 fl oz CORAGEN™ (0.2 lbs. a.i.) per acre per crop per season. The minimum interval between treatments is 5 days.</p> <p>*Suppression only.</p> <p>** For Heliothine control (cotton bollworm and/or tobacco budworm) make the first application at rates of 0.065 - 0.09 lb. ai per acre (5.0 - 7.0 oz product). Subsequent applications can be at rates of 0.045 - 0.09 lb. ai acre (3.5 - 7.0 oz product) depending on pest pressure.</p>						