

Supplemental Labeling

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification



EPA Reg. No. 279-3332

EPA Est. No. 279-NY-1

Active Ingredient:

By Wt.

Bifenthrin: (2 methyl [1,1'-biphenyl]-3-yl)
Methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2, 2-dimethyl-
cyclopropanecarboxylate*11.30%
Imidacloprid: 1-[(6-Chloro-3-pyridonyl)
methyl]-N-nitro -2-Imidazolidinimine11.30%

Other Ingredients:77.40%

100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.
This product contains 2lbs. active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

This label must be in the possession of the user at the time of application. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Brigadier Insecticide before applying. Carefully follow all precautionary statements and application use directions. Use of Brigadier Insecticide according to this supplemental labeling is subject to all use precautions and limitations imposed by the labeling affixed to the container for Brigadier Insecticide

Potato (PHI 21 days) (Foliar uses)

Pests Controlled	Rates of Application
Foliar Application	
Aphid spp. Leafhopper spp.	3.8 – 6.14 fl oz/A (0.06 – 0.096 lbs ai/A)
Banded Cucumber beetle Colorado potato beetle Cucumber beetle European corn borer Grasshopper spp. Looper spp. Flea beetle spp. June beetle Potato psyllid Sugarcane beetle Sweetpotato flea beetle Sweetpotato weevil Tuberworm Whitefringed beetle Whitefly	4.8 – 6.14 fl oz/A (0.075 – 0.096 lbs ai/A)
Restrictions <ul style="list-style-type: none"> • PHI: Do not apply within 21 days of harvest. • Application Interval: Do not make applications less than 7 days apart. • Maximum Amount of Brigadier Insecticide per Foliar Application: 6.14 fl oz/A (0.048 lb ai/a of imidacloprid and 0.048 lb ai/a of bifenthrin) as a foliar application. • Maximum Amount of Brigadier Insecticide per Season: 25.6 fl oz/A (0.2 lb ai/a of imidacloprid and 0.2lb ai/a of bifenthrin) as a foliar application. Two applications are permitted per season. It is permitted to make one at-plant application followed by a foliar application later in the same growing season. • Maximum Amount of imidacloprid per Season: 0.20 lbs ai/A. • Maximum Amount of bifenthrin per Season: 0.50 lbs ai/A Remarks Foliar Application: Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray. Thorough coverage is essential to achieve control.	

Potato (At-plant)

Pests Controlled	Rates of Application
At-Plant Application	
Aphid spp. Colorado potato beetle Flea beetle spp. (adult) Flea beetle spp. (larvae) Japanese beetle (larvae) Leaf hopper spp. Potato psyllid Rootworm spp. White grub spp. Wireworm spp.	16 – 25.6 fl oz/A (0.25 – 0.6 lbs ai/A)
Restrictions <ul style="list-style-type: none"> • PHI: Do not apply within 21 days of harvest. • Application Interval: Do not make applications less than 7 days apart. • Maximum Amount of Brigadier Insecticide per Application: 25.6 fl oz/A (0.2 lb ai/a of imidacloprid and 0.2 lb ai/a bifenthrin) as an at-plant application • Maximum Amount of bifenthrin per Season: 0.30 lbs ai/A as an at-plant application. A total 0.50 lbs ai/A of bifenthrin per season. • Maximum Amount of imidacloprid per Season: 0.20 lbs ai/A. A maximum of 1 at plant applications is permitted per season.	
Remarks At plant applications: In-furrow applications: Apply Brigadier Insecticide as an in-furrow spray on to the seed pieces or seed potatoes	



FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia PA 19103