

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.



2(ee) Recommendation

FOR CONTROL OF KUDZU BUG OR BEAN PLATASPID ON SOYBEANS

Not for use in AK, DC, HI, IN, NH, NM, NY & TX

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

EPA Registration No. 279-3332

CROP	PEST	RATE OF APPLICATION
Soybean	Kudzu bug or Bean Plataspid	6.1 oz/A
<p>DIRECTION FOR USE: Apply 6.1 oz/A BRIGADIER insecticide by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). Thorough coverage is essential to achieve control</p>		
<p>RESTRICTIONS: Do not apply more than 6.1 fluid ounces of product per acre per application. Do not apply more than 17.92 fluid ounces of product per acre per season. Do not sequentially apply BRIGADIER insecticide less than 30 days apart. Do not harvest soybeans within 45 days of a BRIGADIER application for feeding of dry vines. Do not harvest soybeans within 18 days of a BRIGADIER application for feeding of green vines.</p>		
<p>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED BRIGADIER INSECTICIDE LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</p>		

FMC
 FMC Corporation
 Agricultural Products Group
 1735 Market Street
 Philadelphia, PA 19103

Brigadier and FMC - FMC Trademarks
 ©2012 FMC Corporation
 013112