



DuPont™ Resolve® Q

HERBICIDE

GROUP 2 HERBICIDE

For use in Fallow and Field Corn grown for grain or silage

<i>Active Ingredients</i>	<i>By Weight</i>
Rimsulfuron	
N-((4,6-dimethoxypyrimidin-2-yl)aminocarbonyl)-3-(ethylsulfonyl)-2-pyridinesulfonamide	18.4%
Thifensulfuron-methyl	
Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	4.0%
Other Ingredients	77.6%
TOTAL	100.0%

EPA Reg. No. 352-777

EPA Est. No. _____

Nonrefillable Container

Net: _____

OR

Refillable Container

Net: _____

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

Long-sleeve shirt and long pants.

Chemical resistant gloves made of any waterproof material such as nitrile rubber, natural rubber, neoprene rubber, or butyl rubber.

Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

DuPont™ RESOLVE® Q herbicide, also referred to below as RESOLVE® Q or DuPont™ RESOLVE® Q, must be used in accordance with the directions for use on this label; in separately issued labeling or exemptions under FIFRA (Supplemental Labels; Special Local Need Registrations; FIFRA Section 18 exemptions; or as otherwise permitted by FIFRA. Always read the entire label including the Limitation of Warranty and Liability.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical resistant gloves made of any waterproof material such as nitrile rubber, natural rubber, neoprene rubber, or butyl rubber.

Shoes plus socks.

PRODUCT INFORMATION

RESOLVE® Q must be used only in accordance with instructions on this label or in supplemental DuPont publications. DuPont will not be responsible for losses or damage resulting from use of this product in any manner not specified by DuPont.

RESOLVE® Q herbicide is a water soluble granule which is a selective herbicide for burndown and residual control of certain annual grass and broadleaf weeds.

RESOLVE® Q can be tank mixed with a variety of herbicides to improve burndown and residual control; however, the most restrictive label must be followed.

RESOLVE® Q is absorbed through the roots and leaf tissue of plants, rapidly inhibiting the growth of susceptible weeds. Rainfall or sprinkler irrigation is needed to move RESOLVE® Q into the soil. Susceptible weeds will generally not emerge from a preemergence application. In some cases, susceptible weeds may germinate and emerge a few days after application, but growth then ceases and leaves become chlorotic three to five days after emergence. Death of leaf tissue and growing point will follow in some species, while others will remain green, stunted and noncompetitive.

The herbicidal action of RESOLVE® Q may be less effective on weeds stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions or cultural practices.

RESOLVE® Q residual is most effective in controlling weeds when adequate rainfall is received within 5-7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain occurs, use shallow tillage such as rotary hoe to lightly incorporate RESOLVE® Q and make certain corn seeds are below the tilled area.

DuPont™ RESOLVE® Q is best used in a planned sequential application herbicide program, following a preplant/preemergence application of DuPont™ ALLUVEX™, DuPont™ BASIS® Blend, DuPont™ CINCH®, DuPont™ BREAKFREE®, DuPont™ INSTIGATE®, DuPont™ LEADOFF® or DuPont™ PREQUEL® brands, RESOLVE® Q, RESOLVE® SG, and/or other pre-applied corn herbicides. Refer to the label of the respective corn herbicide partner for specific use directions.

For postemergence applications of RESOLVE® Q, if activating rainfall or sprinkler irrigation (>0.5 inch) is not received after application, follow with a cultivation or with a sequential application of DuPont™ ACCENT® Q herbicide, DuPont™ REVULIN™ Q herbicide, DuPont™ STEADFAST® Q, or glyphosate such as ABUNDIT® Extra as needed.

RESOLVE® Q is rainfast in 4 hours.

RESTRICTIONS

Do not apply to popcorn or sweet corn.

Do not apply preemergence or postemergence to seed corn.

Do not apply more than 1.0 oz active ingredient rimsulfuron per acre per year. This includes combinations of fallow, preplant, preemergence and postemergence applications of RESOLVE® Q, as well as rimsulfuron from applications of products such as ALLUVEX™, BASIS® Blend, INSTIGATE®, LEADOFF®, PREQUEL®, RESOLVE® SG and DuPont™ STEADFAST® Q.

Do not use preemergence rates of RESOLVE® Q greater than 1.25 oz product per acre if following with postemergence applications of the rimsulfuron containing product noted above.

Do not apply more than 1.25 ounces of RESOLVE® Q postemergence, per acre per application to field corn, unless instructed to do so by DuPont technical bulletins, fact sheets, or supplemental labeling.

Do not apply to coarse-textured soils (sand, loamy sand or sandy loam) with less than 1% organic matter.

Do not tank mix RESOLVE® Q with “Basagran” or severe crop injury may occur.

Do not tank mix RESOLVE® Q with foliar-applied organophosphate insecticides such as chlorpyrifos (“Lorsban”), malathion, etc, as severe crop injury may occur. To avoid crop injury, apply these products at least 7 days before or 3 days after the application of RESOLVE® Q.

Do not apply the organophosphate insecticide terbufos ("Counter") within 45 days of a preplant or preemergence application of RESOLVE® Q since crop injury may result.

Do not apply RESOLVE® Q within 45 days of crop emergence where the organophosphate insecticide, terbufos ("Counter") was applied since crop injury may occur. Applications made to corn previously treated with chlorpyrifos or other similar organophosphate insecticides may result in unacceptable crop injury.

Do not apply this product through any irrigation system.

Do not use flood or furrow irrigation to apply RESOLVE® Q.

Do not irrigate RESOLVE® Q into coarse soils at planting time when soils are saturated.

Do not graze, feed forage, grain or fodder (stover) from treated areas to livestock within 30 days of RESOLVE® Q application.

Injury or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply RESOLVE® Q or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas
- Prevent drift or spray onto desirable plants
- Do not contaminate any body of water

PRECAUTIONS

Allow at least 4 weeks between preemergence application of RESOLVE® Q and postemergence applications of unsafened rimsulfuron-containing herbicides such as INSTIGATE®.

RESOLVE® Q may interact with certain insecticides previously applied to the crop. Crop response varies with field crop, insecticide used, insecticide application methods, and soil type.

RESOLVE® Q may be applied to corn previously treated with “Fortress”, “SmartChoice”, “Aztec”, or “Force” insecticides, or nonorganophosphate soil insecticides regardless of soil type.

RESOLVE® Q may be applied with pyrethroid type insecticides such as “Asana” or “Warrior” or with diamide type insecticides such as DuPont™ PREVATHON®.

Preplant/Preemergence applications of RESOLVE® Q to field crops where an application of “Lorsban” or “Thimet” is planned may cause unacceptable crop injury, especially on soils of less than 4% organic matter.

Thoroughly clean application equipment immediately after use (See Sprayer Preparation/Cleanup section of this label).

Crop injury may occur following an application of DuPont™ RESOLVE® Q if there is a prolonged period of cold weather and / or in conjunction with wet soils.

Prevent drift or spray to desirable plants.

RESISTANCE MANAGEMENT

RESOLVE® Q, which contains the active ingredients rimsulfuron and thifensulfuron-methyl, is a Group 2 herbicide based on the mode of action classification system of the Weed Science Society of America.

When herbicides with mode of action classifications that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

APPLICATION INFORMATION

Fallow

Rate

Apply RESOLVE® Q at 1.0 - 2.5 ounces per acre.

Timing to Crop & Weeds

RESOLVE® Q may be used as a fallow treatment, in the fall or spring when the majority of weeds have emerged and are actively growing.

Tank Mixtures

RESOLVE® Q may be used as a fallow treatment and may be tank mixed with other herbicides that are registered for use in fallow such as ABUNDIT® Extra, DuPont™ EXPRESS®, DuPont™ PANOFLEX™, glyphosate, paraquat, glufosinate, saflufenacil (Sharpen), 2,4-D LVE, and dicamba herbicides for improved control of emerged weed species. Read and follow all instructions on this label and the labels of any tank mix partner before using any other herbicide in mixtures with RESOLVE® Q. If the instructions on the tank mix partner label conflict with this RESOLVE® Q label, do not use in a tank mixture with RESOLVE® Q.

Field Corn Grown for Grain or Silage - Preplant/Preemergence

Rate

Apply RESOLVE® Q at 1.25 – 2.5 ounce per acre before corn emergence. See cumulative rimsulfuron rate limitation noted in Product Information. RESOLVE® Q at 1.25 – 1.5 ounce per acre fits most preemergence/preplant applications.

Timing to Crop

RESOLVE® Q herbicide may be used in conventional, conservation tillage, or no-till crop management systems and may be applied either preplant, preplant incorporated (less than 2" deep), or preemergence for use in field corn production. Applications of RESOLVE® Q made before weed emergence will provide residual control of labeled weeds. Control of emerged weeds will require the addition of spray adjuvants, and can be further enhanced with additional tank mix partners as noted in this label.

Preplant Incorporated: Apply to the soil and uniformly incorporate in the top two inches of soil before planting using a finishing disc harrow, field cultivator or similar implement capable of providing uniform two inch incorporation. Do not incorporate RESOLVE® Q deeper than 2" or weed control may be reduced.

Preplant/Preemergence Burndown: Apply DuPont™ RESOLVE® Q when weeds are young and actively growing.

The addition of crop oil concentrate or methylated seed oil is recommended for burndown of labeled weeds. When weeds are greater than the maximum height listed or weeds not controlled by RESOLVE® Q are present, the addition of a burndown herbicide such as glyphosate, glufosinate, paraquat, dicamba, and/or 2, 4-D is recommended. If giant ragweed, common cocklebur, henbit, Pennsylvania smartweed or purple deadnettle are present at the time of application, the addition of atrazine will improve control. Observe direction for use and precaution and restrictions on the label of the burndown herbicide. When mixing with liquid nitrogen fertilizer or glyphosate, substitute a non-ionic surfactant for crop oil.

Preemergence: Apply RESOLVE® Q herbicide during planting (behind the planter after furrow closure) or after planting.

Sequential Application - Preemergence

RESOLVE® Q may be used as a sequential application in a planned postemergence weed control program in corn following a preemergence herbicide.

Apply pre products such as atrazine, DuPont™ BASIS® Blend, RESOLVE® Q, DuPont™ CINCH® or DuPont™ BREAKFREE® brands, DuPont™ INSTIGATE®, DuPont™ LEADOFF® or DuPont™ PREQUEL® herbicides. Refer to the preemergence herbicide label for use restrictions, application information, rotational crop guidelines, and cautionary statements prior to applying RESOLVE® Q.

Restriction: Do not apply RESOLVE® Q to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

Field Corn Grown for Grain or Silage – Postemergence

Rate

Apply RESOLVE® Q at 1.25 ounces per acres as a postemergence broadcast application.

Timing to Crop

Apply RESOLVE® Q to corn that is up to 20 inches tall. Do not apply to corn taller than 20 inches or exhibiting 7 or more leaf collars, whichever is more restrictive. While RESOLVE® Q has a wide application window, research has shown best results are obtained when applications are made early postemergence when corn and weeds are small. Target applications to corn that is less than 12" tall for best overall performance.

Applications of RESOLVE® Q made after weed emergence will provide contact control of labeled weeds as well as limited residual control of later emerging weeds.

Timing to Emerged Weeds

Apply RESOLVE® Q when grasses are young and actively growing, but before they exceed the sizes listed on this label.

On "Roundup Ready" corn, glyphosate may be applied with RESOLVE® Q after weeds emerge but before they reach the maximum size listed on the glyphosate herbicide label.

On "Liberty Link" corn, glufosinate may be applied with RESOLVE® Q after weeds emerge but before they reach the maximum size listed on the glufosinate herbicide label.

Applications made to weed sizes greater than those listed on these product labels may result in incomplete control. Grass competition due to incomplete control may reduce yields.

Sequential Application - Postemergence

Apply DuPont™ ACCENT® Q herbicide or DuPont™ REVULIN™ Q herbicide 14 or more days after the RESOLVE® Q application to control grasses that may emerge later in the season. Refer to the ACCENT® Q or REVULIN™ Q label for weeds controlled, proper size of weeds, rates, corn sizes, and other information. When following a RESOLVE® Q application, do not use more than 0.9 ounce per acre of ACCENT® Q or 3.4 ounces per acre of REVULIN™ Q.

Spray Adjuvants

For control of emerged weeds, application of RESOLVE® Q must include a crop oil concentrate, modified seed oil or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer must be used unless specifically prohibited by the tankmix partner labeling. Crop oil concentrate/modified seed oil plus ammonium nitrogen fertilizer is the preferred adjuvant system for RESOLVE® Q for control of emerged weeds. When applied in tank mix combination with a glyphosate or glufosinate herbicide that contains a built-in adjuvant system, ensure the total adjuvant load is equivalent to the recommendations on this label. Select adjuvants authorized for use with both products.

Consult local DuPont fact sheets, technical bulletins, and service policies prior to using other adjuvant systems. Products must contain only EPA-exempt ingredients.

Do not use with spray additives that alter the pH of the spray solution below 5.0 or above 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 – 8.0 allow for optimum stability of RESOLVE® Q.

Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)

- Apply at 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- MSO adjuvants may be used at 0.5% v/v (0.5 gallon per 100 gallons spray solution) if specifically noted on adjuvant product labeling.

- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Nonionic Surfactant (NIS)

- Apply at 0.25% v/v (1 qt per 100 gal spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

Ammonium Nitrogen Fertilizer

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN) such as 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS).
- Do not use liquid nitrogen fertilizer as the total carrier solution after crop emergence.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS and ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.

Weeds Controlled/Suppressed

For best control of emerged weeds, apply RESOLVE® Q to grasses 3 inches or less, broadleaf weeds 4 inches or less and winter annuals/biennials 6 inches or less. Make application prior to flowering.

Refer to the Spray Adjuvants section for additional information on proper adjuvant selection.

Broadleaf & Grass Weeds	Burndown	
	DuPont™ RESOLVE® Q Alone	Residual - RESOLVE® Q
Alfalfa, volunteer	C	NC
Barley, volunteer	C	S
Barnyardgrass	C	C
Bittercress	C	C
Bluegrass, annual	C	C
Brome, downy	S	S
Buckwheat, common	C	NC
Buttercup, smallflower	C	NC
Canada thistle	S	NC
Catchweed bedstraw	C	C
Chamomile, false	NC	C
Chickweed (common, mouseear)	C	NC
Cocklebur	S	S
Crabgrass, large	C ¹	S
Cupgrass, woolly (1")	C	NC
Curly Dock	C	NC
Dandelion (6" diameter)	C	NC
Deadnettle, purple	C	C
Eveningprimrose, cutleaf	C ²	NC
Fescue, tall	S	S
Field pennycress	C	NC
Filaree, redstem	NC	C
Foxtail (bristly, giant, green, yellow)	C	C
Foxtail, Carolina	C	C
Geranium, Carolina	C	NC
Groundsel, common	C	NC
Hemlock, poison (up to 12")	C	S
Henbit	C	C
Knotweed, prostrate	C	NC
Johnsongrass, seedling	S	NC
Kochia	C ³	C ³
Lambsquarters, common	C	C
Marestail (Horseweed)	S ³	C ³
Millet, wild proso	S	NC
Morningglory, ivyleaf	S	S
Mustard (birdsrape, black)	C	C
Mustard, wild	C	NC
Nightshade, hairy	S	S
Panicum, fall	C	S
Parsnip, wild	C	S
Pigweed (prostrate, redroot, smooth)	C ⁴	C
Purslane, common	S	C
Quackgrass	S	NC
Ragweed, common	S	S
Ryegrass, Italian	S ⁴	S ⁴
Shattercane (4")	C	NC
Shepherd's purse	C	NC
Signalgrass, broadleaf	S	C
Smartweed, Pennsylvania	C	S
Smartweed, ladysthumb	C	NC
Stinkgrass	S	NC
Sunflower	C	S
Velvetleaf	C	S
Wallflower, bushy	C	NC
Wheat, volunteer	C	C
Wild oat	S	S
Wild radish	C	NC
Yellow nutsedge	S	NC
Yellow rocket	C	C

C= Control

S= Suppression

NC = No Control

1 = <1/2"

2 = Must add 2,4D LVE or dicamba for control

3 = ALS Sensitive

4 = Resistant biotypes are known to occur

For full season control utilizing a two pass application program (pre followed by post to corn), follow the preemergence application of DuPont™ RESOLVE® Q with a sequential, in-crop application of RESOLVE® Q, DuPont™ REVULIN™ Q or DuPont™ STEADFAST® Q with appropriate tank mix partners.

For full season control utilizing a one pass preemergence application, mix RESOLVE® Q with atrazine-containing grass and broadleaf corn herbicides, such as DuPont™ CINCH®, or DuPont™ BREAKFREE® brands. Depending on the growing season, in-crop post applications may be needed to control late grass and weed escapes.

See Tank Mixtures section of this label for additional information.

When using multiple rimsulfuron-based products in a cropping season, observe 1.0 oz active ingredient per acre limit.

Consult local DuPont representative, fact sheets, technical bulletins, or supplemental labels for additional information.

Postemergence – RESOLVE® Q 1.25 oz/acre with Glyphosate

Glyphosate such as ABUNDIT® Extra may be tank mixed with postemergent applications of RESOLVE® Q when made to corn hybrids containing the "Roundup Ready" gene.

Refer to the Spray Adjuvants section for additional information on proper adjuvant selection.

When used in tank mixture with glyphosate, RESOLVE® Q will deliver improved burndown and/or residual activity on the following weeds:

Alfalfa, volunteer	Nightshade, hairy
Barley, volunteer	Panicum, fall
Barnyardgrass	Pigweed (prostrate, redroot, smooth)
Bluegrass, annual	Purslane, common
Canada thistle	Quackgrass
Chamomile, false	Ragweed, common
Chickweed, common	Ryegrass, Italian
Cocklebur	Sandbur (field, longspine)
Crabgrass, large	Shepherd's purse
Dandelion (6" diameter)	Signalgrass, broadleaf
Filaree, redstem	Smartweed, Pennsylvania
Foxtail (bristly, giant, green, yellow)	Stinkgrass
Henbit	Velvetleaf
Johnsongrass, seedling	Wheat, volunteer
Kochia	Wild buckwheat
Lambsquarters, common	Wild oat
Millet, Wild Proso	Wild radish
Morningglory, ivyleaf	Yellow Nutsedge
Mustard (birdsrape, black, wild)	

DuPont™ RESOLVE® Q 1.25 oz/acre with Glufosinate

RESOLVE® Q may be tank mixed with glufosinate herbicide if applications are made to corn hybrids containing the "Liberty Link" gene. Consult with your seed supplier to confirm the corn hybrid is "Liberty Link" before applying any herbicide containing glufosinate.

When used in a tank mixture with glufosinate herbicide, RESOLVE® Q will deliver improved burndown and/or limited residual activity on the following weeds:

Velvetleaf
Pigweed, redroot
Lambsquarters, common
Foxtail (giant, yellow)

Tank Mixtures - Additional Control of Broadleaf and Grass Weeds

RESOLVE® Q may be tank mixed with other products registered for use in corn. Consult tank mix partner labeling for rate and soil-type restrictions. Read and follow all manufacturer's label instructions for the companion herbicide(s). Do not use a tank mix partner product if its label conflicts with the RESOLVE® Q label.

Ensure the tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as RESOLVE® Q, as well as other products used in the tank mixture.

Read and follow all applicable use directions, precautions, and limitations specified on the respective product labels, technical bulletins, and fact sheets.

RESOLVE® Q may be tank mixed with "Impact" plus atrazine for improved burndown or residual control of several broadleaf weeds including common waterhemp, common ragweed, common lambsquarters, and velvetleaf. When applying mixtures of RESOLVE® Q plus Impact, the use of methylated seed oil is recommended. Refer to Impact label for additional information regarding application timing, tank mixtures, adjuvants, and rotational crops.

RESOLVE® Q may be tank mixed with fluroxypyr herbicide (such as Starane) for improved control of kochia. Use higher labeled rates when weed infestation is heavy. Refer to the specific "Starane" label for application rates, timing and restrictions. RESOLVE® Q may be tank mixed with fluroxypyr and an additional 1/16 to 1/8 pound active ingredient dicamba (such as "Clarity") for broader spectrum weed control.

Tank Mix Compatibility Testing

Perform a jar test prior to tank mixing to ensure compatibility of RESOLVE® Q and other pesticides. Use a clear quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-ups, forms flakes, sludge, gel, oily film or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Mixing Instructions

Fertilizer Carrier Instructions

RESOLVE® Q may be mixed with water or pre-dissolved in water and added to liquid fertilizer for preemergence application. When using liquid fertilizer as the carrier, always pre-slurry RESOLVE® Q in water before adding fertilizer solutions. Add the RESOLVE® Q slurry to the final complete liquid fertilizer mixture – do not add RESOLVE® Q during the fertilizer mixing process. Always use good agitation while adding the RESOLVE® Q slurry to liquid fertilizers and maintain good agitation until sprayed. When using liquid fertilizer as the carrier, conduct a compatibility test with all components prior to mixing.

Do not use with spray additives or liquid fertilizer carriers that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 -8.0 allow for optimum stability of RESOLVE® Q.

Water Carrier Instructions

1. Fill the tank 1/3 to 1/2 full of water.
2. While agitating, add the required amount of RESOLVE® Q.
3. Continue agitation until the RESOLVE® Q is fully dispersed, at least 5 minutes. When the water temperature is 40° F or less, it is important to allow agitation and mixing to occur for the full 5 minutes to ensure the product is completely dissolved.
4. Once the RESOLVE® Q is fully dispersed, maintain agitation and continue filling tank with water. RESOLVE® Q should be thoroughly mixed with water before adding any other material such as water conditioners or other additives.
5. As the tank is filling, add tank mix partners in the proper mixing order.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. At the end of the day, or for extended periods of time between RESOLVE® Q applications, it is recommended to flush boom hoses and lines of spray solution and recharge with clean water. This will aid in proper sprayer cleanout when concluding RESOLVE® Q applications before moving on to spray other products/crops.
8. Apply RESOLVE® Q spray mixture within 48 hours of mixing to avoid product degradation.

If the selected companion herbicide has a ground or surface water advisory, consider this advisory when using the companion herbicide.

Application and Spray Volumes

Ground

Use a minimum of 15 gallons of water per acre (GPA) to ensure thorough coverage of the weeds and the best performance. Use a minimum of 10 GPA for light, scattered stands of weeds. For best performance, select nozzles and pressure that deliver MEDIUM spray droplets, as indicated, for example, by ASABE Standard S572.1. Nozzles that deliver COARSE spray droplets may be used to reduce drift, provided spray volume is increased to maintain coverage on small weeds.

Heavy crop residues may reduce burndown control of emerged weeds if residues impede spray coverage. Higher spray volumes and pressures can improve burndown control in heavy crop residue situations.

For optimal product performance and minimal spray drift, adjust the spray boom to the lowest possible spray height recommended in manufacturers' specifications. Ensure that equipment is set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl. Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

Aerial

Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at a minimum of 2 GPA.

Do not apply during a temperature inversion, when winds are gusty, or when conditions favor poor coverage and/or off-target spray movement.

Aerial application is not permitted in the State of New York.

ROTATIONAL CROP INTERVALS

The following rotational intervals must be observed when using DuPont™ RESOLVE® Q:

1.25 OZ MAXIMUM USE RATE PER ACRE PER YEAR

Rotation Crop	Interval (months)
Corn, field	Anytime
Potatoes	Anytime
Soybeans with BOLT™ technology	Anytime
Sulfonylurea tolerant soybeans	1
Cotton††	1
Tomato	1
Peanuts	1.5
Tobacco	1.5
Cereals, Winter	3
Cereals, Spring (wheat, oats, barley, rye)	9
Alfalfa*†	10
Canola†	10
Corn, pop, seed** or sweet	10
Cucumber	10
Flax	10
Peas	10
Rice	10
Red Clover†	10
Sorghum†	10
Soybeans†††	10
Snap beans, dry beans	10
Sunflower	10
Sugarbeets†	10
Sweet potatoes/yams***	10
Crops Not Listed	18

* On sprinkler irrigated fields in Idaho, Utah, and Northern Nevada it is best to use deep fall tillage such as plowing prior to planting alfalfa. Product degradation may be less on furrow irrigated soils and may result in some crop injury.

**Rotational interval to seed corn is 60 days if applying no more than 1.0 ounce per acre in the fall by December 15.

***On soils with pH 6.5 or less

† 18 months in the Red River Valley region of ND and MN. In all other areas, the rotation intervals should be extended to 18 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

††Except in Oklahoma and Texas west of Route 183, where the rotational interval is 10 months.

†††In the states of AL, AR, GA, KY, LA, MO (bootheel), MS, NC, SC, and TN the recrop interval is 30 days. In the states of KS and OK the counties containing HWY 81 and east and in MO (excluding the bootheel), IL, IN, OH, and WV the counties that contain I-70 and south and the states of DE, MD and VA, the recrop is 60 days.

GREATER THAN 1.25 OZ UP TO 2.5 OZ MAXIMUM USE RATE PER ACRE PER YEAR

Rotation Crop	Interval (months)
Corn, field	Anytime
Potatoes	Anytime
Soybeans with BOLT™ technology	Anytime
Tomato	1
Sulfonylurea tolerant soybeans	4
Cereals, Winter	3
Cereals, Spring (wheat, oats, barley, rye)	9
Corn, pop, seed or sweet	10
Cotton†	10
Cucumber	10
Flax	10
Soybeans	10
Snap beans, dry beans	10
Sunflower	10
Crops Not Listed	18

†The rotation interval should be extended to 18 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

Guidelines for Certain Areas of Oregon and Washington

Field corn grown under sprinkler irrigation with a minimum of 18 inches of water per season. This rotation interval is for sand, loamy sand and sandy loam soils having not more than 1.5% organic matter here a minimum of 18 inches of sprinkler irrigation is used on the previous corn crop. Injury to the rotated crop may occur if less than 18 inches of irrigation is used on the previous field corn crop. For tank mixtures, follow the most restrictive rotational crop guideline.

The following revised rotational intervals must be observed when using DuPont™ RESOLVE® Q on field corn:

Rotation Crop	Interval (months)
Alfalfa	4
Carrots	10
Cucumber	10
Grass, pasture, hay, seed	4
Mint	4
Onions	10
Peas	8

For Rotation to Alfalfa: RESOLVE® Q in field corn not to exceed 1.25 ounces per year in Adams, Grant, Douglas and Lincoln counties of Washington, and RESOLVE® Q in field corn not to exceed 1.88 ounces per acre per year in Benton, Franklin, Klickitat, Walla Walla and Yakima counties in Washington and Morrow and Umatilla counties in Oregon.

For Rotation to Onions and Carrots: RESOLVE® Q in field corn not to exceed 1.88 ounces per acre per year in Adams, Grant, Douglas and Lincoln counties of Washington, and RESOLVE® Q in field corn not to exceed 2.5 ounces per acre per year in Benton, Franklin, Klickitat, Walla Walla and Yakima counties in Washington and Morrow and Umatilla counties in Oregon.

For Rotation to Grass Crops Grown for Seed, Hay or Pasture: RESOLVE® Q in field corn not to exceed 1.88 ounces per acre per year in Adams, Grant, Douglas and Lincoln counties of Washington, and RESOLVE® Q in field corn not to exceed 2.5 ounces per acre per year in Benton, Franklin, Klickitat, Walla Walla and Yakima counties in Washington and Morrow and Umatilla counties in Oregon.

For Rotation to Peas and Mints: RESOLVE® Q in field corn not to exceed 1.88 ounces per acre per year in all areas.

Precaution

RESOLVE® Q should not be used in a tankmix or sequential application program with other soil residual ALS inhibiting herbicides in field corn as the combined effects of these herbicides on the planting of subsequent crops have not been thoroughly investigated and injury to the following rotation crop may occur.

SPRAYER PREPARATION/CLEANUP

It is important that spray equipment is clean and free of previous pesticide deposits before using DuPont™ RESOLVE® Q and then properly cleaned out following application. Clean all application equipment before applying RESOLVE® Q. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of RESOLVE® Q, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

- When cleaning spray equipment before applying RESOLVE® Q, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- When spraying or mixing equipment will be used over an extended period to apply multiple loads of RESOLVE® Q, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight. This will prevent the buildup of dried pesticide deposits, which can accumulate in the application equipment.

After Spraying RESOLVE® Q and Before Spraying Crops Other than Fallow or Field Corn

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of RESOLVE® Q as follows:

1. Empty the tank and drain the sump completely.
2. Spray the tank walls with clean water using a minimum volume of 10% of the tank volume. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.
3. Repeat step 2.

4. Remove the nozzles, screens and the end caps of sprayer booms and clean separately in a bucket containing water. The rinsate solution may be applied back to the crop(s) specified on this label. Do not exceed the maximum labeled use rate. If cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

Notes:

1. Always start with a clean spray tank. Ensure boom sections between end nozzles and the end of the boom are clean of deposits (It is recommended to remove end caps and visually inspect). If needed, thoroughly flush rinse water through the boom sections with the end caps removed to ensure booms are clean and free of any residue or deposits.
2. Steam-cleaning aerial spray tanks is recommended to facilitate the removal of any caked deposits.
3. When RESOLVE® Q is tank mixed with other pesticides, all cleanout procedures for each product should be examined and the most rigorous procedure should be followed.
4. Follow any pre-cleanout guidelines recommended on other product labels.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

Avoiding spray drift is the responsibility of the applicator.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply the largest droplets which are consistent with pest control objectives. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage.

Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions.

A droplet size classification system describes the range of droplet sizes produced by spray nozzles. The American Society of Agricultural and Biological Engineers (ASABE) provide a Standard that describes droplet size spectrum categories defined by a number of reference nozzles (fine, coarse, etc.). Droplet spectra resulting from the use of a specific nozzle may also be described in terms of volume mean diameter (VMD). Coarser droplet size spectra have larger VMD's and lower drift potential.

Controlling Droplet Size - Ground Application

- **Nozzle Type** - Select a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. The use of low-drift nozzles will reduce drift potential.
- **Pressure** - The lowest spray pressures recommended for the nozzle produce the largest droplets. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, using a higher-capacity nozzle instead of increasing pressure results in the coarsest droplet spectrum.
- **Flow Rate/Orifice Size** - Using the highest flow rate nozzles (largest orifice) that are consistent with pest control objectives reduces the potential for spray drift. Nozzles with higher rated flows produce coarser droplet spectra.
- **Boom Application Height** - Applications made at the lowest boom height consistent with pest control objectives, and that allow the applicator to keep the boom level with the application site and minimize bounce, will reduce the exposure of spray droplets to evaporation and wind, and reduce spray drift potential.

Controlling Droplet Size – Aircraft

- **Nozzle Type** - Solid stream, or other low drift nozzles produce the coarsest droplet spectra.
- **Number of Nozzles** - Using the minimum number of nozzles with the highest flow rate that provide uniform coverage will produce a coarser droplet spectrum.
- **Nozzle Orientation** - Orienting nozzles in a manner that minimizes the effects of air shear will produce the coarsest droplet spectra. For some nozzles such as solid stream, pointing the nozzles straight back parallel to the airstream will produce a coarser droplet spectrum than other orientations.
- **Pressure** – Selecting the pressure that produces the coarsest droplet spectrum for a particular nozzle and airspeed reduces spray drift potential. For some nozzle types such as solid streams, lower pressures can produce finer droplet spectra and increase drift potential.

Boom Length And Application Height - Aircraft

Boom Length - Using shorter booms decreases drift potential. Boom lengths are expressed as a percentage of an aircraft's wingspan or a helicopter's rotor blade diameter. Shorter boom length and proper positioning can minimize drift caused by wingtip or rotor vortices.

Application Height - Applications made at the lowest height that are consistent with pest control objectives and the safe operation of the aircraft will reduce the potential for spray drift.

Wind

Drift potential is lowest when applications are made in light to gentle sustained winds (2-10 mph), which are blowing in a constant direction. Many factors, including droplet size and equipment type also determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

Setting up equipment to produce larger droplets to compensate for droplet evaporation can reduce spray drift potential.

Droplet evaporation is most severe when conditions are both hot and dry.

Surface Temperature Inversions

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which may cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Mist or fog may indicate the presence of an inversion. If neither is present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Applications into temperature inversions are prohibited.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are minimizing drift and not interfering with uniform deposition of the product.

Air-Assisted (Air Blast) Field Crop Sprayers

Air-assisted field crop sprayers carry droplets to the target via a downward-directed airstream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application and is configured properly, and that drift potential has been minimized.

Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air assisted field crop sprayer can be used.

Sensitive Areas

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effect of spray drift.

Drift Control Additives

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution.

Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with DuPont™ RESOLVE® Q containing rimsulfuron and thifensulfuron-methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with RESOLVE® Q containing rimsulfuron and thifensulfuron-methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact DuPont at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States. This product includes ingredients that are covered under one or more of the following Bayer CropScience patents: 6,486,096; 6,569,805; 5,922,646; and 5,516,750. Purchase of this product includes a license for use only as specified on this label. Any use, mixture or formulation of this product other than as described on this label is expressly not authorized.

The DuPont Oval Logo, DuPont™, RESOLVE®, ABUNDIT®, ACCENT®, ALLUVEX™, BASIS®, BOLT™, BREAKFREE®, CINCH®, INSTIGATE®, LEADOFF®, PREQUEL®, PREVATHON®, REVULIN™ and STEADFAST® are trademarks or registered trademarks of DuPont, Pioneer or its affiliates.

"Aztec" and "Liberty" are registered trademarks of Bayer CropScience AG

"Basagran" and "Clarity" are registered trademarks of BASF

"Force" and "Warrior" are registered trademarks of Syngenta Group Company

"Counter", "Fortress", "Impact" and "SmartChoice" are registered trademarks of AmVac Chemical Corporation

"Lorsban" and "Starane" are registered trademarks of DowAgroSciences LLC

"Roundup Ready" is a registered trademark of Monsanto Technology LLC

"Asana" is a registered trademark of Sumitomo Chemical Company, Limited

SL - 1986 040516 03-16-16

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read this Limitation of Warranty and Liability before buying or using this product. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by DuPont. User assumes all risks associated with such non-directed use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT [1-888-638-7668]

Internet address: <http://cropprotection.dupont.com/>

© 2008-2016 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.

All rights reserved.