

SECTION 24(C)SPECIAL LOCAL NEED LABELING



Linex[®] 4L

Herbicide

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

EPA Reg. No. 61842-21 EPA SLN No. WA-130002

This label for LINEX[®] 4L Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024.

BROADCAST APPLICATION BEFORE EMERGENCE OF POTATOES FOR CONTROL OF REDROOT PIGWEED, COMMON LAMBSQUARTERS, HAIRY NIGHTSHADE AND BLACK NIGHTSHADE.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered label.

Target Weeds: Redroot pigweed (*Amaranthus retroflexus*), common lambsquarters (*Chenopodium album*), hairy nightshade (*Solanum sarrachoides*) and black nightshade (*Solanum nigrum*).

POTATOES (Preemergence only): Apply a single application of LINEX 4L HERBICIDE after planting but before crop emerges. Plant potato seed at least 2 inches deep. If beds are to be dragged and/or hilled, apply after the final dragging or hilling operation. Apply LINEX 4L HERBICIDE before soil cracking to avoid potential injury to potatoes. Best weed control results may be obtained when application is made to moist soil and before or as weed seedlings emerge. Water incorporation should occur prior to plant emergence. A surfactant may be added to the spray mixture when emerged broadleaf weeds are present.

Soil Texture/Type	LINEX 4L HERBICIDE Rate (pints) per Acre
Coarse: Sandy loam	1.0
Medium: Loam, Silt Loam, Silt, Sandy Clay, Sandy Clay Loam	1.5

RESTRICTIONS / PRECAUTIONS

- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.
- Use of LINEX 4L HERBICIDE is limited to potatoes grown in eastern Washington only. Do not apply this product to potatoes grown in western Washington.
- Do not apply more than one application per year.
- Do not apply to sand or loamy sand soil, as crop injury may result.
- Do not use LINEX 4L HERBICIDE on potatoes grown in soils with > 70% sand and less than 15% clay.
- Do not use on soils with less than 1% organic matter. Soils with organic matter of 1% or less, including calcareous soils, are most susceptible to linuron injury or carryover concerns.
- This product may be applied through the following irrigation systems only: linear (lateral) move, center pivot and solid set. Refer to the EPA registered label for chemigation directions.
- Do not apply by air.
- Do not apply when weather conditions favor drift from the treated area.
- Follow all plant back restrictions on the LINEX 4L HERBICIDE label.
- CROP ROTATION: Do not plant any other crop until 12 months after the last application of linuron as crop injury may result. Any crop registered for the rate of linuron that was applied may be replanted immediately.
- The degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soils low in clay or organic matter to obtain equivalent herbicide performance. Always test this product and any tank-mix combinations on a small scale for varietal responses.
- Results from treatment of emerged weeds vary with rate applied and environmental conditions. Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher.
- This pesticide is toxic to fish and aquatic invertebrates. LINEX 4L HERBICIDE should not be used under this SLN label where impact on threatened or endangered species is likely. You may refer to the WSDA Natural Resources Assessment Section web site (<https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program>) for additional information related to pesticide use and endangered species protection. Information

from the EPA about no-spray buffer zones is available at <https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program/buffers>. Pesticide applicators may use the Washington Department of Fish and Wildlife's SalmonScape mapping tool at <https://apps.wdfw.wa.gov/salmonscape> to determine if listed salmonid species occur in or near a proposed application site. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

- **Ground Water Advisory:** This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- **Surface Water Advisory:** Linuron may contaminate surface water through spray drift or under certain conditions from surface runoff into adjacent surface water bodies (ponds, lakes, streams, etc.). For several weeks post-application, linuron has a high potential to runoff when applied to fields with any of the following conditions: sloping land draining into nearby surface waters; very poorly to somewhat poorly drained soils; areas with extremely shallow ground water; frequently flooded areas; fields with surface water canals or ditches; and highly erodible land cultivated with poor management practices.

WSDA Chemigation Guidance:

- Application off-site is prohibited. The chemigation application must be continuously observed whenever sensitive areas as defined in WAC 16-202-1002(44) (including but not limited to schools, parks, dwellings, occupied buildings or structures, public roadways, and waters of the state) are at risk of being exposed to drift, runoff, or overspray. In order to minimize the potential for application off-site, WSDA recommends that the product only be applied through low pressure irrigation systems (defined as 2 to 35 pounds/square inch measured at the nozzle) with a nozzle release height no higher than 3 feet above the target crop, and that end guns be disabled throughout the application.
- An inspection port or a direct access point is required, and it must be positioned immediately upstream of the irrigation mainline check valve and be of sufficient size to allow visual and manual inspection of the check valve and low pressure drain. The inspection port or access point must have a minimum diameter of four inches, unless an alternative access system is approved by WSDA (WAC 16-202-1012[1]).
- The chemigation application tank cannot be placed within 20 feet of the wellhead or other sensitive areas. Mixing or loading activities cannot occur within 20 feet of the wellhead or other sensitive areas (WAC 16-202-1008[1]).
- WSDA Chemigation Rules (WAC 16-202-1001 through WAC 16-202-1024), are available on the Washington State Legislature website (<http://apps.leg.wa.gov/WAC/default.aspx?cite=16-202>).

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site (<https://agr.wa.gov/departments/pesticides-and-fertilizers/pesticides/waste-pesticide>). Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

For product information, call 1-800-525-2803. You may also contact 1-866-374-1975 for emergency medical treatment information.

This Section 24(c) SLN label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered label. Follow the instructions carefully.

This product when used on potatoes may lead to crop injury, loss, or damage. To the extent consistent with applicable law, Tessenderlo Kerley, Inc makes this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use LINEX 4L HERBICIDE must be made by each individual LINEX 4L HERBICIDE user and/or grower on the basis of possible crop injury from LINEX 4L HERBICIDE, the severity of weed infestation, the cost of alternative weed controls, and other factors.

Read the Warranty and Limitation of Damages on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Warranty and Limitation of Damages on the product label.

Section 24(c)

Tessenderlo Kerley, Inc.
2255 N. 44th Street, Suite 300
Phoenix, AZ 85008 USA
800-525-2803
novasource.com

Linex® and NovaSource® are registered trademarks of Tessenderlo Kerley, Inc
©2013, 2014, 2015, 2017, 2019 Tessenderlo Kerley, Inc. All Rights Reserved

Rev. 04/07/2020

Page 2 of 2

