



SPECIAL LOCAL NEED 24(C) LABELING
LINEX® 4L HERBICIDE CONTROL OF REDROOT PIGWEED,
LAMBSQUARTERS AND CERTAIN OTHER BROADLEAF
WEEDS IN POTATOES IN THE STATE OF IDAHO

Linex® 4L

Agricultural Herbicide

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO
This label for Linex® 4L expires and must not be distributed or used in accordance with
this SLN registration after December 31, 2020.

EPA Reg. No. 61842-21 EPA SLN No. ID-130001
CONTROL OF CUTLEAF REDROOT PIGWEED, LAMBSQUARTERS AND CERTAIN OTHER
BROADLEAF WEEDS IN POTATOES IN THE STATE OF IDAHO

Linuron Flowable

ACTIVE INGREDIENTS:	By Weight
Linuron 3-(3, 4-dichlorophenyl)-1-methoxy-1-methylurea40 .6%
INERT INGREDIENTS:	<u>59.4%</u>
	TOTAL: 100.0%

This product contains 4 lbs of Linuron per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when cleaning of equipment or disposing of equipment wash waters or rinsate.

Ground Water Advisory: This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: Linuron may contaminate surface water through spray drift or under certain conditions, from surface runoff into adjacent surface water bodies (ponds, lakes, streams, etc.) For several weeks post-application, linuron has a high potential to runoff when applied to fields with any of the following conditions: sloping land draining into nearby surface waters; very poorly to somewhat poorly drained soils; areas with extremely shallow ground water; frequently flooded areas; fields with surface water canals or ditches; and highly erodible land cultivated with poor management practices.

DIRECTIONS FOR USE

IMPORTANT NOTICE:

BEFORE BUYING OR USING LINEX® 4L, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THIS SUPPLEMENTAL LABEL AND THE MAIN EPA-REGISTERED LABEL.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Target Pest: Redroot Pigweed (*Amaranthus retroflexus*), Lambsquarters (*Chenopodium album*) and certain other broadleaf weeds.

Application Sites: Potatoes (*Solanum tuberosum*) fields; Apply just before or when weed and grass seedlings emerge. If emerged weeds are present, a light surfactant may be used.

Application Rate: Apply 1 to 2 pints per acre of LINEX 4L only as a preemergence treatment for redroot pigweed, lambsquarters and certain other broadleaf weeds control. As a guide, potatoes grown on calcareous or low organic matter soils should adhere to the lower end of the rate range. See restrictions.

Methods of Application: Ground application with a boom sprayer or chemigation.

Frequency of Applications: Make one application per season. Do not apply more than 2 pints broadcast per acre per year.

REI: 24 hours.

GENERAL INFORMATION

LINEX 4L is a flowable herbicide to be mixed in water and applied as a spray for selective control of weeds on certain crops. The use of LINEX 4L under this Special Local Need registration is specific to application(s) for control of redroot pigweed (*Amaranthus retroflexus*), lambsquarters and certain other broadleaf weeds in potato (*Solanum tuberosum*) plots.

LINEX 4L should be applied prior to soil cracking to help avoid potential plant injury. The degree of control and duration of effect will vary with the amount of chemical applied, soil texture, rainfall and other conditions. Soils high in clay or organic matter require higher dosages than soils low in clay or organic matter to obtain equivalent herbicide performance.

Best results are obtained when application is made to moist soil. Water incorporation should occur prior to plant emergence. Always test this product and any tank-mix combinations on a small scale for varietal responses. Follow all plant back restrictions on the LINEX 4L specimen label.

Results from treatment of emerged weeds vary with rate applied and environmental conditions. Best results are obtained on succulent weeds growing under conditions of high humidity and temperatures of 70°F or higher. Addition of a surfactant to the spray (where recommended) increases contact effects of LINEX 4L.

LINEX 4L is non-corrosive to equipment, non-flammable and non-volatile.

RESTRICTIONS

- DO NOT apply when weather conditions favor drift from the treated area.
- DO NOT apply by air.
- Organic matter of 1% or less, including calcareous soils, are most susceptible to LINEX 4L injury or carryover concerns. Use caution.

For product information, call 1-800-525-2803. You may also contact 1-866-374-1975 for emergency medical treatment information.

This label and the Federal label for this product must be in the possession of the user at the time of pesticide application.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the product label.

Section 24(c)

Tessenderlo Kerley, Inc.
2255 N. 44th Street, Suite 300
Phoenix, AZ 85008 USA
1-800-525-2803
www.novasource.com

Linex is a registered trademark of Tessenderlo Kerley, Inc.
©2013, 2018 Tessenderlo Kerley, Inc. All Rights Reserved.
01/18

