

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ PREVATHON® INSECT CONTROL

EPA Reg. No. 352-844

FOR CONTROL OF CERTAIN PESTS IN GRASS FORAGE, FODDER AND HAY (EPA CROP GROUP 17) IN THE STATES OF COLORADO, IDAHO, MONTANA, OREGON, AND WASHINGTON

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont Prevathon® insect control is recommended at 14.0 - 20.0 fluid ounces per acre (0.047 - 0.067 lb a.i. per acre) for the control of armyworms, corn earworm and sod webworm, and at 8.0 - 20.0 fluid ounces per acre (0.027 - 0.067 lb a.i. per acre) for grasshopper control. For cutworm control and suppression of billbug grubs and European crane fly larvae, apply Prevathon® at 20.0 fluid ounces/acre (0.067 lb a.i. per acre).

Crop	Application Method	Target Pest	Lb ai per acre	fluid ounces product per acre	Last Application (Days to Harvest)	REI (Hours)
Grass Forage, Fodder and Hay: (EPA Crop Group 17) Any grass, Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage	FOLIAR	Beet armyworm Corn earworm Fall armyworm Sod webworm Southern armyworm True armyworm	0.047 - 0.067	14.0 - 20.0	0	4
		Billbug (grubs)* Cutworms European crane fly (larvae)*	0.067	20.0		
		Grasshoppers	0.027 - 0.067	8.0 - 20.0		

USE RESTRICTIONS

Apply higher rates within the listed range for heavier infestations, larger/denser crops or extreme environmental conditions such as rainy weather and high temperatures.

Make no more than 4 applications per acre per calendar year.

Minimum interval between treatments is 7 days.

Do not apply more than 60 fl oz PREVATHON® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Application Instructions

For control of Armyworms, Cutworms, and Sod Webworms, apply at first sign of economic crop damage. Apply Prevathon® as a thorough coverage foliar spray using properly calibrated aerial or ground equipment that provides adequate coverage (minimum of 5 gallons per acre by air, or 10 gallons per acre by ground equipment), or via overhead chemigation in 0.10 to 0.20 acre inch of water. For foliar sprays, increase the spray volume to compensate for the amount of foliage present. For maximum spray penetration in to the root crown area, the use of a silicone surfactant may be useful. For best results with foliar spray applications, delay the next irrigation for at least 24 hours.

*For suppression of European Crane Fly larvae apply between September and early November.

*For suppression of Billbug grubs, apply when overwintered adult Billbugs are first observed. This will usually occur in late April or early May. It is important to move the Prevathon® into the grass root zone. This is best achieved by applying via overhead chemigation in 0.25 to 0.50 acre inch of water, or by immediately following a foliar spray application with 0.25 to 0.50 acre inch of water.

Grasshopper - With foliar sprays, performance is improved with the addition of a Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v). Apply when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. Applications should be made when eggs have hatched and the majority of the grasshopper population is 2nd - 3rd instar nymphs. Once grasshoppers contact and/or ingest PREVATHON® there will be rapid feeding cessation; insect mortality may not occur until a week later or longer. Do not

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2018 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

Precautions

Read and follow all instructions and precautions on the federal Section 3 label before using Prevathon® on grass grown for seed.

The preharvest interval is zero days.

The minimum re-entry interval is 4 hours.

Make no more than 4 applications per acre per calendar year.

Minimum interval between treatments is 7 days.

Do not apply more than 60 fl oz PREVATHON® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Do not make more than two sequential applications of Prevathon® before rotating to another registered insecticide having a different mode-of-action.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.