Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

GrazonNext® HL

EPA Reg. No. 62719-628

2(ee) Recommendation[†]

For Distribution and Use in the State of Oregon

Control of Susceptible Weeds Pre or Post Planting of Certain Annual and Perennial Grasses

ATTENTION

- † This recommendation is permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for GrazonNext HL before applying. Carefully follow all precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of GrazonNext HL is subject to all use precautions and limitations imposed by the label affixed to the container for GrazonNext HL.

Directions for Use

This 2(ee) recommendation is limited to the use sites listed on the main label for GrazonNext HL specialty herbicide. Refer to the main label for use sites.

GrazonNext HL controls susceptible broadleaf plants both pre and postemergence. See the GrazonNext HL main label for a list of susceptible weeds.

PRECAUTIONS

Postemergence applications on newly emerged or established sorghum-sudangrass, teff, crabgrass, and pearl millet are not recommended. Unacceptable injury or death of these grasses may occur with GrazonNext HL applications.

Pre-plant applications for control of susceptible emerged weeds

Apply GrazonNext HL at a maximum of 1.5 pts/acre with a non-ionic surfactant to control susceptible weeds no sooner than 15 days prior to planting tall fescue, orchardgrass, timothy, and annual ryegrass. GrazonNext HL can also be used at 1.5 pts/acre prior to planting sorghum-sudangrass, teff, crabgrass, and pearl millet, but a minimum 30-day interval must be observed between the application and planting.

Post planting applications for control of susceptible emerged weeds

Apply GrazonNext HL at a maximum of 1.5 pts/acre with a non-ionic surfactant to tall fescue, orchardgrass, timothy, and annual ryegrass after seedling plants have at least reached the V3 stage (three collared leaves).

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R392-028 Issued: 07/24/18 Initial printing