

OFFSET™

3.6F FUNGICIDE

GROUP 3 FUNGICIDE

FOR CONTROL OF LISTED DISEASES ON ORNAMENTALS AND GOLF COURSES

ACTIVE INGREDIENT

Tebuconazole* 38.7%

OTHER INGREDIENTS 61.3%

Total **100.0%**

Contains 3.6 pounds tebuconazole per gallon

*alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol (CAS No.: 107534-96-3)

SHAKE WELL BEFORE USING

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INSIDE BOOKLET

EPA Reg. No. 83100-1-83979

EPA Est. No. 069821-CHN-005

Manufactured for:

ROTAM NORTH AMERICA, INC.

4900 Koger Blvd., Suite #140

Greensboro, NC 27407

1-866-927-6826

Net Contents: 1 Gallon

PRODUCT OF CHINA



OFF-01-A051915-REVA052015-1G

TABLE OF CONTENTS

FIRST AID	1
PRECAUTIONARY STATEMENTS	2
PERSONAL PROTECTIVE EQUIPMENT	2
ENVIRONMENTAL HAZARDS	2
DIRECTIONS FOR USE	3
AGRICULTURAL USE REQUIREMENTS	3
NON-AGRICULTURAL USE REQUIREMENTS	3
SPRAY DRIFT MANAGEMENT	4
COMPATIBILITY TEST FOR MIX COMPONENTS/MIXING	4
RESISTANCE MANAGEMENT INFORMATION	5
DISEASE CONTROL IN GOLF COURSE TURF	5-10
DISEASE CONTROL IN CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES	11-12
STORAGE AND DISPOSAL	13
CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY	14

FIRST AID

IF SWALLOWED

- Call poison control center or doctor for treatment advice.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Have person sip a glass of water if able to swallow.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call poison control center or doctor for treatment advice.

Note to Physician

Symptoms of Poisoning and Recommendations for Medical Treatment: The compound does not cause any definite symptoms that would be diagnostic. No specific antidote. Treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Medical Emergency Treatment call your Local Emergency Response Center.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton)
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Users should: User Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Offset 3.6F Fungicide is a soluble concentrate which will control certain pests on the crops listed on this label when applied according to the Directions for Use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements on this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). See specific crop directions for applicable REI.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

APPLICATION INFORMATION FOR GOLF COURSE TURF AND ORNAMENTAL USE

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS AND ESTUARIES.

- Do not apply within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See spray Drift Management section for further information.

Chemigation

- Do not apply this product through any type of irrigation system.

SPRAY DRIFT MANAGEMENT

Make ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures. Do not make ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above ground. Mist or fog may indicate the presence of an inversion in humid areas.

The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

SPRAY VOLUME

For best results, Offset 3.6F Fungicide may be applied in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are recommended depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, equipment calibration should be checked regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

COMPATIBILITY TEST FOR MIX COMPONENTS

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1 1/2 teaspoon for each pint of liquid product of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable agent and use the compatible agent as directed on its label.

MIXING

Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that the Offset 3.6F Fungicide is completely dispersed in the mixture.

Recommended Mixing Procedure:

1. Water. Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2. Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (such as oil concentrates when applicable).
8. Water soluble additives (such as AMS or UAN when applicable).
9. Remaining quantity of water.

RESISTANCE MANAGEMENT INFORMATION

The active ingredient in Offset 3.6F Fungicide is a member of the DMI (Demethylation Inhibitor) fungicide group (FRAC grouping 3) and can exhibit cross-resistance to products with the same mode of action when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. Because the speed and scope of resistant population development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

DISEASE CONTROL IN GOLF COURSE TURF

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Offset 3.6F Fungicide is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Offset 3.6F Fungicide under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85°F.

Offset 3.6F Fungicide can be used for the prevention and control of the diseases mentioned in the table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28-day intervals as indicated. When treating golf course greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of Offset 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq. ft.) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, Offset 3.6F Fungicide should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

RESTRICTIONS:

For use on golf course turf only.

- Do not use on residential lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e. elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use clippings for animal feed.
- Do not exceed 3.6 fl. oz. of Offset 3.6F Fungicide per 1,000 sq. ft. per year.
- Do not apply more than 6 applications per year, except in New York.
- In New York State, do not apply more than 3 applications of 0.6 fl. oz. per 1,000 sq. ft., not to exceed 1.8 fl. oz. of product per 1,000 sq. ft. (2.2 lbs. Tebuconazole/acre) per year.

— GOLF COURSE TURF DISEASE CONTROL —

DISEASE	RATE of Offset 3.6F Fungicide (fl. oz./1,000 sq. ft.)		Notes
	California & New York	Other States	
<p>Dollar Spot (<i>Sclerotinia homeocarpa</i>)</p> <p>Copper Spot (<i>Gloeocercospora sorghi</i>)</p> <p>Powdery Mildew (<i>Erysiphe graminis</i>)</p> <p>Corticium Red Thread (<i>Laetisaria fuciformis</i>)</p> <p>Rusts (<i>Puccinia</i> spp.)</p> <p>Brown Patch/Rhizoctonia Blight, Large Patch (<i>Rhizoctonia solani</i>)</p> <p>Brown Ring Patch (<i>R. circinata</i>)</p> <p>Anthracnose-Basal and Foliar (<i>Colletotrichum cereale</i>)</p> <p>Red Thread (<i>Laetisaria fuciformis</i>)</p> <p>Pink Patch (<i>Limonomyces rosipellis</i>)</p>	0.6	0.6 - 1.1	<p>For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Offset 3.6F Fungicide.</p> <p>Alternate with another fungicide with a different mode of action. A second application may be made after 21 days.</p> <p>California - maintain a retreatment interval of 28 days.</p>

DISEASE	RATE of Offset 3.6F Fungicide (fl. oz./1,000 sq. ft.)		Notes
	California & New York	Other States	
Bermuda Grass decline <i>(Gaeumannomyces graminis var. graminis)</i>	0.6	0.6 - 1.1	<p>Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone.</p> <p>For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information.</p> <p>Apply subsequent applications at 21-day intervals.</p> <p>California - maintain a retreatment interval of 28 days.</p>
Take All Patch <i>(Gaemannomyces graminis)</i>	0.6	0.6 - 1.1	<p>For prevention, apply in the fall when soil temperature reaches 55-65°F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary at 21-day intervals. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.</p> <p>California - maintain a retreatment interval of 28 days.</p>

DISEASE	RATE of Offset 3.6F Fungicide (fl. oz./1,000 sq. ft.)		Notes
	California & New York	Other States	
Gray Leaf Spot <i>(Pyricularia grisea)</i>	0.6	0.6 - 1.1	Apply when conditions are favorable for disease development at 21-day intervals. If using under conditions favoring moderate to heavy disease pressure, Offset 3.6F Fungicide can be tank mixed with a registered contact fungicide at the label rate. California - maintain a retreatment interval of 28 days.
Stripe Smut <i>(Ustilago striiformis)</i>	0.6	0.6 - 1.1	Make a single application to historical disease areas in spring as grass growth begins.
Spring Dead Spot <i>(Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis)</i> Necrotic Ring Spot <i>(Leptosphaeria korrea)</i>	0.6	0.6 - 1.1	For prevention, apply in fall when soil temperature reach 75°F and again in spring under similar soil temperature conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE of Offset 3.6F Fungicide (fl. oz./1,000 sq. ft.)		Notes
	California & New York	Other States	
Fusarium Patch <i>(Fusarium roseum)</i>	0.6	0.6 - 1.1	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 21-day intervals. California - maintain a retreatment interval of 28 days.
Summer Patch <i>(Magnaporthe poae)</i>	0.6	0.6 - 1.1	Apply beginning in the spring. Do not make two consecutive applications of Offset 3.6F Fungicide. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28-day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Zoysia Patch, Large Patch of zoysia <i>(Rhizoctonia solani)</i>	0.6	0.6 - 1.1	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.

DISEASE	RATE of Offset 3.6F Fungicide (fl. oz./1,000 sq. ft.)		Notes
	California & New York	Other States	
Gray Snow Mold/Typhula Blight <i>(Typhula incarnate)</i> Pink Snow Mold/ Microdochium Patch <i>(Microdochium nivalis)</i>	0.6	0.6 - 1.1	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Offset 3.6F Fungicide be tank-mixed with other registered snow mold products for best season long results.
<p>APPLICATION: Apply the specified amount of Offset 3.6F Fungicide in 1.5 to 3.0 gallons of water per 1,000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation.</p> <p>RESTRICTIONS: Do not use clippings for animal feed. Do not exceed 3.6 fl. oz. of Offset 3.6F Fungicide per 1,000 sq. ft. per year. Do not exceed 6 applications per year except in New York. In New York state, do not apply more than 3 applications of 0.6 fl. oz. per 1,000 sq. ft., not to exceed 1.8 fl. oz. of product per 1,000 sq. ft. (2.2 lbs. Tebuconazole/acre) per year.</p>			

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

Offset 3.6F Fungicide can be used in a preventative and curative disease control program for the listed plant types and diseases in the table below. Optimum disease management is obtained when Offset 3.6F Fungicide is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full foliage coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply Offset 3.6F Fungicide at rates of 4-10 fl. oz. per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

ORNAMENTAL USE RESTRICTIONS

- For use on ornamental plants only; not for woodlands or forest management.
- Intended for use only by professional applicators.
- Do not apply more than 10 fl. oz. per acre in a single application.
- Do not apply more than 0.31 gallons (40 fl. oz.) of Offset 3.6F Fungicide (equal to 1.13 lbs. of tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

— ORNAMENTALS DISEASE CONTROL —

PLANTS	DISEASE	APPLICATION	
		TO PREVENT DISEASES	TO TREAT EXISTING DISEASE
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or tank mixing with barrier protectant fungicides is recommended for resistance management.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthracnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew	Petal Blight—Apply 2-3 times per week into the flowers as they open and develop color.	
Ground Covers and Vines	Rust Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.			

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in the original container in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store out of the reach of children, preferably in a locked storage area. Open and handle container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container (5 gallons or less). Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed, by State and local authorities, by burning. If burned, stay out of smoke.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Return to point of sale or offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ROTAM NORTH AMERICA, INC. or Seller. To the extent allowed by applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ROTAM NORTH AMERICA, INC. and Seller harmless for any claims relating to such factors.

ROTAM NORTH AMERICA, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions for use under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or **ROTAM NORTH AMERICA, INC., and to the extent allowed by applicable law, Buyer and User assume the risk of any such use. ROTAM NORTH AMERICA, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, ROTAM NORTH AMERICA, INC. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ROTAM NORTH AMERICA, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ROTAM NORTH AMERICA INC. LIMITED OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

ROTAM NORTH AMERICA, INC. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ROTAM NORTH AMERICA, INC.

Offset is a trademark of the Rotam Group.

Manufactured for:
Rotam North America, Inc.
4900 Koger Blvd.,
Suite #140
Greensboro, NC 27407