



Sevin[®] XLR PLUS

Carbaryl Insecticide

EPA Reg. No. 61842-37

FIFRA §2(ee) RECOMMENDATION

For Control of Citricola Scale

Scientific Name: Citricola Scale (*Coccus pseudomagnoliarum*) on Citrus

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE IN ARIZONA AND CALIFORNIA

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by U.S. EPA.

CITRUS FRUITS

All members of the Citrus Fruit Group 10 including: Calamondin, Citrus citron, Citrus hybrids (*Citrus* spp., includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sweet and sour), Pummelo, Satsuma mandarin, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these.

PEST	RATE OF APPLICATION – Quarts Per Acre	SPECIFIC DIRECTIONS
ARIZONA Citricola scale	5	OBSERVE BEE CAUTION (below) Apply when small, early instar nymphs (crawlers) are present.
CALIFORNIA Citricola scale	5 to 12	Apply July to September for control of small, early instar nymphs (crawlers). Monitor eggs under the adult females during June and July and apply after hatch is complete and nymphs are small, early instar stage.

RESTRICTIONS: CITRUS FRUITS

- Do not apply within 5 days of harvest.
- Do not apply more than once every 14 days.
- Do not apply more than a total of 20 quarts per acre per crop per year.

California:

- The water volume for citricola scale is generally 300-500 gallons per acre and 5 quarts per acre should be sufficient. Higher rates may be useful under conditions of severe pest pressure or with higher water volume applications.
- Restricted-entry interval (REI) = 3 days for application rates ≥ 5 quarts per acre.
- Do not apply more than 12 quarts per acre per application.

Arizona:

- Restricted-entry interval (REI) = 12 hours
- Do not apply more than 5 quarts per acre per application.

BEE CAUTION

Do not apply this product to target crop in bloom. Mowing weed or cover crop bloom prior to applying this product will reduce hazard to bees.

Read the “LIMITATION OF WARRANTY AND LIABILITY” in the label booklet for Sevin XLR.

Sevin[®] and NovaSource[®] are registered trademarks of Tesserlo Kerley, Inc. ©2016 Tesserlo Kerley, Inc. All rights reserved.

052016

TESSENDERLO KERLEY, INC., member of TESSENDERLO GROUP

P. O. Box 15627, Phoenix, Arizona 85060-5627
2255 North 44th Street, Suite 300, Phoenix, Arizona 85008-3279
Tel. (602) 889-8300, Fax (602) 889-8430

