

## RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee)  
AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

### 2(ee) Recommendation

### FOR DISTRIBUTION AND USE ONLY IN TEXAS

IT IS A VIOLATION OF FEDERAL LAW TO  
USE THIS PRODUCT IN A MANNER  
INCONSISTENT WITH ITS LABELING



EPA Reg. No. 279-3426

CROP	PEST	RATE OF APPLICATION
Section 3 Registered Crops*	Boll Weevil	4.0 Fluid Ounces/Acre (0.025 lbs active ingredient/acre)
<b>DIRECTIONS FOR USE:</b> For boll weevil control on volunteer cotton. Apply Mustang Maxx to registered crops (see section 3 label) where volunteer cotton is present. Multiple applications may be necessary to reduce pest populations to acceptable levels. Refer to each crop on the Mustang Maxx Insecticide Section 3 label to determine the maximum rate allowed per season, application intervals and other restrictions.		
<b>*RESTRICTIONS:</b>  *Refer to each crop section on the Mustang Maxx Insecticide Section 3 label for restrictions.		
<b>ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG MAXX® INSECTICIDE, EPA REG. NO. 279-3426, LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</b>		



FMC Corporation

Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Page 1 of 6

Mustang Maxx and FMC - FMC Trademarks  
©2012 FMC Corporation  
10/04/12

**Head and Stem Brassica Vegetables (1 day PHI) including: Broccoli; Chinese Broccoli** (gailon, white flowering broccoli); **Brussels Sprouts; Cauliflower; Cavalo broccolo; Kohlrabi; Cabbage; Chinese Cabbage** (napa); **Chinese Mustard Cabbage** (gai choy). **Leafy Brassica Greens (1 day PHI) including: Broccoli Raab** (rapini) ; **Chinese cabbage** (bok choy); **Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens; Turnip Greens.**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.
Do not make applications less than 7 days apart. A maximum of 24 fl oz of product (0.15 lb ai) may be applied per acre per season.		

**Bulb Vegetables (Allium spp.) (7 day PHI) including: Garlic; Garlic, Great-Headed** (elephant); **Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welch; Shallots, Dry Bulb and Green; Spring Onion or Scallions**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in a minimum of 20 gallons per acre with ground equipment or in a minimum of 3 gallons per acre by aircraft. Begin applications when pests appear and repeat as necessary to maintain control.
Do not make applications less than 7 days apart. Do not apply more than 20 fl oz of product (0.125 lb ai) per acre per season. Do not graze livestock in treated areas or cut treated crops for feed.		

**Corn (Field), Field Corn Grown for Seed, Popcorn**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not make applications less than 3 days apart. Do not apply more than 16 fl oz of product (0.1 lb ai) per acre per season including At-planting plus foliar applications of Mustang MAXX Insecticide. Do not apply within 30 days of harvest for grain and stover and 60 days for forage.		



FMC Corporation

Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Page 2 of 6

Mustang Maxx and FMC - FMC Trademarks  
©2012 FMC Corporation  
10/04/12

**Canola, Crambe, Rapeseed, Borage, Cuphea, Echium, Flax, Gold of Pleasure, Hare's-Ear Mustard, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard, Oil Radish, Poppy Seed, Sesame, and Sweet Rocket (7 Day PHI).**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not make applications less than 7 days apart. Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.		

**Cucurbit Vegetables Group** (1 day PHI) including: **Chayote (fruit); Chinese Waxgourd** (Chinese Preserving Melon); **Citron Melon; Cucumber; Gherkin; Gourd (edible)** (including hyotan, cucuzza, hechima, Chinese orkra); **Mormordica spp.** (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); **Muskmelon** (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, **cantaloupe**, casaba, crenshaw melon, golden pershaw melon, **honeydew melon**, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); **Pumpkin; Summer Squash** (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, **zucchini**); **Winter Squash** (includes butternut squash, calabaza, hubbard squash, acorn squash, and spaghetti squash); **Watermelon** (includes hybrids and varieties).

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not make applications less than 7 days apart. Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.		

**Fruiting Vegetables (except Cucurbits) (1 day PHI) including: Eggplant; groundcherry (Physalis spp.); okra; pepino (Melon pear); pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); tomatillo; tomato.**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not make applications less than 7 days apart. Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.		



FMC Corporation

Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

**Grass Forage, Fodder, and Hay Group and Grass Grown for Seed and Pasture and Rangeland (0 day PHI for forage and hay; 7 day PHI for straw and seed screenings) including:** bahiagrass, barnyardgrass, bentgrass, Bermudagrass, Kentucky bluegrass, big bluestem, smooth brome grass, buffalograss, reed canarygrass, centipedegrass, crabgrass, cupgrass, dallisgrass, sand dropseed, Kentucky fescue, meadow foxtail, eastern gramagrass, side-oats grama, guinea grass, Indian grass, Johnsongrass, lovegrass, napiergrass, oatgrass, orchardgrass, pangolagrass, paspalum, redtop, Italian ryegrass, St. Augustine grass, sprangletop, squirreltailgrass, stargrass, switchgrass, timothy, crested wheatgrass, wildrye grass and zoysia grass. Also included are sudangrass and sorghum forages and their hybrids.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment.
<p>Do not make applications less than 7 days apart for forage and hay; not less than 17 days for straw and seed screenings.</p> <p>Do not spray livestock. Allow application to dry before letting livestock graze on treated area.</p> <p>A maximum of 4 fl oz of product (0.025 lb ai) per acre may be applied per cutting.</p> <p>For hay, a maximum of 16 fl oz of product (0.1 lb ai) per acre per season may be applied. For forage, straw, and seed screenings, a maximum of 20 fl oz of product (0.125 lb ai) per acre per season may be applied.</p> <p>Applications may be made up to 0 days for forage and hay; 7 days for straw and seed screenings.</p>		

**Leafy Vegetables (except Brassica) (1 day PHI):** Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Celery, Chinese; Celtuce; Chervil; Chrysanthemum, edible-leaved and garland; Cilantro (not for use on cilantro grown for seed or coriander); Corn salad; Cress, garden; Cress, upland (yellow rocket, winter cress); Dandelion; Dock (sorrel); Endive (escarole); Fennel, Florence (finochio); Lettuce, head and leaf; Orach; Parsley; Purslane, garden; Purslane, winter; Radicchio (red chicory); Rhubarb; Spinach (including New Zealand and vine, Malabar spinach, Indian spinach); Swiss chard.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.
<p>Do not make applications less than 7 days apart.</p> <p>Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season.</p>		

**Legume Vegetables - Succulent and Dried (except Soybeans), 1 day PHI for succulent shelled or edible-podded peas or beans and 21 day PHI for dried shelled peas or beans**

**Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas (Pisum spp.) including:**

Dwarf Pea; Edible-pod Pea; Snow Pea; Sugar Snap Pea; Pigeon pea; English Pea; Garden Pea; Green Pea; Lentil.

**Succulent Edible-Podded Beans, Succulent Shelled Beans, and Dried Shelled Beans including:** Runner Bean; Snap Bean; Wax Bean; Asparagus Bean; Chinese Longbean; Moth Bean; Yardlong Bean; Jackbean; Soybean (immature seed); Swordbean; Lima Bean; Broad Bean (Fava Bean); Blackeyed Pea; Southern Pea; Grain Lupin; Sweet Lupin; White Lupin; White Sweet Lupin; Field Bean; Kidney Bean; Navy Bean; Pinto Bean; Tepary Bean; Adzuki Bean; Catjang; Cowpea; Crowder Pea; Moth Bean; Mung Bean; Rice Bean; Urd Bean; Chickpea (Garbanzo Bean); Guar; Lablab bean.

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
<p>Do not make applications less than 5 days apart.</p> <p>Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season including at-plant plus foliar applications.</p>		



FMC Corporation

Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

**Peanut (7 day PHI)**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
Do not make applications less than 14 days apart. Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season. Do not graze livestock in treated area. Do not use treated vines or hay for animal feed.		

**Root and Tuber Vegetables Group 1 (except Sugar Beet) (1 day PHI) including: Arracacha; Arrowroot; Artichoke (Chinese and Jerusalem); Garden Beet; Edible Burdock; Edible Canna; Carrot; Cassava (Bitter and Sweet); Celeriac (Celery Root); Chayote (Root); Turnip-Rooted Chervil; Chicory; Chufa; Dasheen (Taro); Ginger; Ginseng; Horseradish; Leren; Turnip-Rooted Parsley; Parsnip; Potato; Oriental Radish (Daikon); Radish; Rutabaga; Salsify (Oyster Plant); Black Salsify; Spanish Salsify; Skirret; Sweet Potato; Tanier (Cocoyam); Turmeric; Turnip; Yam Bean; and Yam (True).**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
Do not make applications less than 4 days apart. Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season. Leaves of Root and Tuber Vegetables (except Sugar Beet tops) cannot be used for food or feed.		

**Sorghum (Grain) and Millet (14 day PHI for grain and stover; 45 day PHI for forage):**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
Do not make applications less than 10 days apart. Do not apply more than 20 fl oz product (0.125 lb ai) per acre per season.		

**Soybeans (21 day PHI):**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
Do not make applications less than 7 days apart. Do not apply more than 24 fl oz of product (0.15 lb ai) per acre per season. Do not graze or harvest treated soybean forage, straw, or hay for livestock feed.		



FMC Corporation

Agriculture Products Group  
 1735 Market Street  
 Philadelphia, PA 19103

**Sugarcane (21 day phi)**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
Do not make applications less than 21 days apart. Do not apply more than 16 fl oz of product (0.1 lb ai) per acre per season.		

**Sunflower, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Stokes Aster, Tallowwood, Tea Oil Plant, and Vernonia (30 day PHI)**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not apply more than 20 fl oz of product (0.125 lb ai) per acre per season including at-plant plus foliar applications. Do not make more than five applications at the maximum application rate per season. Do not graze livestock in treated areas or cut treated crops for feed. Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.		

**Wheat and Triticale (14 day PHI for grain, forage, and hay):**

Insects Controlled	Rate of Application	Method of Application
Boll Weevil	4.0 fl oz/A (0.025 lb ai/A)	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 2 gallons per acre by air.
Do not make applications less than 14 days apart. Do not apply more than 20 fl oz of product (0.125 lb ai) per acre per season.		



FMC Corporation

Agriculture Products Group  
1735 Market Street  
Philadelphia, PA 19103

Page 6 of 6

Mustang Maxx and FMC - FMC Trademarks  
©2012 FMC Corporation  
10/04/12